

Measurement of azimuthal flow of soft and high- p_T charged particles in 5.02 TeV Pb+Pb collisions with the ATLAS detector

Klaudia Burka
on behalf of the ATLAS Collaboration



25.09.2016



Motivation

Azimuthal anisotropy

- ❖ Quark Gluon Plasma (QGP) produced and probed in heavy ion collisions
- ❖ Signatures of QGP: collective expansion, jet quenching, etc.



- ❖ Study QGP properties
 - Particle azimuthal distribution
 - Two/multi – particle correlations



Singles:
$$\frac{dN}{d\phi} \propto 1 + \sum_n 2v_n \cos[n(\phi - \Psi_n)]$$

Pairs:
$$\frac{dN}{d\Delta\phi} \propto 1 + \sum_n 2v_n^a v_n^b \cos[n(\Delta\phi)]$$

- Azimuthal anisotropy results from different pressure gradients in different spatial directions

Pb+Pb@2.76 TeV – results from Run-1


- ❖ v_n harmonics measured with Event Plane method
 - $p_T = 0.5 - 20 \text{ GeV}$
 - $|\eta| < 2.5$
 - centrality 0-80%
- ❖ Flow depends on centrality
 - Biggest asymmetry observed in mid-central events (30-50%)
- ❖ Non-zero v_n observed up to $n=6$
 - Very low viscosity of the system
- ❖ Two particle η/ϕ correlations revealed ridge and double-hump structures
- ❖ $v_{n,n}$ expansion parameters factorize to single particle ones, meaning they come from harmonic flow



Datasets

ATLAS detector → v_n measurement

ATLAS detector



- ❖ 2015: Pb+Pb 5.02 TeV, 0.49 nb^{-1}
- ❖ total luminosity sampled by minimum – bias triggers: $22 \mu \text{ b}^{-1}$

Inner Detector (Pixel+SCT)
Flow measurements is based on charged tracks reconstructed in ID


- $|\eta| < 2.5$
- $2\pi \phi$ acceptance
- $p_T > 0.5 \text{ GeV}$

Pixel detector

SCT detector

ATLAS detector → v_n measurement



ATLAS detector



- ❖ 2015: Pb+Pb 5.02 TeV, 0.49 nb^{-1}
- ❖ total luminosity sampled by minimum – bias triggers: $22 \mu \text{ b}^{-1}$

Forward Calorimeter ($3.2 < |\eta| < 4.9$)

- Centrality definition
- Flow vectors




Analysis procedures

Scalar Product (SP)

❖ **Flow vector:** $Q_n = |Q_n|e^{in\Psi_n} = \frac{1}{S} \sum_j q_{n,j} = \frac{1}{S} \sum_j w_j e^{in\phi_j}$

❖ **Flow vectors are measured in sub-events**

- FCal N and FCal P → Sum over calorimeter towers
- ID → sum over charged tracks



$$v_n\{SP\} = \frac{\langle |q_{n,j}| |Q_n^{N|P}| \cos[n(\phi_j - \Psi_n^{N|P})] \rangle}{\sqrt{\langle |Q_n^N| |Q_n^P| \cos[n(\Psi_n^N - \Psi_n^P)] \rangle}}$$

- ❖ Large eta gap ($|\eta| > 3.2$) to suppress short-range correlations
- ❖ Scalar Product: unambiguous measurement of v_n → always RMS v_n
- ❖ Event Plane: used only to compare to Run-1 results
 - Obtained by Q vectors normalization

Two – particle correlations (2PC)

- ★ Measure pair distributions in $(\Delta \eta, \Delta \phi)$

$$C(\Delta\eta, \Delta\phi) = \frac{S(\Delta\phi, \Delta\eta)}{B(\Delta\phi, \Delta\eta)}$$

signal
background



- ★ Project $S(\Delta \eta, \Delta \phi)$ and $B(\Delta \eta, \Delta \phi)$ to $\Delta \phi$ axis for $|\Delta \eta| > 2$
 - Remove short – range correlations: resonance decays, jet fragmentation etc.
- ★ Divide: $S(\Delta \phi)/B(\Delta \phi)$ to obtain correlation function $C(\Delta \phi)$

$$C(\Delta\phi) = C_0 \left(1 + \sum_{n=1}^{\infty} v_{n,n}(p_T^a, p_T^b) \cos(n\Delta\phi) \right)$$

- ★ Obtain v_n from $v_{n,n}$ using factorization:

$$v_{n,n}(p_T^a, p_T^b) = v_n(p_T^a)v_n(p_T^b)$$


$$v_n(p_T^b) = \frac{v_{n,n}(p_T^a, p_T^b)}{v_n(p_T^a)} = \frac{v_{n,n}(p_T^a, p_T^b)}{\sqrt{v_{n,n}(p_T^a, p_T^a)}}$$



Results

SP Results I: $v_n(p_T)$ dependence

central




peripheral

❖ v_n measured up to $p_T = 25$ GeV

❖ v_2 is dominant and remains positive at high p_T

❖ Flow harmonics measured for $n = 2-7$

2PC Results I: $v_n(p_T)$ dependence





★ v_n measured up to $p_T = 25$ GeV

★ v_2 is dominant and remains positive at high p_T


★ Flow harmonics measured for $n = 2-6$

SP Results II: $v_n(\eta)$ dependence





- ❖ Flow harmonics integrated over narrow p_T intervals:
 - $p_T = 0.5 - 0.6 \text{ GeV}$ and $p_T = 2 - 3 \text{ GeV}$
- ❖ All harmonics show very weak η – dependence
 - In mid-central collisions the integrated v_n over p_T range from 2 to 3 GeV is higher by about 10% in $\eta \approx 0$ compared $\eta \approx \pm 2.5$

SP Results III: v_n centrality dependence




- ❖ v_n dependence on centrality intervals
 - Integrated over narrow p_T intervals as well as the whole p_T range
- ❖ The biggest asymmetry observed in mid-central collisions (30-50 %)
 - elliptic flow is dominant asymmetry, except for the most central bin 0-5%
- ❖ Measurement of v_7 for the first time
 - v_7 is most significant for 10-40% interval

2PC – SP: Methods comparison




- ❖ Both, 2PC and SP, methods are based on two particle correlations
 - In 0-5% interval the SP method gives consistently higher values for v_2
 - For more peripheral collisions $v_2\{\text{SP}\}$ and $v_2\{\text{2PC}\}$ match within 2-5%
 - Similar trend is observed for $n > 2$

SP vs EP: Methods comparison





- ❖ EP results obtained from SP by Q-vector normalization
 - SP measure always RMS v_n
 - EP measure value between $\langle v_n \rangle$ and RMS v_n
- ❖ A small difference is seen for v_2
 - Largest for mid-central 20-50% interval $\rightarrow \sim 3\%$
- ❖ For $n > 2$ the EP and SP results are consistent

2PC Results: Run-1 – Run-2 comparison



- ❖ 2PC shows overall good agreement between results obtained with different system energies
 - Within statistical and systematic uncertainties
- ❖ Consistent with recent ALICE results ([PRL 116 \(2016\) 132302](#))

EP results: Run-1 – Run-2 comparison



- ❖ EP methods shows overall good agreement between results obtained with different system energies
 - Within statistical and systematic uncertainties
- ❖ Consistent with recent ALICE results ([PRL 116 \(2016\) 132302](#))

Summary

- ❖ The first ATLAS measurement of azimuthal anisotropy of charged particles in Pb+Pb collisions at 5.02 TeV using LHC Run-2 data
- ❖ The flow harmonics, v_n , are measured using two methods, SP and 2PC:
 - $n=2-7$
 - wide $p_T = 0.5 - 25$ GeV range
 - $| \eta | < 2.5$
 - centrality 0-80%
- ❖ The first measurement of v_7 harmonic
- ❖ Significant v_2 even at highest p_T
- ❖ The v_n show weak η - dependence
- ❖ The $v_n(p_T)$ values do not change from 2.76 TeV to 5.02 TeV

Backup slides

SP/EP systematics summary table

systematic sources	n harmonic	5 - 10 %		50 - 60 %	
		0.5 - 0.6 GeV	9 - 10 GeV	0.5 - 0.6 GeV	9 - 10 GeV
tracking cuts	v_2	5 (5)	0.2 (0.3)	0.1 (0.1)	0.3 (0.3)
	v_3	6 (6)	0.2 (0.2)	0.2 (0.1)	3 (2)
	v_4	6 (6)	0.4 (0.2)	3 (3)	1 (3)
	v_5	7 (9)	0.2 (1)	2 (2)	3 (2)
	v_6	14 (17)	1 (3)	3 (6)	3 (6)
	v_7	2 (12)	9 (3)	6 (26)	6 (26)
	v_2	0.2 (0.2)	<0.1 (<0.1)	0.2 (0.2)	<0.1 (<0.1)
efficiency variation	v_3	0.2 (0.2)	0.2 (<0.1)	0.3 (0.3)	0.7 (0.5)
	v_4	0.3 (0.3)	0.2 (0.3)	0.3 (0.2)	0.7 (0.5)
	v_5	0.2 (0.2)	<0.1 (0.2)	0.2 (0.2)	1 (3)
	v_6	5 (17)	11 (2)	5 (6)	0.9 (2)
	v_7	3 (3)	0.1 (0.4)	2 (4)	2 (2)
	v_2	0.8 (0.7)	<0.1 (<0.1)	0.2 (0.1)	0.3 (<0.1)
η symmetry	v_3	1 (1)	0.5 (0.3)	0.6 (0.5)	1 (0.5)
	v_4	1 (1)	0.4 (0.9)	2 (5)	4 (9)
	v_5	2 (2)	3 (5)	4 (4)	3 (3)
	v_6	10 (7)	4 (4)	11 (7)	11 (7)
	v_7	11 (15)	11 (15)	15 (12)	
centrality	v_2	1 (1)	1 (1)	0.5 (0.3)	1 (1)
	v_3	0.2 (0.2)	0.2 (<0.1)	0.3 (0.3)	0.7 (0.5)
	v_4	<0.1 (<0.1)	0.4 (0.7)	1 (3)	0.8 (3)
	v_5	2 (2)	0.2 (0.5)	4 (4)	2 (1)
	v_6	2 (1)	2 (2)	2 (3)	2 (3)
	v_7	11 (7)	8 (7)	4 (4)	4 (4)
residual sine term	v_2	0.2 (0.2)	0.1 (<0.1)	0.4 (0.5)	0.3 (0.5)
	v_3	0.5 (0.5)	1 (1)	2 (2)	1 (0.4)
	v_4	1 (2)	0.7 (1)	0.2 (3)	6 (4)
	v_5	3 (4)	0.1 (3)	11 (13)	11 (4)
	v_6	3 (11)	17 (21)	21 (31)	21 (31)
	v_7	34 (26)		35 (43)	
	v_2	2 (2)	1 (1)	0.3 (<0.1)	1 (1)
MC closure	v_3	2 (3)	2 (1)	14 (14)	11 (11)
	v_4	4 (4)	0.5 (1)	1 (3)	5 (9)
	v_5	3 (7)	14 (21)	8 (7)	2 (3)
	v_6	-	-	-	-
	v_7	-	-	-	-
	v_2	0.1 (0.4)	0.7 (1)	0.1 (<0.1)	2 (0.6)
	v_3	1 (2)	2 (2)	0.3 (2)	8 (10)
residual FCal mis-calibration	v_4	2 (3)	4 (6)	3 (2)	0.1 (6)
	v_5	8 (6)	<0.1 (4)	5 (8)	2 (3)
	v_6	17 (5)	5 (17)	28 (3)	28 (3)
	v_7	34 (13)	34 (13)	34 (13)	34 (13)

2PC systematics summary table

systematic sources	n harmonic	5 - 10 %		50 - 60 %	
		0.5–0.6 GeV	6–8 GeV	0.5–0.6 GeV	6–8 GeV
tracking cuts	v_2	8	3	1	1
	v_3	8	3	1	2
	v_4	11	4	3	4
	v_5	16	5	4	5
	v_6	16	8	4	8
efficiency variation	v_2	0.2	<0.1	0.2	<0.1
	v_3	0.2	0.2	0.3	0.7
	v_4	0.3	0.2	0.3	0.7
	v_5	0.2	<0.1	0.2	1.0
	v_6	4.8	11	4.2	0.9
centrality	v_2	1	1	1.5	<0.5
	v_3	0.5	0.5	3	10
	v_4	0.5	0.5	3	10
	v_5	0.5	0.5	3	10
	v_6	0.5	0.5	3	10
MC closure	v_2	6	3	3	1
	v_3	6	3	3	1
	v_4	5	5	5	5
	v_5	6	6	6	6
	v_6	10	10	10	10
event-mixing	v_2	1	1	1	1
	v_3	1	2	1	4
	v_4	5	6	3	6
	v_5	5	10	5	10
	v_6	50	15	50	15

SP/EP methods

Scalar Product:

$$v_{n,j}\{SP\} = Re \frac{\langle q_{n,j} Q_n^{N|P*} \rangle}{\sqrt{\langle Q_n^N Q_n^{P*} \rangle}} = \frac{\langle |q_{n,j}| |Q_n^{N|P}| \cos[n(\phi_j - \Psi_n^{N|P})] \rangle}{\sqrt{\langle |Q_n^N| |Q_n^P| \cos[n(\Psi_n^N - \Psi_n^P)] \rangle}}$$

Event Plane:

$$v_{n,j}\{EP\} = Re \frac{\langle q_{n,j} \frac{Q_n^{N|P*}}{|Q_n^{N|P}|} \rangle}{\sqrt{\langle \frac{Q_n^N}{|Q_n^N|} \frac{Q_n^{P*}}{|Q_n^P|} \rangle}} = \frac{\langle \cos[n(\phi_j - \Psi_n^{N|P})] \rangle}{\sqrt{\langle \cos[n(\Psi_n^N - \Psi_n^P)] \rangle}}$$