

# Spectroscopy and Structure of Baryons

Willibald Plessas

Theoretical Physics / Institute of Physics  
University of Graz, Austria

Humboldt Kolleg

"From the Vacuum to the Universe"

Kitzbühel, Austria, June 27<sup>th</sup>, 2016

Low-energy  
QCD

RCQM

GBE RCQM

Spectroscopy

Light, strange,  
charm, bottom

Structure

Nucleon E.m.

Baryon E.m.

Axial FFs

$\pi NN$ ,  $\pi N\Delta$

$N$  Scalar FF

$N$  Gravitational FF

Summary

# Hadrons at Low Energies

Common methods for dealing with low-energy hadronic phenomena on the basis of QCD nowadays:

- ▶ Lattice QCD (works but faces computational limitations)
- ▶ Chiral perturbation theory (works at low energies, as a systematic expansion but is limited to a few terms)
- ▶ Effective field theories or functional methods (depend on assumptions and regularizations)
- ▶ Effective models, e.g. **constituent-quark models** (depend on assumptions and input parameters)

Here, results for the **spectroscopy** and **structure** (electromagnetic, weak, scalar, gravitational form factors) of **baryons** from the **relativistic constituent-quark model** (RCQM)

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom

Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary

# Aspects Suggestive for Quark Models

Search for a non-perturbative tool to describe/understand

- in a consistent manner
- on the microscopic level
- in accordance with the properties of low-energy QCD such phenomena like

▶ **hadron spectra:** ground states & excitations

▶ **hadron structure:**  $r_E, \mu, g_A; G_E, G_M, G_A, G_P, \dots$   
i.e. electroweak form factors etc.

▶ **resonance excitations:**  $\gamma N \rightarrow N^*, e^- N \rightarrow N^*, \dots$

▶ **resonance decays:**

$\rho \rightarrow \pi\pi, \omega \rightarrow \pi\pi\pi, N^* \rightarrow N\pi, \Delta \rightarrow N\pi, \Lambda^* \rightarrow KN, \dots$

▶ **meson-baryon interactions:**  $\pi - N, K - N, \dots$

▶ **hyperon-hyperon interactions:**  $N - N, N - Y, \dots$   
etc. etc.

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom

Structure  
Nucleon E.m.  
Baryon E.m.

Axial FFs  
 $\pi NN, \pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom

Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary

Low-Energy QCD / Relevant Degrees of Freedom

Relativistic Constituent-Quark Model (RCQM)

Interacting mass operator with  
Goldstone-boson-exchange dynamics

Baryon Spectroscopy

Light, strange, charm, bottom

Baryon Structure

Nucleon e.m. form factors, including flavor analysis

Baryon electromagnetic form factors

Nucleon and baryon axial form factors / charges

Strong  $\pi NN$  and  $\pi N\Delta$  form factors

Nucleon scalar form factor & sigma term

Nucleon gravitational form factors

Summary and Conclusions

Low-energy QCD of  $N_f$  flavors is characterized by:

- spontaneous breaking of chiral symmetry ( $SB_{\chi}S$ ):

$$SU(N_f)_L \times SU(N_f)_R \rightarrow SU(N_f)_V$$

→ appearance of  $(N_f^2 - 1)$  **Goldstone bosons**  $\vec{\phi}$

→ generation of quasiparticles with dynamical mass,  
i.e. **constituent quarks**  $\psi$

- thus (effective) interaction Lagrangian:

$$\mathcal{L}_{\text{int}} \sim ig\bar{\psi}\gamma_5\vec{\lambda}^f \cdot \vec{\phi}\psi$$

A. Manohar and H. Georgi: Nucl. Phys. B 234 (1984) 189

E.V. Shuryak: Phys. Rep. **115**, 151 (1984)

L.Ya. Glozman and D.O. Riska: Phys. Rep. **268**, 263 (1996)


see also:

S. Weinberg: Phys. Rev. Lett. **105**, 261601 (2010)

# Baryons

Baryons are considered as colorless bound states of three constituent quarks.

Here the proton:



- ▶ 'Constituent' quarks are quasiparticles with **dynamical mass**, NOT the original QCD d.o.f. (i.e. 'current' quarks).
- ▶ 'Constituent' quarks are confined and interact via hyperfine interactions associated with  $SB_{\chi}S$ , i.e. **Goldstone-boson exchange**.

## Relativistic quantum mechanics (RQM)

i.e. **Hamiltonian quantum theory** respecting

### **Poincaré invariance**

(theory on a Hilbert space  $\mathcal{H}$  corresponding to a finite number of particles, not a field theory)

### **Invariant mass operator**

$$\hat{M} = \hat{M}_{free} + \hat{M}_{int}$$

### **Eigenvalue equations**

$$\hat{M} |P, J, \Sigma\rangle = M |P, J, \Sigma\rangle \quad , \quad \hat{M}^2 = \hat{P}^\mu \hat{P}_\mu$$

$$\hat{P}^\mu |P, J, \Sigma\rangle = P^\mu |P, J, \Sigma\rangle \quad , \quad \hat{P}^\mu = \hat{M} \hat{V}^\mu$$

## Interacting mass operator

$$\hat{M} = \hat{M}_{free} + \hat{M}_{int}$$

$$\hat{M}_{free} = \sqrt{\hat{H}_{free}^2 - \hat{\vec{P}}_{free}^2}$$

$$\hat{M}_{int}^{rest\ frame} = \sum_{i < j}^3 \hat{V}_{ij} = \sum_{i < j}^3 [\hat{V}_{ij}^{conf} + \hat{V}_{ij}^{hf}]$$

fulfilling the **Poincaré algebra**

$$\begin{aligned} [\hat{P}_i, \hat{P}_j] &= 0, & [\hat{J}_i, \hat{H}] &= 0, & [\hat{P}_i, \hat{H}] &= 0, \\ [\hat{K}_i, \hat{H}] &= -i\hat{P}_i, & [\hat{J}_i, \hat{J}_j] &= i\epsilon_{ijk}\hat{J}_k, & [\hat{J}_i, \hat{K}_j] &= i\epsilon_{ijk}\hat{K}_k, \\ [\hat{J}_i, \hat{P}_j] &= i\epsilon_{ijk}\hat{P}_k, & [\hat{K}_i, \hat{K}_j] &= -i\epsilon_{ijk}\hat{J}_k, & [\hat{K}_i, \hat{P}_j] &= -i\delta_{ij}\hat{H} \end{aligned}$$

$\hat{H}, \hat{P}_i$  ... time and space translations,

$\hat{J}_i$  ... rotations,  $\hat{K}_i$  ... Lorentz boosts



Phenomenologically, baryons with 5 flavors:  $u, d, s, c, b$

$$\Rightarrow H_{free} = \sum_{i=1}^3 \sqrt{m_i^2 + \vec{k}_i^2}$$

$$V^{conf}(\vec{r}_{ij}) = B + C r_{ij}$$

$$V^{hf}(\vec{r}_{ij}) = \left[ V_{24}(\vec{r}_{ij}) \sum_{f=1}^{24} \lambda_i^f \lambda_j^f + V_0(\vec{r}_{ij}) \lambda_i^0 \lambda_j^0 \right] \vec{\sigma}_i \cdot \vec{\sigma}_j$$

- ▶ i.e., for  $N_f = 5$ , we have the exchange of a **24-plet** plus a **singlet** of Goldstone bosons.

L.Ya. Glozman, W. Plessas, K. Varga, and R.F. Wagenbrunn: Phys. Rev. D **58**, 094030 (1998)

J.P. Day, K.-S. Choi, and W. Plessas: arXiv:1205.6918

J.P. Day, K.-S. Choi, and W. Plessas: Few-Body Syst. **54**, 329 (2013)



$$V^{conf}(\vec{r}_{ij}) = B + C r_{ij}$$

$$V_{\beta}(\vec{r}_{ij}) = \frac{g_{\beta}^2}{4\pi} \frac{1}{12m_i m_j} \left\{ \mu_{\beta}^2 \frac{e^{-\mu_{\beta} r_{ij}}}{r_{ij}} - 4\pi \delta(\vec{r}_{ij}) \right\}$$

$$= \frac{g_{\beta}^2}{4\pi} \frac{1}{12m_i m_j} \left\{ \mu_{\beta}^2 \frac{e^{-\mu_{\beta} r_{ij}}}{r_{ij}} - \Lambda_{\beta}^2 \frac{e^{-\Lambda_{\beta} r_{ij}}}{r_{ij}} \right\}$$

$$B = -402 \text{ MeV}, \quad C = 2.33 \text{ fm}^{-2}$$

$$\beta = 24 : \quad \frac{g_{24}^2}{4\pi} = 0.7, \quad \mu_{24} = \mu_{\pi} = 139 \text{ MeV}, \quad \Lambda_{24} = 700.5 \text{ MeV}$$

$$\beta = 0 : \quad \left( \frac{g_0}{g_{24}} \right)^2 = 1.5, \quad \mu_0 = \mu_{\eta'} = 958 \text{ MeV}, \quad \Lambda_0 = 1484 \text{ MeV}$$

$$m_u = m_d = 340 \text{ MeV}, \quad m_s = 480 \text{ MeV},$$

$$m_c = 1675 \text{ MeV}, \quad m_b = 5055 \text{ MeV}$$

Low-energy  
QCD

RCQM

GBE RCQM

Spectroscopy

Light, strange,  
charm, bottom

Structure

Nucleon E.m.

Baryon E.m.

Axial FFs

$\pi NN$ ,  $\pi N\Delta$

$N$  Scalar FF

$N$  Gravitational FF


Summary

Baryon **Excitation Spectra**

and

Mass-Operator **Eigenstates**

# Light Baryon Spectra



red Universal GBE RCQM

green PDG 2013 (experiment)

- Low-energy QCD
- RCQM
  - GBE RCQM
- Spectroscopy
  - Light, strange, charm, bottom
- Structure
  - Nucleon E.m.
  - Baryon E.m.
  - Axial FFs
  - $\pi NN$ ,  $\pi N\Delta$
  - $N$  Scalar FF
  - $N$  Gravitational FF
- Summary



# Strange Baryon Spectra

Low-energy QCD

RCQM

GBE RCQM

Spectroscopy

Light, strange, charm, bottom

Structure

Nucleon E.m.

Baryon E.m.


Axial FFs

$\pi NN$ ,  $\pi N\Delta$

$N$  Scalar FF

$N$  Gravitational FF


Summary



red Universal GBE RCQM

green PDG 2013 (experiment)

# Charm Baryon Spectra



## Left panel – single charm:

**red** Universal GBE RCQM prediction

**green** PDG 2013 (experiment)



## Right panel – double charm:

**green** M. Mattson et al.: Phys. Rev. Lett. 89 (2002) 112001 (SELEX experiment)

**cyan** S. Migura, D. Merten, B. Metsch, and H.-R. Petry: Eur. Phys. J. A 28 (2006) 41 (Bonn RCQM)

**magenta** L. Liu et al.: Phys. Rev. D 81 (2010) 094505 (Lattice QCD)

# Bottom Baryon Spectra



## Left panel – single bottom:

red Universal GBE RCQM prediction

green PDG 2013 (experiment)

## Right panel – double bottom:

green W. Roberts and M. Pervin: Int. J. Mod. Phys. A 23 (2008) 2817 (nonrel. one-gluon-exchange CQM)

orange D. Ebert, R.N. Faustov, V.O. Galkin, and A.P. Martynenko: Phys. Rev. D 66 (2002) 014008 (RCQM)

Low-energy QCD

RCQM

GBE RCQM

Spectroscopy

Light, strange, charm, bottom

Structure

Nucleon E.m.

Baryon E.m.

Axial FFs

$\pi NN$ ,  $\pi N\Delta$

$N$  Scalar FF

$N$  Gravitational FF

Summary

Low-energy QCD

RCQM

GBE RCQM

Spectroscopy

Light, strange, charm, bottom

Structure

Nucleon E.m.

Baryon E.m.


Axial FFs

$\pi NN$ ,  $\pi N\Delta$

$N$  Scalar FF

$N$  Gravitational FF

Summary



**red** Universal GBE RCQM

**green** W. Roberts and M. Pervin: Int. J. Mod. Phys. A 23 (2008) 2817  
(nonrelativistic one-gluon-exchange CQM)

**blue** S. Migura, D. Merten, B. Metsch, and H.-R. Petry: Eur. Phys. J. A 28 (2006) 41 (Bonn RCQM)

**cyan** A.P. Martyntenko: Phys. Lett. B 663 (2008) 317 (RCQM)


**magenta** S. Meinel: Phys. Rev. D 82 (2010) 114502 (lattice QCD)



# Influence of Light-Heavy Q-Q Interaction



- Low-energy QCD
- RCQM
  - GBE RCQM
- Spectroscopy
  - Light, strange, charm, bottom
- Structure
  - Nucleon E.m.
  - Baryon E.m.
  - Axial FFs
  - $\pi NN$ ,  $\pi N\Delta$
  - $N$  Scalar FF
  - $N$  Gravitational FF
- Summary



leftmost cyan levels

confinement only

middle magenta levels

including only light-light GBE

rightmost red levels

including full GBE RCQM

## Mass operator eigenstates

$$\hat{M} |P, J, \Sigma, T, M_T\rangle = M |P, J, \Sigma, T, M_T\rangle$$

represented in configuration space

$$\langle \vec{\xi}, \vec{\eta} | P, J, \Sigma, T, M_T \rangle = \Psi_{PJ\Sigma TM_T}(\vec{\xi}, \vec{\eta})$$

with  $\vec{\xi}$  and  $\vec{\eta}$  the usual Jacobi coordinates.

Picture the baryon wave functions through

## spatial probability density distributions

$$\rho(\xi, \eta) = \xi^2 \eta^2 \int d\Omega_\xi d\Omega_\eta \Psi_{PJ\Sigma TM_T}^*(\xi, \Omega_\xi, \eta, \Omega_\eta) \Psi_{PJ\Sigma TM_T}(\xi, \Omega_\xi, \eta, \Omega_\eta)$$

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom



Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary



# Pictures of Baryons (rest frame)



N GBE CQM




N(1440) GBE CQM




- Low-energy QCD
- RCQM
- GBE RCQM
- Spectroscopy
- Light, strange, charm, bottom
- Structure
- Nucleon E.m.
- Baryon E.m.
- Axial FFs
- $\pi NN, \pi N\Delta$
- N Scalar FF
- N Gravitational FF
- Summary

# Spatial Probability Density Distributions

$\rho(\xi, \eta)$  for the  $\frac{1}{2}^+$  octet baryon ground states  $N(939)$ ,  $\Lambda(1116)$ ,  $\Sigma(1193)$ ,  $\Xi(1318)$ :




$\rho(\xi, \eta)$  for the  $\frac{1}{2}^+$  octet baryon states  $N(1440)$ ,  $\Lambda(1600)$ ,  $\Sigma(1660)$ ,  $\Xi(1690)$ :




T. Melde, W. Plessas, and B. Sengl: Phys. Rev. D **77** (2008) 114002

# Spatial Probability Density Distributions

$\rho(\xi, \eta)$  for the  $\frac{3}{2}^+$  decuplet baryon states  $\Delta(1232)$ ,  $\Sigma(1385)$ ,  $\Xi(1530)$ ,  $\Omega(1672)$ :



$\rho(\xi, \eta)$  for the  $\frac{3}{2}^+$  decuplet baryon states  $\Delta(1600)$ ,  $\Sigma(1690)$ :



multiplet	$(LS)J^P$				
octet	$(0 \frac{1}{2})_1^+$	$N(939)^{100}$	$\Lambda(1116)^{100}$	$\Sigma(1193)^{100}$	$\Xi(1318)^{100}$
octet	$(0 \frac{1}{2})_2^+$	$N(1440)^{100}$	$\Lambda(1600)^{96}$	$\Sigma(1660)^{100}$	$\Xi(1690)^{100}$
octet	$(0 \frac{1}{2})_3^+$	$N(1710)^{100}$		$\Sigma(1880)^{99}$	
octet	$(1 \frac{1}{2})_1^-$	$N(1535)^{100}$	$\Lambda(1670)^{72}$	$\Sigma(1560)^{94}$	
octet	$(1 \frac{1}{2})_2^-$	$N(1650)^{100}$	$\Lambda(1800)^{100}$	$\Sigma(1620)^{100}$	
octet	$(1 \frac{1}{2})_3^-$	$N(1520)^{100}$	$\Lambda(1690)^{72}$	$\Sigma(1670)^{94}$	$\Xi(1820)^{97}$
octet	$(1 \frac{1}{2})_4^-$	$N(1700)^{100}$		$\Sigma(1940)^{100}$	
octet	$(1 \frac{1}{2})_5^-$	$N(1675)^{100}$	$\Lambda(1830)^{100}$	$\Sigma(1775)^{100}$	$\Xi(1950)^{100}$
decuplet	$(0 \frac{3}{2})_3^+$	$\Delta(1232)^{100}$	$\Sigma(1385)^{100}$	$\Xi(1530)^{100}$	$\Omega(1672)^{100}$
decuplet	$(0 \frac{3}{2})_4^+$	$\Delta(1600)^{100}$	$\Sigma(1690)^{99}$		
decuplet	$(1 \frac{3}{2})_1^-$	$\Delta(1620)^{100}$	$\Sigma(1750)^{94}$		
decuplet	$(1 \frac{3}{2})_2^-$	$\Delta(1700)^{100}$			
singlet	$(1 \frac{1}{2})_1^-$	$\Lambda(1405)^{71}$			
singlet	$(1 \frac{1}{2})_2^-$	$\Lambda(1520)^{71}$			
singlet	$(0 \frac{1}{2})_2^+$	$\Lambda(1810)^{92}$			

T. Melde, W. Plessas, and B. Sengl: Phys. Rev. D **77**, 114002 (2008)

See also the PDG: Chin. Phys. C **38**, 090001 (2014)

# SU(3) Flavor Multiplets – New

## Classification of baryon resonances by the PDG since 2010

(results from the GBE relativistic CQM marked by asterisks)

$J^P$	$(D, L_N^P) S$	Octet members			Singlets
$1/2^+$	$(56, 0_0^+)$	$1/2 N(939)$	$\Lambda(1116)$	$\Sigma(1193)$	$\Xi(1318)$
$1/2^+$	$(56, 0_2^+)$	$1/2 N(1440)$	$\Lambda(1600)$	$\Sigma(1660)$	$\Xi(1690)^\dagger$
$1/2^-$	$(70, 1_1^-)$	$1/2 N(1535)$	$\Lambda(1670)$	$\Sigma(1620)$	$\Xi(?)$ $\Lambda(1405)$
				$\Sigma(1560)^\dagger$	
$3/2^-$	$(70, 1_1^-)$	$1/2 N(1520)$	$\Lambda(1690)$	$\Sigma(1670)$	$\Xi(1820)$ $\Lambda(1520)$
$1/2^-$	$(70, 1_1^-)$	$3/2 N(1650)$	$\Lambda(1800)$	$\Sigma(1750)$	$\Xi(?)$
				$\Sigma(1620)^\dagger$	
$3/2^-$	$(70, 1_1^-)$	$3/2 N(1700)$	$\Lambda(?)$	$\Sigma(1940)^\dagger$	$\Xi(?)$
$5/2^-$	$(70, 1_1^-)$	$3/2 N(1675)$	$\Lambda(1830)$	$\Sigma(1775)$	$\Xi(1950)^\dagger$
$1/2^+$	$(70, 0_2^+)$	$1/2 N(1710)$	$\Lambda(1810)$	$\Sigma(1880)$	$\Xi(?)$ $\Lambda(1810)^\dagger$
$3/2^+$	$(56, 2_2^+)$	$1/2 N(1720)$	$\Lambda(1890)$	$\Sigma(?)$	$\Xi(?)$
$5/2^+$	$(56, 2_2^+)$	$1/2 N(1680)$	$\Lambda(1820)$	$\Sigma(1915)$	$\Xi(2030)$
$7/2^-$	$(70, 3_3^-)$	$1/2 N(2190)$	$\Lambda(?)$	$\Sigma(?)$	$\Xi(?)$ $\Lambda(2100)$
$9/2^-$	$(70, 3_3^-)$	$3/2 N(2250)$	$\Lambda(?)$	$\Sigma(?)$	$\Xi(?)$
$9/2^+$	$(56, 4_4^+)$	$1/2 N(2220)$	$\Lambda(2350)$	$\Sigma(?)$	$\Xi(?)$

PDG: J. Phys. G **37**, 075021 (2010); Phys. Rev. D **86**, 010001 (2012);



Chin. Phys. C **38**, 090001 (2014)



# $N$ and $\Delta$ Rest-Frame Wave Functions

Rest-frame **spatial distribution of constituent quarks**  
in terms of 3-body Jacobi coordinates  $\vec{\xi}$  and  $\vec{\eta}$ :

$$\rho(\xi, \eta) = \xi^2 \eta^2 \int d\Omega_\xi d\Omega_\eta \Psi_{PJ\Sigma TM_T}^*(\xi, \Omega_\xi, \eta, \Omega_\eta) \Psi_{PJ\Sigma TM_T}(\xi, \Omega_\xi, \eta, \Omega_\eta)$$



Units on abscissa and ordinates are [fm]




The **root-mean-square radius** (in the rest frame):


$$r_{\text{rms}} = \sqrt{\langle r_i^2 \rangle} = \left( \int d^3 r_i \langle P = 0, J, \Sigma | \hat{r}_i^2 | P = 0, J, \Sigma \rangle \right)^{1/2}$$

Is NOT an **observable**! Is NOT **relativistically invariant**!

→ Idea about the **spatial distribution** of constituent quarks.



$$r_{\text{rms}}^N = 0.304 \text{ fm}$$



$$r_{\text{rms}}^\Delta = 0.390 \text{ fm}$$


# Root-Mean-Square Radii

The **root-mean-square radius** (in the rest frame):


$$r_{\text{rms}} = \sqrt{\langle r_i^2 \rangle} = \left( \int d^3 r_i \langle P = 0, J, \Sigma | \hat{r}_i^2 | P = 0, J, \Sigma \rangle \right)^{\frac{1}{2}}$$

Is NOT an **observable**! Is NOT **relativistically invariant**!

→ Idea about the **spatial distribution** of constituent quarks.



$$r_{\text{rms}}^N = 0.304 \text{ fm}$$



$$r_{\text{rms}}^\Delta = 0.390 \text{ fm}$$

Exp.:  $r_E^p \sim 0.88 \text{ fm}$   
 $(r_E^n)^2 \sim -0.12 \text{ fm}^2$

$r_E^{\Delta^{++}} = r_E^{\Delta^+} = r_E^{\Delta^-} = 0.656 \text{ fm}$   
 $r_E^{\Delta^0} = 0 \text{ fm}$

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom

Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary

Baryon **Reactions**

&

Baryon **Structure**

# Various Baryon Reactions

Matrix elements of a transition operator  $\hat{O}$  between baryon eigenstates  $|P, J, \Sigma, T, T_3, Y\rangle$

$$\langle P', J', \Sigma', T', T'_3, Y' | \hat{O} | P, J, \Sigma, T, T_3, Y \rangle$$

- $\hat{O} \dots \hat{J}_{\text{em}}^\mu \rightarrow$  electromagnetic FF's
- $\dots \hat{A}_{\text{axial}}^\mu \rightarrow$  axial FF's
- $\dots \hat{\Theta}^{\mu\nu} \rightarrow$  gravitational/tensor FF's
- $\dots \hat{S} \rightarrow$  scalar FF
- $\dots \hat{D}_\lambda^\mu \rightarrow$  strong FF's and hadronic decays

To be calculated from microscopic three-quark ME's

$$\langle p'_1, p'_2, p'_3; \sigma'_1, \sigma'_2, \sigma'_3; f'_{i'_1}, f'_{i'_2}, f'_{i'_3} | \hat{O} | p_1, p_2, p_3; \sigma_1, \sigma_2, \sigma_3; f_{i_1}, f_{i_2}, f_{i_3} \rangle$$

$\uparrow$   
boosted 3-body states


$\uparrow$   
boosted 3-body states

# Rel. Dynamics: Poincaré Transf. Generators

Invariant hypersurfaces in Minkowski space  $\rightsquigarrow$  3 forms

P.A.M. Dirac: Rev. Mod. Phys. 21 (1949) 392

instant




$$x^0 = 0$$

$J_i, P_j$

$$P^0 = H, K_1, K_2, K_3$$

point




$$x^2 - a^2 = 0$$

$J_i, K_j$

$$P^0, P^1, P^2, P^3$$

front



$$x^0 - x^3 = 0$$

$P^+, \vec{P}_\perp, E^1, E^2, J_z, K_z$

$$P^-, F^1, F^2$$

interaction-free and interaction-dependent generators.

There are 2 and only 2 more forms discussed by Leutwyler & Stern with less, namely only **three** interaction-free generators!


H. Leutwyler and J. Stern: Ann. Phys. 112 (1978) 94

Covariant predictions for:

- ▶ **Electromagnetic** nucleon form factors  
 $G_E^p(Q^2)$ ,  $G_M^p(Q^2)$ ;  $G_E^n(Q^2)$ ,  $G_M^n(Q^2)$
- ▶ **Electric radii** and **magnetic moments**  
 $r_E^p$ ,  $\mu^p$ ;  $r_E^n$ ,  $\mu^n$

→ Comparison to experiment

## Elastic electron scattering:



## Invariant form factors:

$$F_{\Sigma'\Sigma}^{\nu}(Q^2) = \langle P', J, \Sigma', T, M_T | \hat{J}_{\text{em}}^{\nu} | P, J, \Sigma, T, M_T \rangle$$

$$\text{with } Q^2 = -q^2; \quad q^{\mu} = P^{\mu} - P'^{\mu}$$

# Transition Matrix Elements in Point Form

Incoming baryon state:  $|V, M, J, \Sigma\rangle \hat{=} |P, J, \Sigma\rangle$   
 Outgoing baryon state:  $|V', M', J', \Sigma'\rangle \hat{=} |P', J', \Sigma'\rangle$   
 Transition operator:  $\hat{O} = \hat{J}_{em}^\mu$





$$\begin{aligned}
 & \langle V', M', J', \Sigma' | \hat{J}_{em}^\mu | V, M, J, \Sigma \rangle = \\
 & = \frac{2}{MM'} \sum_{\sigma_i \sigma'_i} \sum_{\mu_i \mu'_i} \int d^3 \vec{k}_2 d^3 \vec{k}_3 d^3 \vec{k}'_2 d^3 \vec{k}'_3 \\
 & \times \sqrt{\frac{(\sum_i \omega'_i)^3}{\prod_i 2\omega'_i}} \prod_{\sigma'_i} D_{\sigma'_i \mu'_i}^{* \frac{1}{2}} \{R_W [k'_i; B(V')]\} \Psi_{M' J' \Sigma'}^* (\vec{k}'_1, \vec{k}'_2, \vec{k}'_3; \mu'_1, \mu'_2, \mu'_3) \\
 & \times \langle p'_1, p'_2, p'_3; \sigma'_1, \sigma'_2, \sigma'_3 | \hat{J}_{rd}^\mu | p_1, p_2, p_3; \sigma_1, \sigma_2, \sigma_3 \rangle \\
 & \times \sqrt{\frac{(\sum_i \omega_i)^3}{\prod_i 2\omega_i}} \prod_{\sigma_i} D_{\sigma_i \mu_i}^{\frac{1}{2}} \{R_W [k_i; B(V)]\} \Psi_{MJ\Sigma} (\vec{k}_1, \vec{k}_2, \vec{k}_3; \mu_1, \mu_2, \mu_3) \\
 & \times 2MV_0 \delta^3 (M\vec{V} - M'\vec{V}' - \vec{q})
 \end{aligned}$$

where  $p_i = B_c(V)k_i$ ,  $p'_i = B_c(V')k'_i$ , and  $\omega_i = \sqrt{\vec{k}_i^2 + m_i^2}$



# Electromagnetic Nucleon Form Factors

## Covariant predictions of the GBE CQM:







Low-energy QCD  
RCQM  
GBE RCQM  
Spectroscopy  
Light, strange, charm, bottom  
Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF  
Summary



# Electromagnetic Nucleon Form Factors

## Covariant predictions of the GBE CQM:



R.F. Wagenbrunn, S. Boffi, W. Klink, W. Plessas, and M. Radici: Phys. Lett. **B511** (2001) 33

M. Rohmoser: Diploma Thesis, Univ. of Graz, 2013

Low-energy QCD  
RCQM  
GBE RQM  
Spectroscopy  
Light, strange, charm, bottom  
Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary

# Nucleon Electric Radii and Magnetic Moments

Electric radii  $r_E^2$  [fm<sup>2</sup>]

Baryon	GBE PFSM	Experiment
$p$	0.82	$0.7692 \pm 0.0123$ <sup>1)</sup> $0.70870 \pm 0.00113$ <sup>2)</sup>
$n$	-0.13	$-0.1161 \pm 0.0022$

<sup>1)</sup> CODATA value (PDG)

<sup>2)</sup> Pohl et al.: Nature **466** (2010) 213

Magnetic moments  $\mu$  [n.m.]

Baryon	GBE PFSM	Experiment
$p$	2.70	2.792847356
$n$	-1.70	-1.9130427

K. Berger, R.F. Wagenbrunn, and W. Plessas: Phys. Rev. D **70**, 094027 (2004)

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom

Structure  
Nucleon E.m.  
Baryon E.m.

Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary

# Nucleon $r_E^2$ and $\mu$ – Nonrelativistic !!!

Electric radii  $r_E^2$  [fm<sup>2</sup>]

Baryon	GBE PFSM	GBE NR1A	Experiment
$p$	0.82	0.10	$0.7692 \pm 0.0123$ <sup>1)</sup> $0.70870 \pm 0.00113$ <sup>2)</sup>
$n$	-0.13	-0.01	$-0.1161 \pm 0.0022$

<sup>1)</sup> CODATA value (PDG)

<sup>2)</sup> Pohl et al.: Nature **466** (2010) 213

Magnetic moments  $\mu$  [n.m.]

Baryon	GBE PFSM	GBE NR1A	Experiment
$p$	2.70	2.74	2.792847356
$n$	-1.70	-1.82	-1.9130427

K. Berger, R.F. Wagenbrunn, and W. Plessas: Phys. Rev. D **70**, 094027 (2004)

Low-energy QCD  
 RCQM  
 GBE RCQM  
 Spectroscopy  
 Light, strange, charm, bottom  
 Structure  
 Nucleon E.m.  
 Baryon E.m.  
 Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
 N Scalar FF  
 N Gravitational FF

Summary

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom


Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
 $N$  Scalar FF  
 $N$  Gravitational FF

Summary

## Nucleons $N$


# Proton Electric Form Factor

$$G_E^p = \frac{2}{3} G_E^u - \frac{1}{3} G_E^d$$




# Neutron Electric Form Factor

$$G_E^n = \frac{2}{3} G_E^d - \frac{1}{3} G_E^u$$



# Proton Magnetic Form Factor


$$G_M^p = \frac{2}{3} G_M^u - \frac{1}{3} G_M^d$$





# Neutron Magnetic Form Factor

$$G_M^n = \frac{2}{3} G_M^d - \frac{1}{3} G_M^u$$



# Conclusions from Nucleon Flavor Analysis

- ▶ **Flavor analysis of nucleon e.m. form factors** in a relativistically invariant framework (point form).
- ▶ The **GBE RCQM** predicts flavor contributions in reasonable agreement with **experimental data**.
- ▶ The GBE RCQM relies on  $\{QQQ\}$  degrees of freedom only; no explicit  $\{QQQQ\bar{Q}\}$  etc.
- ▶ No explicit **meson-cloud effects** are included.
- ▶ No **strangeness content** in the nucleon for the low momentum transfers considered here.
- ▶ With respect to  $F_2^d/F_2^u$  three different phenomenological analyses give **distinct answers**.
- ▶ Details:  
M. Rohrmoser, Ki-Seok Choi, and W. Plessas: arXiv:1110.3665  
W. Plessas: Mod. Phys. Lett. A **28**, 136022 (2013)

# $\Delta$ and Hyperon E.m. Form Factors

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom

Structure  
Nucleon E.m.  
**Baryon E.m.**  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF


Summary

$\Delta$

$\Lambda, \Sigma, \Xi$

$\Sigma^*, \Xi^*, \Omega$


# Electric $\Delta$ Form Factors



GBE RCQM: Ki-Seok Choi: PhD Thesis, Univ. Graz, 2011

Lattice QCD: C. Alexandrou et al. Phys. Rev. D **79** (2009) 014507

# Magnetic $\Delta$ Form Factors



GBE RCQM: Ki-Seok Choi: PhD Thesis, Univ. Graz, 2011

Lattice QCD: C. Alexandrou et al. Phys. Rev. D **79** (2009) 014507

## Axial **Charges** and Axial **Form Factors**

of

$N$  Ground State and  $N^*$  Resonances

as well as

$\Delta, \Sigma, \Xi, \Sigma^*, \Xi^*$

Low-energy

QCD

RCQM

GBE RCQM

Spectroscopy

Light, strange,  
charm, bottom

Structure

Nucleon E.m.

Baryon E.m.

Axial FFs



$\pi NN, \pi N\Delta$

$N$  Scalar FF

$N$  Gravitational FF

Summary

## Covariant predictions of the GBE RCQM:



$$g_A^{GBE} = 1.15 \quad \text{vs.} \quad g_A^{exp} = 1.2695 \pm 0.0029$$

L.Ya. Glozman, M. Radici, R.F. Wagenbrunn, S. Boffi, W. Klink, and W. Plessas: Phys. Lett. B **516**, 183 (2001)

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom

Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
 $N$  Scalar FF  
 $N$  Gravitational FF

Summary

# Microscopic Description


of

# Meson-Baryon Interaction Vertices



# Meson-Baryon Interaction Vertices

## Interaction vertices



$$F_{i \rightarrow f} = (2\pi)^4 \langle f | \mathcal{L}_I(0) | i \rangle \equiv \langle V', M', J', \Sigma' | \hat{D}_{\text{rd}}^\pi | V, M, J, \Sigma \rangle$$

where



$$\langle p'_1, p'_2, p'_3; \sigma'_1, \sigma'_2, \sigma'_3 | \hat{D}_{\text{rd}}^\pi | p_1, p_2, p_3; \sigma_1, \sigma_2, \sigma_3 \rangle = 3\mathcal{N}_S \frac{ig_{qqm}}{2m_1 (2\pi)^{\frac{3}{2}}} \bar{u}(p'_1, \sigma'_1) \gamma_5 \gamma_\mu \lambda_m u(p_1, \sigma_1) \tilde{q}^\mu 2p_{20} \delta(\vec{p}_2 - \vec{p}'_2) 2p_{30} \delta(\vec{p}_3 - \vec{p}'_3) \delta_{\sigma_2 \sigma'_2} \delta_{\sigma_3 \sigma'_3}$$

and

$$G_{\pi NN}(Q^2) = \frac{1}{f_{\pi NN}} \frac{m_\pi \sqrt{2\pi}}{\sqrt{2M_N}} \frac{\sqrt{E'_N + M'_N}}{E'_N + M'_N + \omega} \frac{F_{i \rightarrow f}}{Q_z}$$

$$G_{\pi N\Delta}(Q^2) = -\frac{1}{f_{\pi N\Delta}} \frac{3\sqrt{2\pi}}{2} \frac{m_\pi}{\sqrt{E'_N + M'_N} \sqrt{2M_\Delta}} \frac{F_{i \rightarrow f}}{Q_z}$$

# $\pi NN$ and $\pi N\Delta$ Interaction Vertices



T. Melde, L. Canton, and W. Plessas: Phys. Rev. Lett. **102**, 132002 (2009)

Low-energy QCD  
RCQM  
GBE RCQM  
Spectroscopy  
Light, strange, charm, bottom  
Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF  
Summary



# Form-Factor Parametrizations

$$G(\vec{q}^2) = \frac{1}{1 + \left(\frac{\vec{q}}{\Lambda_1}\right)^2 + \left(\frac{\vec{q}}{\Lambda_2}\right)^4}$$

$$G(Q^2) = \frac{1}{1 + \left(\frac{Q}{\Lambda}\right)^2}$$

		RCQM	SL	PR		LIU	ERK	ALX
Structure	$\frac{f_N^2}{4\pi}$	0.0691	0.08	0.075		0.0649	0.0481	0.0412
	$\Lambda_1$	0.451	0.453	0.940	$\Lambda$	0.747	0.614	1.65
	$\Lambda_2$	0.931	0.641	1.102		-	-	-
$\Delta$	$\frac{f_\Delta^2}{4\pi}$	0.188	0.334	0.478				
	$\Lambda_1$	0.594	0.458	0.853				
	$\Lambda_2$	0.998	0.648	1.014				

T. Melde, L. Canton, and W. Plessas: Phys. Rev. Lett. **102**, 132002 (2009)

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom

Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary

# Nucleon **Scalar** Form Factor and $\pi$ -N **Sigma** Term

## $S(Q^2)$

Invariant ME of  $\hat{S} = \frac{m_u + m_d}{2}(\bar{u}u + \bar{d}d)$ :


$$\langle P' J \Sigma' | \hat{S} | P J \Sigma \rangle = \bar{U}(P') U(P) S(Q^2)$$

## $\sigma_N$

Scalar form factor at zero momentum transfer:

$$\sigma_N = S(Q^2 = 0)$$

# Nucleon Scalar Form Factor $S(Q^2)$



red line

scalar form factor

green line

u-contribution to scalar form factor

blue line

d-contribution to scalar form factor

Low-energy  
QCD

RCQM

GBE RCQM

Spectroscopy

Light, strange,  
charm, bottom

Structure

Nucleon E.m.

Baryon E.m.

Axial FFs

$\pi NN$ ,  $\pi N\Delta$

$N$  Scalar FF

$N$  Gravitational FF

Summary

Prediction of the Goldstone-boson-exchange RCQM:

$$\sigma_N \approx 8.0 \text{ MeV}$$

To be compared, e.g., with:

$$\sigma_N \approx 35 \pm 5 \text{ MeV} \quad [1]$$

$$18 \pm 5 \text{ MeV} \quad [2]$$

$$31 \pm 7 \text{ MeV} \quad [3]$$

etc.

[1] J. Gasser and H. Leutwyler: Phys. Rep. 87 (1982) 77

[2] S. Güsken et al. (SESAM Coll.): Phys. Rev. D 59 (1999) 054504

[3] R. Horsley et al. (QCDSF-UKQCD Coll.): Phys. Rev. D 85 (2012) 034506

etc.

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom


Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary

# Gravitational Form Factors of the Nucleon



# Gravitational Form Factors



Invariant ME of **energy-momentum tensor**  $\hat{\Theta}^{\mu\nu}$ :

$$\langle P' J \Sigma' | \hat{\Theta}^{\mu\nu} | P J \Sigma \rangle = \bar{U}(P') \left[ \gamma^{(\mu} \bar{P}^{\nu)} A(Q^2) + \frac{i}{2M} \bar{P}^{(\mu} \sigma^{\nu)} B(Q^2) + \frac{q^\mu q^\nu - q^2 g^{\mu\nu}}{M} C(Q^2) \right] U(P)$$

$$A(Q^2) \sim \langle P' J \Sigma' | \Theta^{00} | P J \Sigma \rangle$$

Low-energy

QCD

RCQM

GBE RCQM

Spectroscopy

Light, strange,  
charm, bottom

Structure

Nucleon E.m.

Baryon E.m.

Axial FFs


$\pi NN$ ,  $\pi N\Delta$

$N$  Scalar FF

$N$  Gravitational FF

Summary

# Nucleon Gravitational Form Factor $A(Q^2)$



# Summary and Conclusions

- ▶ Surprisingly **good agreement** of predictions by the  $\{QQQ\}$  GBE RCQM with experimental data (wherever such data are available)
- ▶ **Small deviations** left in some observables, such as electric radii and magnetic moments
- ▶ Surprisingly **good agreement** of predictions by the GBE RCQM with lattice-QCD results
- ▶ Most important symmetries of the GBE RCQM:
  - ▶ **SB $\chi$ S**
  - ▶ **Lorentz invariance**
  - ▶ **time-reversal invariance**
  - ▶ **current conservation**
- ▶ The **non-relativistic** quark model **does not work** in any instance

## Graz

K. Berger, J.P. Day, K.-S. Choi, L. Glozman, T. Melde,  
M. Rohrmoser, R.C. Schardmüller, B. Sengl,  
R.F. Wagenbrunn  
(Theoretical Physics, University of Graz)

## Pavia

S. Boffi and M. Radici  
(INFN, Sezione di Pavia)

## Padova

L. Canton  
(INFN, Sezione di Padova)

## Iowa City

W. Klink  
(Department of Physics, University of Iowa, USA)

Low-energy  
QCD

RCQM  
GBE RCQM

Spectroscopy  
Light, strange,  
charm, bottom

Structure  
Nucleon E.m.  
Baryon E.m.  
Axial FFs  
 $\pi NN$ ,  $\pi N\Delta$   
N Scalar FF  
N Gravitational FF

Summary

Thank you very much  
for  
your attention!