

Alignment of the CMS Tracker

Latest Results from LHC Run-II

Gregor Mittag
on behalf of the CMS Collaboration

Deutsches Elektronen-Synchrotron (DESY)

22nd International Conference on Computing in High Energy and Nuclear Physics
12 October 2016



Precise tracking is key to CMS physics performance

➤ What alignment precision is needed?

- Track- p_T resolution

$$\frac{\delta p_T}{p_T} = \underbrace{C_1 \cdot p_T}_{\text{position res.}} \oplus \underbrace{C_2}_{\text{multiple scat.}}$$

$$\text{with } C_1 \propto \frac{\sigma_{\text{meas}}}{B \cdot L^2 \sqrt{N}}$$


- Effective position resolution

$$\sigma_{\text{meas}} \propto \sigma_{\text{hit}} \oplus \sigma_{\text{align}}$$

- Intrinsic hit-position resolution

$$\sigma_{\text{hit}} \approx 9 \mu\text{m (pixel)}$$

$$\sigma_{\text{hit}} \approx 20 - 60 \mu\text{m (strip)}$$



Need to keep $\sigma_{\text{align}} \ll \sigma_{\text{hit}}$

CMS DETECTOR

Total weight : 14,000 tonnes
Overall diameter : 15.0 m
Overall length : 28.7 m
Magnetic field : 3.8 T

STEEL RETURN YOKE
12,500 tonnes

SILICON TRACKERS
Pixel ($100 \times 150 \mu\text{m}$) $\sim 16\text{m}^2 \sim 66\text{M}$ channels
Microstrips ($80 \times 180 \mu\text{m}$) $\sim 200\text{m}^2 \sim 9.6\text{M}$ channels

SUPERCONDUCTING SOLENOID
Niobium titanium coil carrying $\sim 18,000\text{A}$


MUON CHAMBERS
Barrel: 250 Drift Tube, 480 Resistive Plate Chambers
Endcaps: 468 Cathode Strip, 432 Resistive Plate Chambers


PRESHOWER
Silicon strips $\sim 16\text{m}^2 \sim 137,000$ channels

FORWARD CALORIMETER
Steel + Quartz fibres $\sim 2,000$ Channels

CRYSTAL
ELECTROMAGNETIC
CALORIMETER (ECAL)
 $\sim 76,000$ scintillating PbWO_4 crystals

HADRON CALORIMETER (HCAL)
Brass + Plastic scintillator $\sim 7,000$ channels







- 1440 silicon pixel modules
 - 3D hit-position measurements

- 15 148 silicon strip modules (24 244 sensors)
 - 2D measurements ($r\phi$ direction)
 - In some layers: additional modules rotated by 100 mrad

- Difference between real and assumed geometry affects track measurement
- **Idea:** track-hit residuals r between predicted and measured hit positions as a measure of misalignment




- Difference between real and assumed geometry affects track measurement
- **Idea:** track-hit residuals r between predicted and measured hit positions as a measure of misalignment




Track-based alignment

- Difference between real and assumed geometry affects track measurement
- **Idea:** track-hit residuals r between predicted and measured hit positions as a measure of misalignment
- Simply moving module by $-r$ means
 - Change of position (alignment) parameters
 - Change of track parameters
 - Change of other residuals
- Tracks correlate alignment parameters
- Need many tracks to determine parameters for all tracker modules






- Difference between real and assumed geometry affects track measurement
- **Idea:** track-hit residuals r between predicted and measured hit positions as a measure of misalignment
- Simply moving module by $-r$ means
 - Change of position (alignment) parameters
 - Change of track parameters
 - Change of other residuals
- Tracks correlate alignment parameters
- Need many tracks to determine parameters for all tracker modules



Alignment challenge


- Deriving the optimal positions, orientations, and surface deformations that minimize residuals
- Two independent approaches in CMS complementing each other
 - HipPy (local fit)
 - MillePede-II (global fit)







- Split cosmic tracks in horizontal plane at point of closest approach to interaction region
- Differences in track quantities between the two parts indicate misalignment

- Derived updated alignment with 0 T cosmic data prior to 2016 data-taking start-up
- Mean & RMS values show **reduced bias of updated alignment** wrt. 2015 geometry
 - 2015-EOY geometry no longer valid due to temperature and magnetic field changes
 - **Performance** of updated geometry **very close to ideal** Monte Carlo




- Unbiased track-hit residuals of 1 M collision events @ 3.8 T
- Pixel-detector position known to be **very sensitive to condition changes**
- Updated alignment produced using 3.8 T cosmic-ray and collision data
- EOY-2015 no longer valid for 2016 data, mainly due to temperature and magnetic field changes
- RMS values show **improvement over the 2015 geometry by a factor of 5**




- Study unbiased residuals between tracks and primary vertices
 - In longitudinal and transverse plane
 - In bins of η and ϕ

- Cosmic-ray and collision data @ 3.8 T are used for alignment
- Sample of 1 M events collected through minimum bias triggers @ 3.8 T used for validation
- Systematic z-offset of the pixel half-shells is corrected by the updated alignment

- Since this year, CMS employs automatic procedure to continuously monitor high-level-structure movements in the pixel tracker
 - Occur for example due to temperature changes or changes of the magnetic field
- Online alignment of the high-level structures for each run with more than 20 000 events
 - Measuring the movements relative to the geometry used in data processing
- Geometry is automatically updated, if alignment corrections exceed certain thresholds






➤ Run 276318 was taken shortly after a **magnet ramp** during July 4 2016

- Resulted in **movements of the pixel** detector structures
- **Up to 30 μm** wrt. the geometry used in data processing

➤ Red horizontal lines indicate the thresholds to trigger updates


- Histogram color changes from green to orange if limit is exceeded



➤ Geometry was updated afterwards

- Much improved performance during run 276327
- Only small residual movements
 - Below 2 μm in x, y direction
 - Below 10 μm in z direction

➤ Such plots are routinely produced per run as part of the CMS data-quality monitoring







➤ Evolution of the pixel's high-level-structure movements

- Covers period from June 21 to July 12 2016, corresponding to 7 fb^{-1}
- Error bars represent the statistical uncertainties
- Grey bands represent runs during which CMS magnet was not at 3.8 T
- Vertical dashed lines illustrate updates of the alignment





- Typical movements during magnet-cycles are smaller than $50 \mu\text{m}$ in x and y , and smaller than $150 \mu\text{m}$ in z

- CMS has two independent algorithms in place to solve the highly non-trivial alignment task
- Tracker-alignment **updates in 2016 significantly improved the performance** at start-up and during data taking
- Geometry changes due to temperature and magnetic-field changes are nicely compensated
- An **automatic online calibration** workflow successfully put in place
 - **Clear improvement** of the prompt reconstruction
 - Note: Ultimate performance needed for reprocessing still requires more fine-grained alignment
- References
 - *CMS Tracker Alignment Performance Results Summer 2016*, CMS-DP-2016-063, <http://cds.cern.ch/record/2221746/>
 - *Alignment of the CMS tracker with LHC and cosmic ray data*, 2014 JINST 9 P06009, [doi:10.1088/1748-0221/9/06/P06009](https://doi.org/10.1088/1748-0221/9/06/P06009)
 - *Alignment of the CMS silicon tracker during commissioning with cosmic rays*, 2010 JINST 5 T03009, [doi:10.1088/1748-0221/5/03/T03009](https://doi.org/10.1088/1748-0221/5/03/T03009)





Back-Up








Distribution of median residuals (DMR) – Pixel





Distribution of median residuals (DMR) – Strip





















- Local linearisation of track model and minimisation requiring $d\chi^2(a)/da \stackrel{!}{=} 0$
 - Linear equation system: $\mathbf{C}\mathbf{a} = \mathbf{b}$ with $\mathbf{a}^T = (\Delta\mathbf{p}, \Delta\mathbf{q})$
- Track parameters \mathbf{q} in part of data only
 - Block structure in \mathbf{C}
- Only interested in alignment parameters \mathbf{p}
 - Problem can be reduced to $\mathbf{C}'\Delta\mathbf{p} = \mathbf{b}'$
 - Solution provides alignment parameters
 - All correlations still taken into account
- \mathbf{C}' , \mathbf{b}' by solving $N_{\text{track pars}} \times N_{\text{track pars}}$ per track
- Dramatic cost reduction:

$$N_{\text{align pars}}^2 + N_{\text{tracks}} \cdot N_{\text{track pars}}^2 \ll (N_{\text{align pars}} + N_{\text{tracks}} \cdot N_{\text{track pars}})^2$$
 - Full-scale alignment performed within $\lesssim 24$ h



http://cms.desy.de/sites/site_cms/content/e53612/e234315/e241962/e241763/posterCCP2010_brokenlines.pdf