


Heavy-quark energy-loss via angular correlations between heavy and light mesons

M. Rohrmoser, P.-B. Gossiaux, T. Gousset, J. Aichelin
from SUBATECH, Nantes

July 25, 2016
4th Heavy-Ion Jet Workshop, École Polytechnique, Palaiseau

Some samples of R_{AA}



[CMS Collaboration, CMS-PAS-HIN-16-001.]

Overview

Approach in 2 directions:

Observables

Strategy of the analysis:

- 1 heavy-light-particle (angular) correlations:
overall medium effects?
- 2 Search for origin of differences:
specific shower processes + individual parton branchings.
- 3 Extract medium dependent quantities from global results.

Production of heavy-quark showers:

situation	vacuum	Inelastic	Elastic
Description	splitting functions	model A	model B
In-medium energy-loss		additional branchings	transfer shower → medium

Mechanisms

In-medium propagation: inelastic scattering

Model A:

[Th. Renk: Phys.Rev.C **78**, 034908 (2008)]

Virtuality increases/no changes in 3-momenta per small timesteps Δt :




$$\begin{aligned} Q &\mapsto \sqrt{Q^2 + \hat{q}\Delta t}, \\ \vec{p} &\mapsto \vec{p}, \\ E &\mapsto \sqrt{E^2 + \hat{q}\Delta t}. \end{aligned} \tag{1}$$

\Rightarrow 3-momenta in shower only changed due to additional radiation!

(Azimuthal) Angular correlations

Correlations of **heavy quark** & **any light particle**:

$$\cos(\Delta\phi) = \frac{\vec{p}_{h\perp} \cdot \vec{p}_{l\perp}}{\|\vec{p}_{h\perp}\| \|\vec{p}_{l\perp}\|},$$





(Azimuthal) Angular correlations

Correlations of **heavy** quark & **any** light particle:


$$\cos(\Delta\phi) = \frac{\vec{p}_{h\perp} \cdot \vec{p}_{l\perp}}{\|\vec{p}_{h\perp}\| \|\vec{p}_{l\perp}\|},$$

from physical viewpoint pretty equivalent to

$$\cos(\Delta\theta) = \frac{\vec{p}_h \cdot \vec{p}_l}{\|\vec{p}_h\| \|\vec{p}_l\|}.$$




$$\frac{dN}{d\cos(\Delta\theta)}$$



$$\Delta Q^2 = \int_{\tau_{ini}}^{\tau_{fin}} dt \hat{q}(t)$$

Angular Broadening

$\langle \Delta\theta \rangle$ [rad]




Boxes: half-width of $\frac{dN}{d\Delta\theta}$ distribution.

More sensitive observables?

Contributions from different processes


Dominant: Small number of branchings $N_S!$



Processes of 1 quark \rightarrow 2 gluons: $\approx 10\%$


Individual shower contributions to angular correlations

→ Look at contributions from different topologies/processes with different numbers of emitted particles, e.g.:




Individual shower contributions to angular correlations

→ Look at contributions from different topologies/processes with different numbers of emitted particles, e.g.:




Individual shower contributions to angular correlations

→ Look at contributions from different topologies/processes with different numbers of emitted particles, e.g.:



Individual shower contributions to angular correlations

→ Look at contributions from different topologies/processes with different numbers of emitted particles, e.g.:




medium effect:
2nd splitting,
mostly (i.e.: for
smaller angles)

inversion of
angular ordering
in the medium



Which observable for a full shower?

Angular Ordering?

Compare angles between momenta of a light particle and:
the heavy particle... $\Delta\theta$ \rightarrow contain heavy quark branchings
the entire jet... $\Delta\theta_{\text{jet}}$ \rightarrow "history" of previous branchings




contributions from 3 splittings:





Angular Ordering?

Compare angles between momenta of a light particle and:
the heavy particle. . . $\Delta\theta$ \rightarrow contain heavy quark branchings
the entire jet. . . $\Delta\theta_{\text{jet}}$ \rightarrow "history" of previous branchings



results for arbitrary N_S :



\rightarrow indication for weakened angular ordering

black curves: $\langle \Delta\theta(\Delta\theta_{\text{jet}}) \rangle$

Conclusions for observables from model A

Observables



- angular broadening verified.
- refined analysis of broadening
→ small angles affected most strongly.
- indications for angular ordering violations.

Conclusions for observables from model A

Observables




- angular broadening verified.
- refined analysis of broadening
→ small angles affected most strongly.
- indications for angular ordering violations.

model A  model B

In-medium propagation: elastic scattering

Model B:

Forces transverse and parallel to incident 3-momenta \vec{p} + changes in particle energy; $Q=\text{constant}$:



$$\vec{p} = (\vec{0}, p) \mapsto \vec{p}' = (\vec{p}_\perp, p_\parallel),$$

$$p_\perp = \sqrt{\hat{q}\Delta t}, \quad p_\parallel = p - A\Delta t. \quad (2)$$

transverse momentum
transfer

(longitudinal)
drag force

$$A = \frac{\hat{q}}{\kappa T},$$

$A \dots$ drag force ,

$T \dots$ Temperature (medium) ,

$\kappa \dots$ proportionality constant .

(3)


[H. Berrehrh, P. B. Gossiaux, J. Aichelin, W. Cassing, E. Bratkovskaya: *Phys. Rev. C* **90**, 064906 (2014)]

\Rightarrow 3-momenta changed via stochastic force from medium, but no additional radiation!

In-medium propagation: elastic scattering

Model B:

Forces transverse and parallel to incident 3-momenta \vec{p} + changes in particle energy; $Q=\text{constant}$:



Energy \rightarrow Medium

$$\vec{p} = (\vec{0}, p) \mapsto \vec{p}' = (\vec{p}_\perp, p_\parallel),$$

$$p_\perp = \sqrt{\hat{q}\Delta t}, \quad p_\parallel = p - A\Delta t. \quad (2)$$

transverse momentum transfer

(longitudinal) drag force

$$A = \frac{\hat{q}}{\kappa T},$$

A ... drag force,

T ... Temperature (medium),


κ ... proportionality constant.

(3)

[H. Berrehrh, P. B. Gossiaux, J. Aichelin, W. Cassing, E. Bratkovskaya: Phys. Rev. C90, 064906 (2014)]

\Rightarrow 3-momenta changed via stochastic force from medium, but no additional radiation!


Differences in $\Delta\theta$ for models A and B?



Angular Broadening: different energy dependencies for **model A** and **B**?

Differences in $\Delta\theta$ for models A and B?

$\langle\Delta\theta\rangle$ [rad]



Angular Broadening: different energy dependencies for **model A** and **B**?




Summary

- Angular correlations as possible way to study medium effects!
- 2 mechanisms of energy loss simulated:
inelastic (**model A**) and elastic scattering (**model B**).
- Angular broadening reflected in results!
- ...allows to distinguish hot and dense medium from vacuum...
- ...and maybe different energy-loss mechanisms from one another
(further, ongoing studies)!

Thank you for your attention!

Backup

Branching angles



inversion
of angular
ordering in
the medium