



Low-mass dark matter searches with the LZ experiment

Maria Elena Monzani
on behalf of the LZ
Collaboration

SLAC
NATIONAL ACCELERATOR LABORATORY


SLAC, Apr 29, 2016




© 1912 W. B. Perkins

111717

LZ Detector Overview




Dual-Phase Xenon TPC




- **Electron recoil (ER) background rejection:**
 - Charge/light ratio (aka “S2/S1”)
- **3D event reconstruction:**
 - Vertical coordinate from drift time
 - Horizontal coordinates from S2 light pattern
- **External backgrounds:**
 - Fiducial volume cuts
- **Neutron rejection:**
 - Multiple scattering
 - Tagged in skin/veto


State of the Art and Projected Sensitivity



How to probe the low-mass range?



High Statistics Calibrations from LUX



Electron Recoil (ER)
(Tritiated methane)


+

Nuclear Recoil (NR)
(DD neutron gun)


↓

In-situ calibrations,
unpreced. accuracy,
lower NR threshold

NR Calibrations: towards Lower Threshold



WIMP Sensitivity with Lower Threshold



Light Dark Matter Searches with LXe





	S1 + S2	S2 - only
Nuclear Recoils	<ul style="list-style-type: none">• Vanilla WIMPs	<ul style="list-style-type: none">• Light(er) WIMPs• Asymmetric Dark Matter
Electron Recoils	<ul style="list-style-type: none">• \simkeV axion-like particles	<ul style="list-style-type: none">• subGeV hidden sector models (Rouven's talk)

Axion-Like Particles (ALP)




- Sensitivity to axions and ALPs via the axioelectric effect:
 - Nonrelativistic galactic ALPs (DM candidates)
 - ALPs emitted by brems/Compton in the Sun
- Technique pioneered in Xenon100 (see arXiv:1404.1455)



S2-only analysis: Light(er) WIMPs



Cross Section Upper Limits



S2-only analysis: Experimental Challenges



- Xenon10 results were described by Rouven yesterday
- Going to a larger detector doesn't make this any easier
 - No S2-only results from Xenon100 or LUX so far...
- Limited background rejection with S2-only analysis
 - No S1/S2 discrimination, no Z coordinate available
- Electrons can be captured by impurities in drift volume
 - Depth-dependent effect... but we don't know Z coordinate
- Incomplete extraction from liquid phase
 - Uniformity issues? increase extraction field?
- Single electron background, difficult to model/subtract
 - Correlated with larger events (at least to some degree)
 - Electrons trapped under the liquid level? increase field?
 - Imperfections in the grids? can we make better grids?

S2-only analysis: Experimental Challenges




- Xenon10 results were described by Rouven yesterday
- Going to a larger detector doesn't make this any easier
 - No S2-only results from Xenon100 or LUX so far...
- Limited background rejection with S2-only analysis
 - No S1/S2 discrimination, no Z coordinate available
- Electrons can be captured by impurities in drift volume
 - Depth-dependent effect... but we don't know Z coordinate
- Incomplete extraction from liquid phase
 - Uniformity issues? increase extraction field?
- Single electron background, difficult to model/subtract
 - Correlated with larger events (at least to some degree)
 - Electrons trapped under the liquid level? increase field?
 - Imperfections in the grids? can we make better grids?

Single Electron Sensitivity in LZ



- **Single electron signal depends on field configuration:**
 - expected >50 p.e./electron (x2 higher than Xenon10)
 - expected 97.6% extraction efficiency for electrons



Parameter	value
Gate-Anode separation (and tolerance)	13.0 mm (± 0.2 mm)
Gas gap (and tolerance)	8.0 mm (± 0.2 mm)
Field in LXe (GXe)	5.2 kV/cm (10.2 kV/cm)
Electron emission probability	97.6 %
S2 photon yield	820 ph/e
S2 width FWHM	1.2 μs
Detailed modeling	
S2 photon yield	910 ph/e
S2 photon <i>rms</i>	2.0 %
S2 width FWHM	1.0 μs to 2.0 μs ^a

^a The larger value is for diffusion-broadened S2 pulses from interactions near the cathode (see Figure 3.6.4).


LZ TDR, in preparation



- **Reaching HV specs has proven elusive in all LXe detectors**
 - very extensive fields/grid R&D/testing in progress at SLAC



SLAC Noble Liquid Test Platform





and Test Platform















Conclusions



- **Low-energy calibrations drastically improved across the field:**
 - Lower thresholds in LUX
 - Sensitivity to lighter DM particles
- **S2-only analysis is always challenging due to backgrounds**
 - No S2/S1 or Z-coordinate cut
- **Single electron sensitivity will be greatly enhanced in LZ**
 - Single electron background a possible concern (large area)
 - Tackling the single electron background very aggressively
 - Includes full-scale grid testing