



Contribution ID: 53

Type: Oral

Salicylideneaniline-Functionalized Poly(*m*-phenyleneethynylene)s as Fluorescent Turn-On Chemosensors for Cations

Tuesday 29 November 2016 14:00 (15 minutes)

Two different series of conjugated polymer, poly(*m*-phenyleneethynylenes) (*m*-PPEs) containing different amounts of salicylideneaniline moieties (50% and 100%) have been synthesized via a post-functionalization of aniline group on *m*-PPEs backbone. PPEs are successfully prepared in excellent yield (90-99%) and spectroscopically characterized the structure by ^1H , ^{13}C NMR and FTIR exhibited signals that reasonably correlate with the desired polymer. The resulting polymers displayed weak orange emission at 560 nm and undergo remarkable turn-on bright blue fluorescent emission at 450 nm response to Fe^{2+} , Fe^{3+} , Al^{3+} and Cr^{3+} without any change with other cations.

Primary author: Ms THAVORNSIN, Nopparat (Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand.)

Co-author: Mr SUKWATTANASINITT, Mongkol (Nanotec-CU Center of Excellence on Food and Agriculture, Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand.)

Presenter: Ms THAVORNSIN, Nopparat (Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand.)

Session Classification: Heron 1

Track Classification: Other related topics