



Contribution ID: 266

Type: **Invited Speaker**

## First-principles study on two dimensional dichalcogenides for hydrogen production

*Monday 28 November 2016 10:55 (20 minutes)*

Hydrogen is a promising candidate for the clean energy carrier that may replace fossil fuels. For the production of hydrogen, water splitting with efficient catalysts has been intensively studied over the past decades. Platinum, which is known to be the best catalyst for water splitting, is too expensive to be used in large-scale applications. Therefore, numerous earth-abundant materials have been investigated as a replacement of Pt. Recently, transition metal dichalcogenides (TMDs), most notably MoS<sub>2</sub>, are receiving a great deal of attention as a novel catalyst for water splitting. Although the basal plane of TMDs are efficient as catalysts, it was found recently that the sulfur vacancy in MoS<sub>2</sub> can increase the catalytic activity for hydrogen evolution.

In this presentation, motivated by the previous work, we explore the detailed mechanism for hydrogen production from the sulfur vacancy in MoS<sub>2</sub> and calculate the activation energies along the reaction path. Furthermore, we evaluate the catalytic efficiency of vacancy sites in various TMDs and suggest TMDs that may show high catalytic effects in hydrogen evolution reaction.

**Primary authors:** KIM, Kye Yeop (Department of Materials Science and Engineering Seoul National University); Prof. HAN, Seungwu (Department of Materials Science and Engineering Seoul National University)

**Co-author:** LEE, Joohee (Department of Materials Science and Engineering Seoul National University)

**Presenter:** Prof. HAN, Seungwu (Department of Materials Science and Engineering Seoul National University)

**Session Classification:** Heron 1

**Track Classification:** Theory and simulation related to nanosystem