



The Cockcroft Institute  
of Accelerator Science and Technology

# Cross-talk between experiments and LSS losses at FCC-hh


Rob Appleby  
Cockcroft Institute/University of Manchester

DRAFT



## MDI activities


- Experimental Interaction Region
  - **One of critical areas defining FCC-hh performance**
- Design tasks of EuroCirCol IR Work Package
  - **Coordination**
    - JAI/Oxford (lead), CERN, task 3.1
  - **Development of the interaction region lattice**
    - JAI/Oxford (lead), CERN, task 3.2
  - **Design of machine detector interface**
    - CI/Manchester (lead), INFN, CERN, task 3.3
  - **Study of beam-beam interaction**
    - EPFL (lead), CERN, task 3.4





## Cross-talk between IRs

- Given the 50 TeV on 50 TeV protons collisions at a Lumi of 5E34, experimental cross-talk is a possible issue at FCC-hh.
- Here we use the DPMJETIII generator in its upgraded form, valid at this energy, to look at potential cross-talk contributions.
- We take IPA to IPB as a representative high lumi to other experiment, with ~5k of straight and arc between them.
  - (We use IPA to IPB for ease...same results as IPG->IPH)
- This preliminary study looks at
  - Inelastic protons through beam pipe vacuum (i.e. optics)
  - Elastic protons through beam pipe vacuum (i.e. optics)
  - Muons travelling through rock from IPA to IPB.
  - (Muons through beam pipe vacuum automatically)




# p-p EVENT GENERATOR (DPMJET-III) DEVELOPMENT


PhD thesis of A. Fedynitch  
supervisors R. Engel (KIT) and A. Ferrari

- implementation of *new parton distributions functions and improved hard scatter*
- all Regge parameters re-fitted to match cross-sections from low energy up to LHC as good as possible
- Exists inside development version of FLUKA
- Crucial for loads on the inner triplet (CERN) and experiment cross-talk (Manchester)**



ATLAS average  $p_T$



7 TeV



LHCb forward energy flow









The Cockcroft Institute  
of Accelerator Science and Technology

[Rates are for prompt  
particles (just after  
IP), with  $E > 10$  GeV]

# 50 TeV-50 TeV pp collision particle content




Mean protons per primary 0.89  
Mean energy of protons 14757 GeV  
Mean mu- per primary 0.007  
Mean energy of mu- 34 GeV  
Mean CH per primary 17.4  
Mean energy of CH 373 GeV  
Mean gammas per primary 13.8  
Mean energy of gammas 209 GeV








The Cockcroft Institute  
of Accelerator Science and Technology


# 50 TeV-50 TeV pp collision inelastic protons and muons



(For 1M collisions)






# The simulation framework





The Cockcroft Institute  
of Accelerator Science and Technology

## Optics and lattice properties



Optics:  
 $L^*=45\text{m}$  optics with matched IRA  
TAS at 50mm, length 750m  
 $\Beta^*=0.3\text{m}$   
NO crossing angle (hence this dispersion)  
Norm emit  $2.2\text{E-}6\text{ m}$





$L=5\text{E}34$ ,  $\sigma_{\text{ie}}=108\text{mb}$



The Cockcroft Institute  
of Accelerator Science and Technology

# Inelastic protons : loss patterns

$L^*=45\text{m}$  optics with matched IRA  
 $\text{Beta}^*=0.3\text{m}$ , NO crossing angle  
Losses assume  $L=5E34$ ,  $\sigma_{\text{ie}}=108\text{mb}$









The Cockcroft Institute  
of Accelerator Science and Technology

# Inelastic protons : loss rates

$L^*=45\text{m}$  optics with matched IRA  
 $\text{Beta}^*=0.3\text{m}$ , NO crossing angle  
Losses assume  $L=5E34$ ,  $\sigma_{\text{ie}}=108\text{mb}$



Note - losses are per element, i.e. the high p/s on Q1 are distributed along the magnet length



- losses consistent with estimation from FLUKA group ( $\sim 5E8$  p/s with crossing angle)
- losses are exceed by pion loss, which will be shielded by liner
- losses will be reduced by crossing angle and impact of detector dipole
- losses will create local showers – to be studied




The Cockcroft Institute  
of Accelerator Science and Technology

# Inelastic protons : IPA->IPB

IPB energy distribution



L\*=45m optics with matched IRA  
Beta\*=0.3m, NO crossing angle  
Losses assume L=5E34, sigma-ie=108mb



**Particles at IPB PER BX [nom]: 6.2**

**Particles at IPB PER BX [ult]: 37.4**

Particles at IPB PER SEC [nom]: 311613300


Particles at IPB PER SEC [ult]: 11218078800

Power [W] at IPB [nom]: 1575

Power [W] at IPB [ult]: 9451

**Mean energy of protons at IPB [GeV]: 49962**

**X-STDDEV=Around 100 um**





## Inelastic protons : On TAS


Proton loss rate on TAS [p/s] [nom] = 1958202000.

Proton power rate on TAS [W] [nom] = 61 (due to low energy protons)

Proton loss rate on TAS [p/s] [ult] = 11749212000

Proton power rate on TAS [W] [ult] = 365

Picture will be changed by detector dipole and crossing angle





## Elastic protons

Particles at IPB PER BX [nom]: 85

Particles at IPB PER BX [ult]: 512

Particles at IPB PER SEC [nom]: 4268878200

Particles at IPB PER SEC [ult]: 1.536796152e+11


Power [W] at IPB [nom]:

Power [W] at IPB [ult]:

Mean energy of protons at IPB [GeV]: 50000.0

Conclusion : elastics stay well within beampipe and TAS (as LHC)

-> Emittance growth (as LHC)



## Muons through rock : theory

Muon stopping power for high energy muons can be described by

$$\left\langle -\frac{dE}{dx} \right\rangle = a(E) + b(E)E$$

Where E is the energy, a(E) models electronic stopping power (ionization and excitation) and b(E) is due to radiative processes, such that

$$b = b_{brems} + b_{pair} + b_{nucl}$$

Note b(E) is << a(E) for most materials, when E <= 100 GeV.

The range in the continuous slowing down approximation is given by

$$R(E) = \int_{E_0}^E (a(E') + b(E')E')^{-1} dE'$$

Which is what we use for FCC. However we can do the integral at high energy, when a(E) and b(E) are constant to get

$$R(E) \sim \frac{1}{b} \ln\left(1 + \frac{E}{E_c}\right)$$

Where the electronic and radiative losses are equal at the 'critical energy'

$$a(E_c) = E_c b(E_c)$$



The Cockcroft Institute  
of Accelerator Science and Technology


## Muons through rock : *a* and *b*

[http://pdg.lbl.gov/2015/AtomicNuclearProperties/HTML/standard\\_rock.html](http://pdg.lbl.gov/2015/AtomicNuclearProperties/HTML/standard_rock.html)

T [MeV]	p [MeV/c]	Ionization	brems	pair	photon	nucl	Radloss	dE/dx	CSDA	Range	delta	beta	dE/dx_R
									[g/cm^2]				
1.000E+00	1.457E+01	2.643E+00	0.000E+00	0.000E+00	4.771E-05	4.771E-05	5.286E+00	2.311E-03	0.0000	0.13661	4.060E+01		
1.200E+00	1.597E+01	3.517E+01	0.000E+00	0.000E+00	4.780E-05	4.780E-05	3.517E+01	7.619E-03	0.0000	0.14944	3.517E+01		
1.400E+00	1.726E+01	3.112E+01	0.000E+00	0.000E+00	4.789E-05	4.789E-05	3.112E+01	1.368E-02	0.0000	0.16119	3.112E+01		
1.700E+00	1.903E+01	2.667E+01	0.000E+00	0.000E+00	4.802E-05	4.802E-05	2.667E+01	2.413E-02	0.0000	0.17725	2.667E+01		
2.000E+00	2.066E+01	2.342E+01	0.000E+00	0.000E+00	4.816E-05	4.816E-05	2.342E+01	3.616E-02	0.0000	0.19186	2.342E+01		
2.500E+00	2.312E+01	1.960E+01	0.000E+00	0.000E+00	4.838E-05	4.838E-05	1.960E+01	5.961E-02	0.0000	0.21376	1.960E+01		
3.000E+00	2.536E+01	1.694E+01	0.000E+00	0.000E+00	4.860E-05	4.860E-05	1.694E+01	8.713E-02	0.0000	0.23336	1.673E+01		
3.500E+00	2.742E+01	1.498E+01	0.000E+00	0.000E+00	4.883E-05	4.883E-05	1.498E+01	1.186E-01	0.0000	0.25120	1.462E+01		
4.000E+00	2.935E+01	1.347E+01	0.000E+00	0.000E+00	4.905E-05	4.905E-05	1.347E+01	1.539E-01	0.0000	0.26763	1.302E+01		
4.500E+00	3.116E+01	1.227E+01	0.000E+00	0.000E+00	4.927E-05	4.927E-05	1.227E+01	1.928E-01	0.0000	0.28290	1.176E+01		
5.000E+00	3.289E+01	1.129E+01	0.000E+00	0.000E+00	4.950E-05	4.950E-05	1.129E+01	2.353E-01	0.0000	0.29720	1.074E+01		
5.500E+00	3.453E+01	1.047E+01	0.000E+00	0.000E+00	4.972E-05	4.972E-05	1.047E+01	2.814E-01	0.0000	0.31066	9.902E+00		

Standard rock data:



Quantity	Value
$\langle Z/A \rangle$	0.50000
Specific gravity	2.650 g cm <sup>-3</sup>
Critical energy	49.13 MeV (for e-), 47.74 MeV (for e+)
Muon critical energy	693. GeV






The Cockcroft Institute  
of Accelerator Science and Technology

# Muon range through rock



3. Chord through FCC-hh ring  
circum=100 km. r=15.9 km.  
 $C=2\pi(5.964 \text{ km}/100 \text{ km}) = 0.37 r$



$$\text{Chord} = 2r \sin(c/2) \\ = 5.92 \text{ km}$$

Max energy is 7.9 TeV  
Max range is 2.2 km

Do not expect many muons through rock

Needs checking with Monte Carlo to include fluctuations and straggling

Note this is prompt muons. Decay muons being looked at now



## Summary

- Cross-talk between IPs is a concern for FCC-hh
- Using a new version of DPMJETIII for an IP source and MADX/calcs we're able to study these effects
- **Inelastic protons through beam pipe vacuum** give a rate of protons per BX at the IPB of 6 for nominal parameters and 37 for ultimate parameters, with a spot of 100 um.
  - May cause some background due to position spread...to be checked
  - Proton losses in the LSS are high but expected... protection from TAS and TAN and liner (but not highest contribution, which is pions)
  - Maybe TCL-style collimations further down would be useful for DS protection
- **Elastic protons through beam pipe vacuum** give rise to a rate of protons per BX at the IPB of 85 for nominal parameters and 512 for ultimate parameters, with a spot comparable to the nominal beam size. This will lead to emittance growth.
- **Muons travelling through rock** from IPA to IPB will only have the range in extreme cases to give a meaningful cross-talk flux
- **Muons, due to energy, will not last for long in the optics**, if produced heading in the forward region
- Both calculations would benefit from shower codes, in the proton case to check local effects of high proton loss and in the muon case to properly include range straggling and fluctuations.
  - E.g. potential muon background from inelastic losses close to IPB.
- For Rome, will add
  - decay muons to tunneling
  - crossing angle to proton transport
  - Detector spectrometer and compensator magnets