

Applications of Accelerators


CERN Introductory Accelerator School
Budapest, 2016.

Dr. Suzie Sheehy

John Adams Institute for Accelerator Science
University of Oxford

- *“A beam of particles is a very useful tool...”*

-Accelerators for Americas Future
Report, pp. 4, DoE, USA, 2011



There are roughly 35,000 accelerators in the world
(Above 1 MeV...)


Outline

1. Medical imaging and treatment
2. Industrial uses of accelerators
3. Synchrotron light sources
4. Neutron sources
5. Energy and security applications
6. Historical & cultural applications

1. Medical Applications

- Around 1/3 of people in the will die from cancer...
- But diagnosis is no longer a death sentence!

Patients cured by the major cancer treatment modalities



Reference
 Cancer Services Collaborative 2002
www.nhs.uk/npat

X-ray radiotherapy

Linac

Foil to produce x-rays

Collimation system






Image: copyright Varian medical systems



7

Charged Particle Therapy



- Greater dose where needed
- Less morbidity for healthy tissue
- Less damage to vital organs

8

Energy loss in materials

The relativistic version of the formula reads:

$$-\frac{dE}{dx} = \frac{4\pi}{m_e c^2} \cdot n z^2 \cdot \left(\frac{e^2}{4\pi\epsilon_0}\right)^2 \cdot \left[\ln\left(\frac{2m_e c^2 \beta^2}{I}\right) - \beta^2 \right]$$

$$\text{where } -\frac{dE}{dx} = \frac{4\pi n z^2}{m_e v^2} \cdot \left(\frac{e^2}{4\pi\epsilon_0}\right)^2 \cdot \left[\ln\left(\frac{2m_e v^2}{I}\right) \right]$$

where

$$\beta = v / c$$

v velocity of the particle

E energy of the particle

x distance travelled by the particle

c speed of light

z particle charge

e charge of the electron

m_e rest mass of the electron

n electron density of the target

I mean excitation potential of the target



ϵ_0 vacuum permittivity

High speed -> small energy loss
Low speed -> high energy loss


Proton therapy

– “Hadron therapy” = Protons and light ions

- Used to treat localised cancers
- Less morbidity for healthy tissue
- Less damage to vital organs
- Particularly for childhood cancers




A few developments




Spot Scanning



Proton Radiography




Gantries



FFAG Accelerators



Dielectric Wall Accelerators




Laser Plasma Accelerators



HEP community can contribute accelerators AND other expertise!

Radioisotope production

- Accelerators (compact cyclotrons or linacs) are used to produce radio-isotopes for medical imaging.
- 7-11MeV protons for short-lived isotopes for imaging
- 70-100MeV or higher for longer lived isotopes



- Positron emission tomography (PET) uses Fluorine-18, half life of ~110 min



- Fluorodeoxyglucose or FDG carries the F18 to areas of high metabolic activity
- 90% of PET scans are in clinical oncology

Radiopharmaceuticals

p, d, ³He, ⁴He
beams

Isotopes used for PET, SPECT and
Brachytherapy etc...




TABLE 2.1. THE RADIOISOTOPES THAT HAVE BEEN USED AS TRACERS IN THE PHYSICAL AND BIOLOGICAL SCIENCES

Isotope	Isotope	Isotope
Actinium-225	Fluorine-18	Oxygen-15
Arsenic-73	Gallium-67	Palladium-103
Arsenic-74	Germanium-68	Sodium-22
Astatine-211	Indium-110	Strontium-82
Beryllium-7	Indium-111	Technetium-94m
Bismuth-213	Indium-114m	Thallium-201
Bromine-75	Iodine-120g	Tungsten-178
Bromine-76	Iodine-121	Vanadium-48
Bromine-77	Iodine-123	Xenon-122
Cadmium-109	Iodine-124	Xenon-127
Carbon-11	Iron-52	Yttrium-86
Chlorine-34m	Iron-55	Yttrium-88
Cobalt-55	Krypton-81m	Zinc-62
Cobalt-57	Lead-201	Zinc-63
Copper-61	Lead-203	Zirconium-89
Copper-64	Mercury-195m	
Copper-67	Nitrogen-13	

2. Industrial accelerators

15

Ion implantation



Images courtesy of Intel


- Electrostatic accelerators are used to deposit ions in semiconductors.

16


Electron beam processing

In the US, potential markets for industrial electron beams total \$50 billion per year.

- 33% Wire cable tubing
- 32% Ink curing
- 17% shrink film
- 7% service
- 5% tires
- 6% other



<http://rscnuclearcable.com/capabilities.htm>



When polymers are cross-linked, can become:

- stable against heat,
- increased tensile strength, resistance to cracking
- heat shrinking properties etc

Equipment sterilisation

Manufacturers of medical disposables have to kill every germ on syringes, bandages, surgical tools and other gear, without altering the material itself.

E-beam sterilisation works best on simple, low density products.


Advantages: takes only a few seconds (gamma irradiation can take hours)

Disadvantages: limited penetration depth, works best on simple, low density products (syringes)



The IBA rhodotron – a commercial accelerator used for e-beam sterilisation

Food irradiation



'Cold pasteurisation' or 'electronic pasteurisation'

Uses electrons (from an accelerator) or X-rays produced using an accelerator.

The words 'irradiated' or 'treated with ionising radiation' must appear on the label packaging.

In the US all irradiated foods have this symbol



Foods authorised for irradiation in the EU:



Lower dose




Higher dose

19


Other uses in industry...

- Hardening surfaces of artificial joints
- Removal of NO_x and SO_x from flue gas emissions
- Scratch resistant furniture

Treating waste water or sewage
Purifying drinking water
(Without additional chemicals...)



Irradiating topaz and other gems with electron beams to change the colour



<http://www.nrc.gov/reading-rm/doc-collections/fact-sheets/irradiated-gemstones.html>

<http://www.symmetrymagazine.org/article/october-2009/cleaner-living-through-electrons>



20

3. Synchrotron Light Sources




Image courtesy of ESRF

Synchrotron radiation is emitted by charged particles when accelerated radially



Produced in synchrotron radiation sources using bending magnets, undulators and wigglers

THE ELECTROMAGNETIC SPECTRUM



Synchrotron radiation: microwaves to hard x-rays (user can select)

High flux = quick experiments!


Pulsed structure = resolution of processes down to picoseconds

X-Ray crystallography

2014 was the International Year of Crystallography

Protein crystallography is a standard technique at synchrotron light sources (Diamond light source has 5 beamlines devoted to it)

The hardest part is forming the crystal...



For some great overview videos of crystallography, see:
<http://www.richannel.org/collections/2013/crystallography>

Hard condensed matter science
 Applied material science
 Engineering
 Chemistry
 Soft condensed matter science
 Life sciences
 Structural biology
 Medicine
 Earth and science
 Environment
 Cultural heritage
 Methods and instrumentation

© CCLRC



Diffraction pattern from pea lectin

25

Synchrotron Radiation Science

Biology

Reconstruction of the 3D structure of a nucleosome (DNA packaging) with a resolution of 0.2 nm



In 1990 scientists determined the structure of a strain of foot & mouth virus using Daresbury SRS.

Archeology/Heritage

A synchrotron X-ray beam at the SSRL facility illuminated an obscured work erased, written over and even painted over of the ancient mathematical genius Archimedes, born 287 B.C. in Sicily.



Using X-Ray induced fluorescence


26




4. Neutron Spallation Sources



<https://youtu.be/VESMU7JfVHU?t=21>

‘Neutrons tell you where atoms *are* and what atoms *do*’






ISIS Accelerators and Targets

- H⁻ ion source (17 kV)
- 665 kV H⁻ RFQ
- 70 MeV H⁻ linac
- 800 MeV proton synchrotron
- Extracted proton beam lines
- Targets
- Moderators

Pulsed beam of 800 MeV
(84% speed of light) protons
at 50 Hz
Average beam current
is 230 μA (2.9 × 10¹³ ppp)

184 kW on target (148 kW to
TS-1 at 40 pps, 36 kW to TS-2 at
10 pps).

$$P = 800[MV] \times 230[\mu A] = 184[kW]$$



Calculating beam power

- Power = Work/time $P = \frac{W}{T}$
- Work = force x distance $W = Fd$
- Force on particle in an electric field $F = qE$
- We know the electric field is (voltage/distance) and the protons (charge +1) have gained 800 MeV, so V=800MV.
- Also know current = charge/time

$$P = 800[MV] \times 230[\mu A] = 184[kW]$$





Image courtesy ISIS, STFC.

Unblocking oil pipes

- **Asphaltenes** are a complex mixture of molecules that can sometimes **block oil pipes**
- Research to more easily **predict** and **prepare** for the formation of asphaltene deposits
- Result in **fewer blockages** and **big savings** for the oil industry.





“ISIS allowed us to understand more clearly how asphaltenes aggregate, an important observation from a flow assurance point of view and should allow more efficient extraction of hydrocarbons in the future.”
 –Edo Boek, Schlumberger Cambridge Research, Senior Research Scientist



Stresses in Airbus A380 Wing


- Aircraft manufacturer Airbus has used ISIS since 2006
- Research into aluminium alloy weld integrity for aircraft programmes
- Residual stresses from welding cause weaknesses and the possibility of cracks
- ISIS neutrons look deep inside engineering components to measure stress fields



“Residual stress measurement at ISIS has been invaluable in researching and developing existing and novel material manufacturing and processing techniques .”
 - Richard Burguete, Airbus Experimental Mechanics Specialist

Understanding infant lung structure


- Natural lung surfactant allows oxygen into the bloodstream
- Absence in premature babies causes breathing difficulties
- ISIS mimicked change in lung capacity to discover how proteins and phospholipids act together
- Helping to develop synthetic lung surfactants which can be more precisely targeted at clinical needs to help save babies' lives



“ISIS is the premier place in the world to work with neutrons and liquid surfaces. In collaboration with the University of Queensland we were able to discover how proteins and phospholipids act together to enable lung function.”
 - Dr Stephen Holt, ISIS neutron scientist

Fast neutron testing for the semiconductor industry


- **Atmospheric neutrons** collide with microchips and **upset microelectronic devices** every few seconds
- **300 x greater effect** at **high altitude**
- **ISIS enables manufacturers to mitigate against the problem** of cosmic radiation
- **Increased confidence** in the quality and safety of aerospace electronic systems



“ISIS is one of few facilities in the world capable of producing enough very high energy neutrons to perform accelerated testing.”
 -Andrew Chugg, MBDA, SEEDER consortium

5. Energy and Security Applications

Cargo scanning



Cargo containers scanned at ports and border crossings

Accelerator-based sources of X-Rays can be far more penetrating (6MV) than Co-60 sources.

Container must be scanned in 30 seconds.

Image source: Varian medical systems




Image: dutch.euro

37

Materials testing for fusion

Source: IFMIF.org

“deuterium-tritium nuclear fusion reactions will generate neutron fluxes in the order of $10^{18} \text{ m}^{-2}\text{s}^{-1}$ with an energy of 14.1 MeV that will collide with the first wall of the reactor vessel”

International Fusion Material Irradiation Facility (IFMIF)

40 MeV
2 x 125mA linacs
CW deuterons, 5MW each
Beams will overlap onto a liquid Li jet
To create conditions similar to in a fusion reactor

To de-risk IFMIF, first a test accelerator ‘LIPAc’ is being built




Installation of ‘LIPAc’ test accelerator has started in Japan

38

Accelerator Driven Systems




Transmutation of nuclear waste isotopes or energy generation



Major challenges for accelerator technology in terms of beam power (>10MW) and reliability

6. Historical and cultural applications


Radiocarbon Dating



For more accuracy, isolate C-14 from other isotopes
 "AMS" = Accelerator Mass Spectrometry

41


Accelerators can study art




Patch of Grass, spring 1887, F583/JH1263, KM 105.264 (30,8 x 39,7 cm), Kröller-Müller Museum (Photo: Rik Klein Gotink)

This painting "Patch of grass" by Vincent van Gogh was the first one analysed by a particle accelerator

Used X-ray fluorescence technique
 Distribution of Hg and Sb pigment allowed a reconstruction of underlying image



It showed a portrait of a woman underneath




http://www.javno.com/en-best-seller/van-gogh-first-victim-of-particle-bombarding_185316

Accelerators can help spot art forgeries

Ion Beam Analysis shows us the chemical composition of pigments used in paint

This allows art historians to compare them with paints available to artists like Leonardo da Vinci



Accelerators in archaeology

The interior of samples can be studied using accelerators without destroying them

Pottery from Armenia, dating back to 1300 BC, is set up for a synchrotron experiment




Image: Argonne National Laboratory


Accelerators can make food taste better

NEW INSIGHTS INTO
CHOCOLATE



Of the six possible crystal forms, the fifth (form V) produces the best quality chocolate

Cadbury used X-rays from a particle accelerator to study how cocoa crystallises



Finally, just one more application...

Detecting wine fraud

Use ion beam to test the bottle of “antique” wine – chemical composition of the bottle compared to a real one.

“In a recent and spectacular case, American collector William Koch sued a German wine dealer, claiming four bottles – allegedly belonging to former U.S. president Thomas Jefferson – purchased for 500,000 dollars, were fake. The case has yet to be settled.”

- <http://www.cosmosmagazine.com>

HOME » NEWS » SCIENCE » SCIENCE NEWS

Atomic boffins spot fake wines

 0  0  0  0  0  Email

Bottles are zapped with beams of charged ions generated by a particle accelerator

Bottles are zapped with beams of charged ions generated by a particle accelerator

 By Roger Highfield, Science Editor
7:25PM BST 03 Sep 2008

A rare wine merchant has joined forces with nuclear scientists to develop a 21st-century tool for unmasking counterfeit vintage wines.




Next time someone asks you what accelerators are for...

*“A beam of the right particles with the right energy at the right intensity can shrink a tumor, produce cleaner energy, spot suspicious cargo, make a better radial tire, clean up dirty drinking water, map a protein, study a nuclear explosion, design a new drug, make a heat-resistant automotive cable, diagnose a disease, reduce nuclear waste, detect an art forgery, implant ions in a semiconductor, prospect for oil, date an archaeological find, package a Thanksgiving turkey or...
...discover the secrets of the universe.”*




-Accelerators for Americas Future Report, pp. 4, DoE, USA, 2011

Accelerators in the world >24000



- High Energy Accelerators of more than 1 GeV
- Ion implantation
- Electron cutting and welding
- Electron beam and X-ray irradiators (sterilization)
- Ion Beam analysis (including AMS)
- Radioisotope production (including PET)
- Non destructive testing (including security)
- Neutron generators (including sealed tubes)
- Synchrotron radiation



Engines of Discovery.
A Century of Particle Accelerators.
Andrew Sessler, Edmund Wilson

Source (2007):
<http://www.worldscientific.com/worldscibooks/10.1142/6272>