



Contribution ID: 39

Type: **Talk**

## **Initial source size and transverse momentum fluctuations in the event-by-event relativistic heavy-ion collisions**

*Saturday, 17 October 2009 10:00 (30 minutes)*

We investigate the initial transverse size of the source, which comes directly from the Glauber treatment of the earliest stage of relativistic heavy-ion collisions. After the hydrodynamic evolution stage the fluctuations in the transverse velocity flow at the hadronic freeze-out are transformed into the even-by-event fluctuations of the average transverse momentum. The Glauber phase is simulated by GLISSANDO and followed by a realistic hydrodynamic evolution stage. The statistical hadronization is performed by the THERMINATOR. We describe the  $p_T$  fluctuations at RHIC, in particular the magnitude of the effect, its centrality dependence, and the weak dependence on the incident energy. The results show that the observed event-by-event  $p_T$  fluctuations are mainly caused by the initial source size fluctuations

**Primary author:** CHOJNACKI, Mikołaj (Institute of Nuclear Physics, Krakow - Poland)

**Presenter:** CHOJNACKI, Mikołaj (Institute of Nuclear Physics, Krakow - Poland)

**Session Classification:** LHC predictions

**Track Classification:** Expectations for the Observations at the LHC: Theory & Experiment