

Non-Abelian Vortex in Four Dimensions as a Critical String on a Conifold

P. Koroteev, M. Shifman and A. Yung

1 Introduction

Shifman and Yung, 2015:

Non-Abelian vortex in $\mathcal{N} = 2$ supersymmetric QCD can behave as a critical superstring

4D $\mathcal{N} = 2$ QCD: $U(N = 2)$ gauge group and $N_f = 4$ quark flavors.

Non-Abelian vortex strings

Non-Abelian strings were suggested in $\mathcal{N} = 2$ $U(N)$ QCD

Hanany, Tong 2003

Auzzi, Bolognesi, Evslin, Konishi, Yung 2003

Shifman Yung 2004


Hanany Tong 2004

Z_N Abelian string: Flux directed in the Cartan subalgebra, say for
 $SO(3) = SU(2)/Z_2$

$$flux \sim \tau_3$$

Non-Abelian string : **Orientational zero modes**

Rotation of color flux inside $SU(N)$.



Idea:

Non-Abelian string has more moduli than Abrikosov-Nielsen-Olesen (ANO) string.

It has translational + orientational moduli

We can fulfill the criticality condition:

- The solitonic non-Abelian vortex has six orientational moduli, which, together with four translational moduli, form a ten-dimensional space ($N = 2, N_f = 4$).
- For $N_f = 2N$ 2D world sheet theory on the string is conformal.

Most of solitonic strings are "thick".

Transverse size = $\frac{1}{m}$, where m is the typical mass of bulk excitations.

ANO string: Nambu-Goto action

$$S_{\text{NG}} = T \int d^2\sigma \left\{ \sqrt{h} + O\left(\frac{\partial^n}{m^n}\right) \right\}$$

where T is string tension and

$$h = \det(\partial_\alpha x^\mu \partial_\beta x_\mu)$$

Polchinski-Strominger, 1991: Without higher derivative terms
the world sheet theory is not UV complete

Higher derivative terms at weak coupling, $g \ll 1$

$$O\left(\frac{\partial^n}{m^n}\right), \quad m \sim g\sqrt{T}$$

At $J \sim 1$ $\partial \rightarrow \sqrt{T}$


Thus higher derivative terms

$$\rightarrow \left(\frac{T}{m^2}\right)^n$$

blow up at weak coupling!

Polyakov: string surface become "crumpled".

String grows short and thick.



QUESTION:

Can we find **any** example of a 4D field theory which supports **thin** vortex strings?

Non-Abelian vortex in $\mathcal{N} = 2$ QCD with $U(2)$ gauge group and $N_f = 4$ flavors is critical.

Given that for non-Abelian vortex low energy world sheet theory is critical

Shifman, Yung 2015: Conjecture that

Thin string regime

$$T \ll m^2$$

is actually satisfied at strong coupling $g_c^2 \sim 1$.

$$m(g) \rightarrow \infty, \quad g^2 \rightarrow g_c^2$$

Higher derivative corrections can be ignored

2 Non-Abelian vortex strings

Bulk theory: 4D $\mathcal{N} = 2$ QCD with Fayet-Iliopoulos term.

For $U(N)$ gauge group in the bulk we have 2D $CP(N - 1)$ model on the string

$CP(N - 1) \implies U(1)$ gauge theory in the strong coupling limit

$$S_{CP(N-1)} = \int d^2x \left\{ |\nabla_\alpha n^P|^2 + \frac{e^2}{2} (|n^P|^2 - \beta)^2 \right\},$$

where n^P are complex fields $P = 1, \dots, N$,

Condition

$$|n^P|^2 = \beta = \frac{4\pi}{g^2},$$

imposed in the limit $e^2 \rightarrow \infty$

More flavors \Rightarrow semilocal non-Abelian string

The orientational moduli described by a complex vector n^P (here $P = 1, \dots, N$),

$\tilde{N} = (N_f - N)$ size moduli are parametrized by a complex vector ρ^K ($K = N + 1, \dots, N_f$).

The effective two-dimensional theory is the $\mathcal{N} = (2, 2)$ weighted CP model

$$S_{\text{WCP}} = \int d^2x \left\{ |\nabla_\alpha n^P|^2 + |\tilde{\nabla}_\alpha \rho^K|^2 + \frac{e^2}{2} (|n^P|^2 - |\rho^K|^2 - \beta)^2 \right\},$$

$$P = 1, \dots, N, \quad K = N + 1, \dots, N_f.$$

The fields n^P and ρ^K have charges $+1$ and -1 with respect to the auxiliary $U(1)$ gauge field

$$e^2 \rightarrow \infty$$

3 From non-Abelian vortices to critical strings

String theory

$$\begin{aligned} S &= \frac{T}{2} \int d^2\sigma \sqrt{h} h^{\alpha\beta} \partial_\alpha x^\mu \partial_\beta x_\mu \\ &+ \int d^2\sigma \sqrt{h} \left\{ h^{\alpha\beta} \left(\tilde{\nabla}_\alpha \bar{n}_P \nabla_\beta n^P + \nabla_\alpha \bar{\rho}_K \tilde{\nabla}_\beta \rho^K \right) \right. \\ &\left. + \frac{e^2}{2} \left(|n^P|^2 - |\rho^K|^2 - \beta \right)^2 \right\} + \text{fermions}, \end{aligned}$$

where $h^{\alpha\beta}$ is the world sheet metric. It is independent variable in the Polyakov formulation.

What about necessary conditions for thin string?

- Conformal invariance

$$b_{WCP} = N - \tilde{N} = 0 \Rightarrow N = \tilde{N}, \quad N_f = 2N$$

- Critical dimension =10

Number of orientational + size degrees of freedom
 $= 2(N + \tilde{N} - 1) = 2(2N - 1)$

$$4 + 2(2N - 1) = 4 + 6 = 10, \quad \text{for } N = 2$$

Our string is BPS so we have $\mathcal{N} = (2, 2)$ supersymmetry on the world sheet.

For these values of N and \tilde{N} the target space of the weighted $CP(2, 2)$ model is a non-compact Calabi-Yau manifold studied by Witten and Vafa, namely conifold.

Strings in the $U(N)$ theories are stable; they cannot be broken. Thus, we deal with the closed string.

For closed string moving on Calabi-Yau manifold $\mathcal{N} = (2, 2)$ world sheet supersymmetry ensures $\mathcal{N} = 2$ supersymmetry in 4D.

This is expected since we started with 4D QCD with $\mathcal{N} = 2$ supersymmetry.

Type IIB string is a chiral theory and breaks parity while Type IIA string theory is left-right symmetric and conserves parity.

Our bulk theory conserves parity \Rightarrow we have **Type IIA superstring**

There is self-duality in 4D bulk theory

$$\tau \rightarrow \tau_D = -\frac{1}{\tau}, \quad \tau = \frac{4\pi i}{g^2} + \frac{\theta_{4D}}{2\pi},$$

We conjectured that the string becomes thin at $g^2 \rightarrow g_c^2 \sim 1$.

It is natural to expect that $g_c^2 = 4\pi =$ self-dual point.

$$m^2 \rightarrow T \times \begin{cases} g^2, & g^2 \ll 1 \\ \infty, & g^2 \rightarrow 4\pi \\ 16\pi^2/g^2, & g^2 \gg 1 \end{cases},$$

In 2D theory on the string self-dual point is $\beta = 0$

Conifold develops conical singularity.

4 4D Graviton

Our goal:

Study massless states of closed string propagating on

$$R_4 \times Y_6, \quad Y_6 = \text{conifold}$$

and interpret them as hadrons in 4D $\mathcal{N} = 2$ QCD.

Massless 10D graviton

$$\delta G_{\mu\nu} = \delta g_{\mu\nu}(x) g_6(y), \quad \delta G_{\mu i} = B_\mu(x) g_i(y), \quad \delta G_{ij} = g_4(x) \delta g_{ij}(y)$$

Lichnerowicz equation

$$(\partial_\mu \partial^\mu + \Delta_6) g_4(x) g_6(y) = 0$$

Expand $g_6(y)$ in eigenfunctions

$$-\Delta_6 g_6(y) = \lambda_6 g_6(y), \quad \lambda_6 = \text{mass}$$

Consider massless states $\lambda_6 = 0$

$$-\Delta_6 g_6(y) = 0.$$

Solutions of this equation for Calabi-Yau manifolds are given by elements of Dolbeault cohomology $H^{(p,q)}(Y_6)$, where (p, q) denotes numbers of holomorphic and anti-holomorphic indices in the form. The dimensions of these spaces $h^{(p,q)}$ are called Hodge numbers for a given Y_6 .

For 4D graviton $g_6(y)$ is scalar

$$-D_i \partial^i g_6 = 0$$

The only solution is

$$g_6(y) = \text{const}$$

Non-normalizable on non-compact conifold Y_6 .

No 4D graviton == good news!

We do not have gravity in our 4D $\mathcal{N} = 2$ QCD

5 Kahler form deformations

Consider 4D scalar fields

Lichnerowicz equation on Y_6

$$D_k D^k \delta g_{ij} + 2R_{ikjl} \delta g^{kl} = 0.$$

Solutions = Kahler form deformations or complex structure deformations.

Kahler form deformations = variations of 2D coupling β

D -term condition in weighted CP(2,2) model

$$|n^P|^2 - |\rho^K|^2 = \beta, \quad P = 1, 2, \quad K = 1, 2$$

Resolved conifold

The effective action for $\beta(x)$ is

$$S(\beta) = T \int d^4x h_\beta (\partial_\mu \beta)^2,$$

where

$$h_\beta = \int d^6y \sqrt{g} g^{li} \left(\frac{\partial}{\partial \beta} g_{ij} \right) g^{jk} \left(\frac{\partial}{\partial \beta} g_{kl} \right)$$

Using explicit Calabi-Yau metric on resolved conifold we get

$$h_\beta = (4\pi)^3 \frac{5}{6} \int dr r = \infty$$

β - non-normalizable mode

Physical nature of non-normalizable modes

Gukov, Vafa, Witten 1999: Non-normalizable moduli = coupling constants in 4D

- 4D metric do not fluctuate. It is fixed to be flat. "Coupling constants."
- 2D coupling β is related to 4D coupling g^2 . Fixed. Non-dynamical.

Another option:

Large $y_i \Rightarrow$ large n^P and ρ^K

Non-normalizable modes are not localized on the string.

Unstable states. Decay into massless perturbative states.

Higgs branch: $\dim \mathcal{H} = 4N\tilde{N} = 16$.

6 Deformation of the complex structure

D -term condition

$$|n^P|^2 - |\rho^K|^2 = \beta, \quad P = 1, 2, \quad K = 1, 2$$

Construct U(1) gauge invariant "mesonic" variables"

$$w^{PK} = n^P \rho^K.$$

$$\det w^{PK} = 0$$

Take $\beta = 0$

Complex structure deformation \Rightarrow Deformed conifold

$$\det w^{PK} = b$$

b – complex modulus

The effective action for $b(x)$ is

$$S(\beta) = T \int d^4x h_b (\partial_\mu b)^2,$$

where

$$h_b = \int d^6y \sqrt{g} g^{li} \left(\frac{\partial}{\partial b} g_{ij} \right) g^{jk} \left(\frac{\partial}{\partial \bar{b}} g_{kl} \right)$$

Using explicit Calabi-Yau metric on deformed conifold we get

$$h_b = (4\pi)^3 \frac{4}{3} \log \frac{T^2 L^4}{|b|}$$

For Type IIA string b is a part of hypermultiplet.

Another complex scalar \tilde{b} comes from 10D 3-form.

$$S(b) = T \int d^4x \left\{ |\partial_\mu b|^2 + |\partial_\mu \tilde{b}|^2 \right\} \log \frac{T^4 L^8}{|b|^2 + |\tilde{b}|^2}$$


7 Monopole-monopole baryon

Weak coupling


Strings in the $U(N)$ theories are stable; they cannot be broken. Thus, we deal with the closed string.

Quarks are condensed in the bulk theory. Therefore, monopoles are confined.

In $U(N)$ gauge theories the confined monopoles are junctions of two non-Abelian vortex strings.



a



b

Monopole-antimonopole meson

Monopole-monopole baryon

Stringy states are massive, with mass $\sim \sqrt{T}$.

Strong coupling

Global group of the 4D QCD:

$$SU(2) \times SU(2) \times U(1)$$

$U(1)$ - "baryonic" symmetry.

b -hypermultiplet: $(1, 1, 2)$

Logarithmically divergent norm \implies Marginal stability at $\beta = 0$

b -state can decay into massless bi-fundamental (screened) quarks living on the Higgs branch.

8 Conclusions

- In $\mathcal{N} = 2$ supersymmetric QCD with gauge group $U(2)$ and $N_f = 4$ quark flavors non-Abelian BPS vortex behaves as a critical fundamental superstring.
- Massless closed string state b associated with deformations of the complex structure of the conifold \implies monopole-monopole baryon.
- Successful tests of our gauge-string duality:
 - $\mathcal{N} = 2$ supersymmetry in 4D QCD
 - Absence of graviton and unwanted vector fields.
 - Massless monopole-monopole baryon is present only at $\beta = 0$ and cannot be continued to weak coupling.