

Double Higgs Production in VBF

Fady Bishara



17 January 2017




1st FCC Physics Workshop

Based on: 1611.03860 FB, R. Contino, and J. Rojo

Motivation


- Is EWSB in the SM minimal?
- Is EWSB linearly realised?
- If EWSB is non-linearly realised, can we test this?
- Non-linearities \rightarrow harder mhh tail \rightarrow handle on the background

Double Higgs production



HH production at pp colliders

- VBF cross-section at the LHC is small ~ 2 [fb] w/o \mathcal{BR} s (100[fb] at the FCC)
- But, is a unique probe of the EWSB mechanism



Is EWSB (non-)linearly realised?

[Grinstein and Trott: [0704.1505]

[Contino, Grojean, Moretti, Piccinini, Rattazzi: 1002.1011]

$$\Sigma = e^{i\sigma^a \pi^a / v}$$

$$\mathcal{L} \supset \frac{1}{2} (\partial_\mu h)^2 - V(h)$$

$$+ \frac{v^2}{4} \text{Tr} (D_\mu \Sigma^\dagger D^\mu \Sigma) \left[1 + 2c_V \frac{h}{v} + c_{2V} \frac{h^2}{v^2} + \dots \right]$$

$$- m_i \bar{\psi}_{Li} \Sigma \left(1 + c_\psi \frac{h}{v} + \dots \right) \psi_{Ri} + \text{h.c.},$$

$$V(h) = \frac{1}{2} m_h^2 h^2 + c_3 \frac{1}{6} \left(\frac{3m_h^2}{v} \right) h^3 + c_4 \frac{1}{24} \left(\frac{3m_h^2}{v^2} \right) h^4 + \dots$$

- In the minimal SM, linear realization
→ $c_V = c_{2V} = c_3 = 1$ and all ... terms vanish
- Measuring $c_{2V} \neq 1 \rightarrow$ non-linearity!

A concrete example

- In minimal $SO(5)/SO(4)$ models, the couplings c_V and c_{2V} are given by Agashe et al. [hep-ph/0412089]
Contino et al. [hep-ph/0612048]

$$c_V = \sqrt{1 - \xi}, \quad c_{2V} = 1 - 2\xi$$

where $\xi = v^2/f^2$

- And, looking at the longitudinal vector boson scattering we see that

$$\mathcal{A}(V_L V_L \rightarrow hh) \simeq \frac{\hat{s}}{v^2} (c_{2V} - c_V^2)$$

- Choose a benchmark with $c_{2V} = 0.8$ (to roughly correspond to $\xi = 0.1$ which is at the boundary of exclusion by ATLAS) ATLAS: [1509.00672]


Sensitivity to δ_{c_3} and $\delta_{c_{2V}}$

- To illustrate, consider total σ before and after cuts with $\sigma/\sigma_{\text{SM}} = 1 + a\delta + b\delta^2$
- Sensitivity to $\delta_{c_{2V}}$ (δ_{c_3}) is enhanced (suppressed) by the cuts

$$\delta_i \equiv c_i - 1$$

dashed : before cuts

solid : after cuts



Kinematic cuts & b-tagging

Final cuts:

		14 TeV	100 TeV
Acceptance cuts	p_{T_j} (GeV) \geq	25	40
	p_{T_b} (GeV) \geq	25	35
	$ \eta_j \leq$	4.5	6.5
	$ \eta_b \leq$	2.5	3.0
VBF cuts	$ \Delta y_{jj} \geq$	5.0	5.0
	m_{jj} (GeV) \geq	700	1000
	Central jet veto: $p_{T_{j_3}}$ (GeV) \leq	45	65
	m_{hh} (GeV) \geq	500	1000


B-tagging parameters:

$$\varepsilon(b\text{-tag}) = 0.75, \quad \varepsilon(c\text{-mistag}) = 0.1, \quad \varepsilon(q, g\text{-mistag}) = 0.01$$

Scale invariant tagging


Gouzevitch et al. [1303.6636]

- Key feature: m_{hh} tail is harder when $c_V^2 \neq c_{2V}$
- Signal events will have boosted Higgs pairs \rightarrow handle to reduce background



Higgs reconstruction

Resolved



Boosted








Figure credit: Juan Rojo

- ▷ 4 small- R b-tagged jets
- ▷ Consider hardest 6
- ▷ $h_1 \leftrightarrow$ b-jet pair with $\min\{|m_{bb} - 125|\}$
- ▷ $h_2 \leftrightarrow$ b-jet pair with $\min\{|m_{bb} - m_{h_1}|\}$


- ▷ 2 large- R jet \supset 2 b-quarks each
- ▷ $h_1 \leftrightarrow$ large- R jet with $\min\{|m_j - 125|\}$
- ▷ $h_2 \leftrightarrow$ large- R with $\min\{|m_{j_2} - m_{j_1}|\}$

Taming the background




Detector coverage at an FCC

- VBF jet with max η peaks ~ 5
- If coverage only extended to $|\eta| < 5$, would lose $\sim 50\%$ of signal events






The hh invariant mass distribution

14 TeV





100 TeV



Results: probability intervals on δc_{2V}

$$\delta c_{2V} \equiv c_{2V} - 1$$




	68% probability interval on δc_{2V}	
	$1 \times \sigma_{\text{bkg}}$	$3 \times \sigma_{\text{bkg}}$
LHC ₁₄	[-0.37, 0.45]	[-0.43, 0.48]
HL-LHC	[-0.15, 0.19]	[-0.18, 0.20]
FCC ₁₀₀	[0, 0.01]	[-0.01, 0.01]

	95% probability upper limit on μ	
	$1 \times \sigma_{\text{bkg}}$	$3 \times \sigma_{\text{bkg}}$
LHC ₁₄	109	210
HL-LHC	49	108
FCC ₁₀₀	12	23

Validity of the EFT description

- If NP is characterized by coupling g_* and scale Λ
- One expects $\delta_{c_{2V}} \approx g_*^2 v^2 / \Lambda^2$ See, e.g., [Giudice, Grojean, Pomarol, Rattazzi: hep-ph/0703164]
- Saturating the strong coupling limit then gives
 $\delta_{c_{2V}} \approx 16\pi^2 v^2 / \Lambda^2$

This procedure was outlined in [Contino, Falkowski, Goertz, Grojean, Riva: 1604.06444]







Summary & outlook

- Double Higgs production in VBF is useful to constrain $hhVV$ couplings
- Boosted kinematics gives a crucial handle to tame backgrounds
- 20% precision achievable at the HL-LHC reaching the 1% level at a 100 TeV FCC


Thank you!

Scale invariant tagging



Populating the tail in our analysis

- Sensitivity is driven by the tail. Therefore good modelling is imperative



$$d\sigma/db_j / (\sum_i d\sigma/db_i) \sim 10^{-6}$$

→ need 1M events to get 1 event in this bin.

→ Accounting for efficiency of all cuts and requiring 100 events here means need to generate 10^{12} events!

- Solution: generate weighted events and fit the background
- For signal, can also put generation cut on m_{hh} but this does not work for background