

Moduli of heterotic G2 systems

Magdalena Larfors

Uppsala University

COST MP1210 Conference & 22nd European Workshop on String Theory
23.02.2017

X. de la Ossa, ML, E. Svanes (1607.03473 & work in progress)

Motivation and summary

Heterotic string compactifications: long history of particle physics models

- Minkowski 4D $\mathcal{N} = 1$ vacua:
 - Calabi–Yau manifolds with vector bundles
 - Strominger–Hull systems with flux and bundles.
- Moduli stabilisation? No RR flux.
- How many moduli?

Moduli of these configurations needed to determine the low-energy physics.

This talk: heterotic compactifications on G_2 structure manifolds

- Heterotic 4D $\mathcal{N} = 1/2$ domain wall solutions
(may “uplift” non-perturbatively to non-SUSY *AdS* solutions.)
- Infinitesimal moduli of heterotic systems with G_2 holonomy/structure
(some relevance also for (non-SUSY) M-theory and type II compact.)
- Moduli of heterotic system captured by Atiyah-like bundles.

Heterotic supersymmetric vacua

Heterotic string to $\mathcal{O}(\alpha')$

- Bosonic fields: Metric G , B-field B , dilaton ϕ , gauge field A
- Fermionic fields: Gravitino Ψ_M , dilatino λ , gaugino χ

Compactifications

- $\mathcal{M}_{10} = \mathcal{M}_E \times X$: SUSY \iff nowhere vanishing spinor η on X
- Killing spinor equations

$$\begin{aligned}\nabla_H \eta &= \left(\nabla_M + \frac{1}{8} \not{H}_M \right) \eta = 0 \\ \left(\not{\nabla} \hat{\phi} + \frac{1}{12} \not{H} \right) \eta &= 0 \\ \not{F} \eta &= 0\end{aligned}$$

where $\not{\nabla} = \gamma^M \nabla_M$, etc.

- Bianchi identity $dH = \frac{\alpha'}{4} (\text{tr} R \wedge R - \text{tr} F \wedge F)$

4D Heterotic $\mathcal{N} = 1$ Minkowski vacua

6D Geometry:

- $\nabla_H \eta = 0 \iff X$ has conformally balanced $SU(3)$ structure
 $\iff d(e^{-2\phi}\Psi) = 0 = d(e^{-2\phi}\omega \wedge \omega)$.
 - No H -flux $\iff X$ is Calabi–Yau $d\Psi = 0 = d\omega$.
- Candelas, et.al.:85, Hull:86; Strominger:86, Ivanov, Papadopoulos:00; Gauntlett, et.al.:03,...*

Gauge fields \rightarrow vector bundle V


- $\not{F}\eta = 0 \implies$
 - ▶ $F^{(0,2)} = F^{(2,0)} = 0 \rightsquigarrow$ holomorphic V
 - ▶ HYM equation $F \lrcorner \omega = 0 \rightsquigarrow$ polystable holomorphic V
- Candelas, et.al.:85, Donaldson:85, Uhlenbeck, Yau:86, Li, Yau:87,...*

4D Heterotic $\mathcal{N} = 1/2$ DW vacua

Lukas et al:10, 11, 12, 13; Gray, ML, Lüst:12, de la Ossa, ML, Svanes:14,16

7D Geometry

4D domain wall vacuum: $\mathcal{M}_{10} = \mathcal{M}_4 \times_W X(r) \equiv \mathcal{M}_3 \times Y$
 $\mathcal{M}_4 = \mathcal{M}_3 \times \mathbb{R}$, \mathcal{M}_3 AdS or Minkowski



4D Heterotic $\mathcal{N} = 1/2$ DW vacua

Lukas et al:10, 11, 12, 13; Gray, ML, Lüst:12, de la Ossa, ML, Svanes:14,16

7D Geometry:

- $\nabla_H \eta = 0 \iff Y$ has integrable G_2 structure

$$\iff \boxed{d(e^{-2\phi}\psi) = 0, \quad d\phi = \tau_0 \psi + \frac{3}{2} d\phi \wedge \varphi + *\tau_3.}$$

for positive 3-form φ and $\psi = *\varphi$.

- No H -flux $\iff Y$ has G_2 holonomy

Embed $SU(3)$ in G_2 : $\varphi = dr \wedge \omega(r) + \text{Re}(\Psi(r))$.

G_2 torsion classes: Fernandez-Gray:82, Chiossi-Salamon:02

Gauge fields \rightarrow vector bundle V

- $\not{F}\eta = 0 \implies$ instanton bundle: $\boxed{F \wedge \psi = 0.}$

Infinitesimal Moduli of Heterotic SUSY Vacua

- 4D $\mathcal{N} = 1$ vacua: deformations of (X, V, H)
 - ▶ X : conformally balanced complex 3-fold
 - ▶ V : holomorphic polystable gauge bundle
 - ▶ H : α' corrected BI
- “Atiyah class stabilization”

Infinitesimal moduli

=

Deformations of holomorphic structure on extension bundle of Atiyah type.

Atiyah:57, Kodaira, Spencer:58,60, Candelas, de la Ossa:91, Becker,et.al:05,06, Anderson,et.al:10,11,13, Fu, Yau:11,Anderson, Gray, Sharpe:14, de la Ossa, Svanes:14, Garcia-Fernandez,et.al:13,15,...

- 4D $\mathcal{N} = 1/2$ vacua: deformations of (Y, V, H)
 - ▶ Y integrable G_2 structure manifold
 - ▶ V : instanton gauge bundle
 - ▶ H : α' corrected BI

de la Ossa, ML, Svanes:16 + in progress, Clarke,et.al:16

Infinitesimal Moduli: Heterotic G_2 systems

Naive moduli space

Y : integrable G_2 structure manifold ($H = 0$: G_2 holonomy)

V : instanton gauge bundle

- $\partial_t \psi, \partial_t \varphi$: geometric moduli

Joyce:96, Dai–Wang–Wei:03, de Boer–Naqvi–Shomer:05,...

- $\partial_t A$: Vector bundle moduli $H^1(Y, \text{End}(V))$
- $\partial_t B$: deformations of B -field, $H = dB + \alpha'(\dots)$

Infinitesimal Moduli: Heterotic G_2 holonomy system

Geometric moduli for G_2 holonomy

$$\partial_t \psi = \frac{1}{3!} M_t^a \wedge \psi_{bcda} dx^{bcd}, \quad M_t^a = M_t b^a dx^b$$

$$\partial_t \varphi = -\frac{1}{2} M_t^a \wedge \varphi_{bca} dx^{bc}$$

- Diffeomorphisms:

$$\mathcal{L}_V \psi = -\frac{1}{3!} (d_\theta V^a) \wedge \psi_{bcda} dx^{bcd}$$

where d_θ is a connection for TY -valued forms.

- Preserve $d\psi = 0 = d\varphi$: constraints on $d_\theta M_t$

- Compact G_2 manifold:

$$\mathcal{T}\mathcal{M}_{G_2\text{hol}} \cong H_d^3(Y) \subset H_{d_\theta}^1(Y, TY)$$

- Canonical G_2 cohomology (integrable G_2 str.)

Reyes-Carrion:93, Fernandez-Ugarte:98

Infinitesimal Moduli: Heterotic G_2 holonomy system

G_2 “Atiyah” class stabilization

Instanton condition $F \wedge \psi = 0$: couples bundle and geometric moduli

$$\check{d}_A(\partial_t A) = -\check{\mathcal{F}}(\Delta_t) .$$

$\Delta_t \in \Lambda^1(Y, TY)$, \check{d}_A , $\check{\mathcal{F}}$: project to G_2 $\mathbf{7}$ irrep

- “Atiyah” map

$$\begin{aligned} \mathcal{F} : \Lambda^p(Y, TY) &\longrightarrow \Lambda^{p+1}(Y, \text{End}(V)) \\ \Delta &\mapsto \mathcal{F}(\Delta) = -F_{ab} dx^b \wedge \Delta^a . \end{aligned}$$

Bianchi identity $d_A F = 0 \implies \check{\mathcal{F}}$ is a map in cohomology

Corrected moduli space for bundle and geometric moduli:

$$\mathcal{T}\mathcal{M}_{(Y,V)} \subset H_{\check{d}_A}^1(\text{End}(V)) \oplus \ker(\check{\mathcal{F}})$$

Infinitesimal Moduli: Heterotic G_2 holonomy system

Infinitesimal moduli space for bundle and geometry:

$$\mathcal{T}\mathcal{M}_{(Y,V)} \subset H_{d_A}^1(\text{End}(V)) \oplus \ker(\check{\mathcal{F}})$$

Remark 1: B -field deformations

- Infinitesimal moduli of G_2 -holonomy metrics only spans part of $H_{d_\theta}^1(Y, TY)$:

$$H_{d_\theta}^1(Y, TY) \cong \check{\mathcal{H}}^1(Y, TY) = \check{\mathcal{S}}^1(Y, TY) \oplus \check{\mathcal{A}}^1(Y, TY)$$

- $\check{\mathcal{A}}^1(Y, TY)$ is spanned by $\partial_t B$
- All $\partial_t B$ are in the kernel of $\check{\mathcal{F}}$.
- Thus easily incorporate B -field deformations in the infinitesimal moduli space.

Infinitesimal Moduli: Heterotic G_2 holonomy system

Infinitesimal moduli space for bundle, geometry and B-field:

$$\mathcal{TM}_{(Y,V,B)} \cong H_{d_A}^1(\text{End}(V)) \oplus \ker(\check{\mathcal{F}})$$

Remark 2: Extension bundle

- Use the G_2 Atiyah map $\check{\mathcal{F}}$ to define a new bundle

$$0 \longrightarrow \text{End}(V) \longrightarrow E \longrightarrow TY \longrightarrow 0,$$

- E has connection \mathcal{D}_E :

$$\mathcal{D}_E = \begin{pmatrix} \check{d}_A & \check{\mathcal{F}} \\ 0 & \check{d}_\theta \end{pmatrix}.$$

- $\mathcal{D}_E^2 = 0 \iff \check{\mathcal{F}}(\check{d}_\theta(\Delta)) + \check{d}_A(\check{\mathcal{F}}(\Delta)) = 0$ (cf Atiyah algebroid for $\mathcal{N} = 1$).
- $H_{\mathcal{D}_E}^1(Y, E)$ is the moduli space:

$$0 \rightarrow H_{d_A}^1(Y, \text{End}(V)) \rightarrow H_{\mathcal{D}_E}^1(E) \rightarrow H_{d_\theta}^1(Y, TY) \xrightarrow{\check{\mathcal{F}}} H_{d_A}^2(Y, \text{End}(V)) \rightarrow \dots$$

Infinitesimal Moduli: Heterotic integrable G_2 system

Geometric moduli for integrable G_2 structure

- Diffeomorphisms:

$$\mathcal{L}_V \psi = -\frac{1}{3!} (d_\theta V^a) \wedge \psi_{bcda} dx^{bcd}$$

where d_θ is a connection for TY -valued forms.

- Preserve $\tau_2 = 0$:

$$(d_\theta \check{\Delta}_t^a) \wedge \psi_{bcda} dx^{bcd} = 0$$

- $\partial_t \varphi \implies$ Variational constraints on torsion

$\check{\mathcal{F}}, \check{\mathcal{R}}$ maps

SUSY + BI \implies EOM if θ is an instanton connection: $R(\theta) \wedge \psi = 0$.

- \mathcal{R} map: completely analogous to \mathcal{F} .
- Extra moduli for connection variations.
Related to field redefinitions as for $\mathcal{N} = 1$ system?
- $\check{\mathcal{F}}, \check{\mathcal{R}}$ in fact map **all** geometric moduli to $\check{d}_A, \check{d}_\theta$ -closed forms

Infinitesimal Moduli: Heterotic integrable G_2 system

Corrected moduli space for bundle, instanton and geometric moduli:

$$H_{d_A}^1(\text{End}(V)) \oplus H_{d_\theta}^1(\text{End}(TY)) \oplus \ker(\check{F} + \check{R})$$

Last equation to bring in: Bianchi identity $dH = \frac{\alpha'}{4}(\text{tr}R \wedge R - \text{tr}F \wedge F)$

- Recall from $\mathcal{N} = 1$: *Anderson, et.al:14, de la Ossa, Svanes:14*
Vary BI together with $H = d^c \omega \rightsquigarrow \text{map } \mathcal{H} : \Lambda^p(X, E) \longrightarrow \Lambda^{p+1}(X, T^*X)$
 - ▶ \mathcal{H} well-defined in cohomology
 - ▶ finite-dim moduli space

- $\mathcal{N} = 1/2$ *de la Ossa, ML, Svanes:17XX*
Vary BI together with $H = \frac{1}{6} \tau_0 \varphi - \tau_1 \lrcorner \psi - \tau_3 \rightsquigarrow \text{map } \mathcal{H}$
 - ▶ \mathcal{H} well-defined in cohomology ✓
 - ▶ finite-dim moduli space ✓
 - ▶ moduli \sim deformations of Atiyah-like bundle ✓

Compare heterotic generalised geometry.

Clarke, Garcia-Fernandez, Tipler:16

Conclusions and outlook

Conclusions

- 4D heterotic $\mathcal{N} = 1/2$ DW solutions
 - ▶ Y Integrable G_2 structure $\supset G_2$ holonomy
 - ▶ X Conformally balanced (non-complex) $SU(3)$ structure
- Infinitesimal moduli of G_2 holonomy manifold Y w. instanton bundle V :

$$H_{d_A}^1(\text{End}(V)) \oplus \ker(\check{\mathcal{F}})$$

- **Infinitesimal moduli captured by $H_{D_E}^1(Y, E)$ of extension bundle**

$$0 \longrightarrow \text{End}(V) \longrightarrow E \longrightarrow TY \longrightarrow 0 ,$$

- Similar structure for moduli of heterotic int. G_2 system (Y, V, H) .

Outlook

- Relation of $SU(3)$ and G_2 structure moduli spaces for domain wall solutions.
- Relevance for deformations of M-theory and type II string compactifications.

Conclusions and outlook

Conclusions

- 4D heterotic $\mathcal{N} = 1/2$ DW solutions
 - ▶ Y Integrable G_2 structure $\supset G_2$ holonomy
 - ▶ X Conformally balanced (non-complex) $SU(3)$ structure
- Infinitesimal moduli of G_2 holonomy manifold Y w. instanton bundle V :

$$H_{d_A}^1(\text{End}(V)) \oplus \ker(\check{\mathcal{F}})$$

- **Infinitesimal moduli captured by $H_{D_E}^1(Y, E)$ of extension bundle**

$$0 \longrightarrow \text{End}(V) \longrightarrow E \longrightarrow TY \longrightarrow 0 ,$$

- Similar structure for moduli of heterotic int. G_2 system (Y, V, H) .

Outlook

- Relation of $SU(3)$ and G_2 structure moduli spaces for domain wall solutions.
- Relevance for deformations of M-theory and type II string compactifications.

Thank You

Manifolds with G_2 structure

Fernandez–Gray:82, Chiossi–Salamon:02

Decomposition of forms

$\Lambda^k(Y)$ decomposes into $\Lambda_p^k(Y)$, p denotes G_2 irrep. Find these using φ :

Example: $\Lambda^1 = \Lambda_7^1 = T^*Y \cong TY$

\implies any $\beta \in \Lambda^2$ decomposes as $\beta = \alpha \lrcorner \varphi + \gamma$, where $\alpha \in \Lambda^1$ and $\gamma \lrcorner \varphi = 0$

$$\Lambda^0 = \Lambda_1^0,$$

$$\Lambda^1 = \Lambda_7^1 = T^*Y \cong TY,$$

$$\Lambda^2 = \Lambda_7^2 \oplus \Lambda_{14}^2,$$

$$\Lambda^3 = \Lambda_1^3 \oplus \Lambda_7^3 \oplus \Lambda_{27}^3.$$

Canonical G_2 cohomology

Decomposition of de Rham cohomology

Reyes-Carrion:93, Fernandez-Ugarte:98

Analogue of Dolbeault operator on a complex manifold: project d onto G_2 irreps.

- The differential operator \check{d} is defined by

$$\check{d}_0 = d, \quad \check{d}_1 = \pi_7 \circ d, \quad \check{d}_2 = \pi_1 \circ d.$$

- $\tau_2 = 0 \iff \check{d}^2 = 0$, so can construct differential, elliptic complex

$$0 \rightarrow \Lambda^0(Y) \xrightarrow{\check{d}} \Lambda^1(Y) \xrightarrow{\check{d}} \Lambda_7^2(Y) \xrightarrow{\check{d}} \Lambda_1^3(Y) \rightarrow 0$$

- $H_{\check{d}}^*(Y)$ is “canonical G_2 -cohomology of Y ”.

This generalizes to TY -valued forms: Elliptic complex

$$0 \rightarrow \Lambda^0(TY) \xrightarrow{\check{d}_\theta} \Lambda^1(TY) \xrightarrow{\check{d}_\theta} \Lambda_7^2(TY) \xrightarrow{\check{d}_\theta} \Lambda_1^3(TY) \rightarrow 0$$

with finite-dim cohomology groups $H_{\check{d}_\theta}^p(Y, TY)$, if $R(\theta) \wedge \psi = 0$

Infinitesimal moduli: $\mathcal{N} = 1$

Impose full set of $\mathcal{N} = 1$ constraints: “Atiyah class stabilization”

- Stability $F \wedge \omega \wedge \omega$: stable under first order deformations
- Holomorphic bundle $F \wedge \Psi = 0$
- Instanton connection: $R(\nabla) \wedge \Psi = 0$
- α' corrected BI

\rightsquigarrow equivalent to deformations of the holomorphic structure on an extension bundle of Atiyah type.

Atiyah:57, Donaldson:85, Uhlenbeck, Yau:86 Li, Yau:87, Fu, Yau:11,
Anderson, Gray, Lukas, Ovrut:10,11,13, Anderson, Gray, Sharpe:14, de la Ossa, Svanes:14,
Garcia-Fernandez, *et.al*:13,15...