

Contribution ID: 4

Type: not specified

## Laser-Driven High Energy Alpha Beam Interaction with Solid p11B to Achieve Fusion Ignition by Alpha Heating

Wednesday 30 November 2016 12:00 (20 minutes)

Laser-Driven High Energy Alpha Beam Interaction with Solid p11B to Achieve Fusion Ignition by Alpha Heating

S.D. Moustaizis1, P. Lalousis2, H. Hora3, S. Eliezer4 and G. Korn 5

1 Technical University of Crete, Lab of Matter Structure and Laser Physics, Chania, Crete, Greece

2 Institute of Electronic Structure and Laser FORTH, Heraklion, Greece

3 Department of Theoretical Physics, University of New South Wales, Sydney 2052, Australia

4Nuclear Fusion institute, Polytechnique University of Madrid, ETSII, Madrid 28006, Spain

5Institute of Physics, ELI-Beamlines, CSO Prague, Czech Republic & Max-Planck-Institute for Quantum Optics,

Garching, Germany

ABSTRACT

We present for the first time numerical results on fusion ignition process produced by laser-driven high energy alpha beam interaction with compressed solid p11B fuel. Relevant results on alpha measurements from nuclear reactions, induced by laser-driven proton beam interaction with 11B plasma [1, 2], justify extensive numerical investigations on fusion burning feasibility of pB11 fuel using a multi-fluid code [3, 4, 5, 6]. The consideration of a new physical process, termed as alpha avalanche effect [7, 8] contribute to enhance the alpha heating effect and consequently the reaction rate of the fusion process. The p11B nuclear fusion reaction is attractive not only because is an aneutronic reaction but because have the advantages to produce three alphas with total energy 8.9 MeV. But the main difficulty for fusion ignition in solid density p11B targets is that the cross section for nuclear fusion reactions is efficient for energies higher than 250 keV. The proposed concept, to overcome this difficulty, is to inject an energetic alpha beam in the compressed target and simulate the temporal evolution of the temperature and the reaction rate of the p11B fuel due to alpha heating effect. The initial high energy alpha beam could be produced by ultra-short and high intensity laser beam interaction with thin DT solid targets (see fig.3 of ref.9). The high energy alpha beam interact with a p11B plasma compressed 4-10 times the initial solid density and with low initial temperaturature. The numerical results show that the maximum of the reaction rate is achieved tens of nsec after the injection of the alpha beam. After this time interval the reaction rate decreases due to the depletion of the plasma ion density. The temperature corresponding to the maximum of the reaction rate is about 200 keV. The time corresponding to the maximum reaction rate depend strongly on the density of the injected alpha beam, the initial plasma temperature and the compression factor of the p11B fuel. In the near future Petawatt or Exawatt -Zetawatt laser systems [10, 11] and especially the IZEST project and the fiber based laser system investigate for the ICAN project, will be able to attain intensities up to 1025 W/cm2 and 1029 W/cm2 with pulse duration of the order of attoseconds or zettaseconds. These projects will enable unique applications on laser-driven ion beam acceleration with high current [12, 13, 14]. The numerical results of this work promote the development of new high efficiency and high power fibre laser systems like ICAN in order to generate high density alpha beams.

[1] Labaune, C., S. Deprieraux, S. Goyon, C. Loisel, G. Yahia & J. Rafelski Nature Communications 4, 2506 (2013).
[2] G. Korn, D. Margarone, A. Picciotto, Boron-Proton Nuclear-Fusion Enhancement induced inBoron-dopped Silicon Targets by low Contrast Pulsed Laser Lecture at the IZEST conference, Paris, Romanian Embassy, 19 September 2014.

A. Picciotto et al. Physical Review X 4, 031030 (2014).

[3] P. Lalousis, H. Hora and S. Moustaizis, Laser and Particle Beams 32, 409 (2014).

[4\_H. Hora, G. Miley, P. Lalousis, S. Moustaizis, K. Clayton and D. Jonas. Efficient Generation of Fusion Flames Using PW-ps Laser Pulses for Ultrahigh Acceleration of Plasma Blocks by Nonlinear (Ponderomotive) Forces, IEEE Transaction of Plasma Science 42, 640-644 (2014).

[5] H. Hora, P. Lalousis, Shalom Eliezer, G. H. Miley, S. Moustaizis, G. Mourou. 10 kilotesla magnetic field confinement combined with ultra-fast laser accelerated plasma blocks for initiating fusion flames. Abstract of an oral presentation at the Physics-Congress Canberra/Australia, see arXiv 1412.4190. (11 December 2014).

[6]P. Lalousis, S. Moustaizis, H. Hora and G.H. Miley, Kilotesla Magnetic Assisted Fast Laser Ignited Boron-11 Hydrogen Fusion with Nonlinear Force Driven Ultrahigh Accelerated Plasma Blocks, Journal of Fusion Energy, 34 62-67 (2015).

[7]\_H. Hora et al, Petawatt laser pulses for proton-boron high gain fusion with avalanche reactions excluding problems of nuclear radiation. SPIE Conf. Proceedings No. 9515, paper 9515-44 (2015).

[8] S. Eliezer, H. Hora, G. Korn, N. Nissin and J. M. Martinez Val, "Avalance proton-boron fusion based on elastic nuclear collisions", Physics of Plasmas 23, 050704, 2016

[9]\_ R. Banati, H. Hora, P. Lalousis, S. Moustaizis.. "Ultrahigh laser acceleration of plasma blocks with ultrahigh ion density for fusion and hadron therapy", Jour. Intense Pulsed Lasers & Application in Adv. Physics 4, No. 1, 11-16 (2014).

[10]\_T. Tajima and G. Mourou, "Zettawatt-exawatt lasers and their applications in ultrastrong field physics", Physical Review Special Topics –Accelerators and Beams, Vol 5, 0310301, 2002

[11]\_G. Mourou, T. Tajima and S. Bulanov, Reviews of Modern Physics 78, 309 (2006)

[12]\_G. Mourou, T. Tajima and Jens Limpert, "The future is fibre accelerators", Nature Photonics, Vol. 7, April 2013.

[13] \_ T. Esirkepov, M. Borgesi, S. V. Bulanov, G. Mourou and T. Tajima, "High Efficiency Relativistic-Ion Generation in Laser-Piston Regime", Phys. Rev. Lett. Vol.92, no 17, 175003-2, April 2004.

[14]\_T. Tajima, G. Mourou and S. Gals, "Arrangement for generating a proton beam and an installation for transmutation of nuclear wastes", European Patent Application, EP 2 709 429 A1, 2012.

Author: MOUSTAIZIS, Stavros (Technical University of Crete)

**Presenter:** MOUSTAIZIS, Stavros (Technical University of Crete)

Session Classification: Socio-Economic Impacts

Track Classification: Socio-Economic Impacts