

# Non-exponential decay in a quantum field theoretical treatment

Francesco Giacosa  
**UJK Kielce – Goethe U Frankfurt**

**ICNFP 2017**

OAC, Kolymbari, Crete, 17-29/8/2017

# Outline



1. General discussion of the decay law
2. Non-exponential decay: experiments
3. Theory: from Lee Hamiltonian to QFT
4. Decay of a moving particle. Is the usual Einstein-formula correct?

# Part 1: General discussion

# Exponential decay law

- $N_0$  : Number of unstable particles at the time  $t = 0$ .

$$N(t) = N_0 e^{-\Gamma t}, \quad \tau = 1/\Gamma \text{ mean lifetime}$$

Confirmed in countless cases!

- For a single unstable particle:

$$p(t) = e^{-\Gamma t}$$

is the survival probability for a single unstable particle created at  $t=0$ .  
(Intrinsic probability, see Schrödinger's cat).

For small times:  $p(t) = 1 - \Gamma t + \dots$

# Basic definitions



Let  $|S\rangle$  be an unstable state prepared at  $t = 0$ .

Survival probability amplitude at  $t > 0$ :

$$a(t) = \langle S | e^{-iHt} | S \rangle$$

Survival probability:  $p(t) = |a(t)|^2$

Rep. Prog. Phys., Vol. 41, 1978. Printed in Great Britain

**Decay theory of unstable quantum systems**

L FONDA, G C GHIRARDI and A RIMINI

# Deviations from the exp. law at short times

Taylor expansion of the amplitude:

$$a(t) = \langle S | e^{-iHt} | S \rangle = 1 - it \langle S | H | S \rangle - \frac{t^2}{2} \langle S | H^2 | S \rangle + \dots$$

$$a^*(t) = \langle S | e^{iHt} | S \rangle = 1 + it \langle S | H | S \rangle - \frac{t^2}{2} \langle S | H^2 | S \rangle + \dots$$


It follows:

$$p(t) = |a(t)|^2 = a^*(t)a(t) = 1 - t^2 \left( \langle S | H^2 | S \rangle - \langle S | H | S \rangle^2 \right) + \dots = 1 - \frac{t^2}{\tau_Z^2} + \dots$$

$$\text{where } \tau_Z = \frac{1}{\sqrt{\langle S | H^2 | S \rangle - \langle S | H | S \rangle^2}} .$$

$p(t)$  decreases quadratically (not linearly);  
no exp. decay for short times.  
 $\tau_Z$  is the 'Zeno time'.

**Note:** the quadratic behavior holds for any quantum transition, not only for decays. It is an absolutely general property.



## Time evolution and energy distribution (1)

The unstable state  $|S\rangle$  is not an eigenstate of the Hamiltonian  $H$ .  
Let  $d_S(E)$  be the energy distribution of the unstable state  $|S\rangle$ .

Normalization holds:  $\int_{-\infty}^{+\infty} d_S(E) dE = 1$

$$a(t) = \int_{-\infty}^{+\infty} d_S(E) e^{-iEt} dE$$

In stable limit :  $d_S(E) = \delta(E - M) \rightarrow a(t) = e^{-iMt} \rightarrow p(t) = 1$

## Time evolution and energy distribution (2)



Breit-Wigner distribution:

$$d_s(E) = \frac{\Gamma}{2\pi} \frac{1}{(E - M)^2 + \Gamma^2 / 4} \rightarrow a(t) = e^{-iM_0 t - \Gamma t / 2} \rightarrow p(t) = e^{-\Gamma t}.$$


The Breit-Wigner energy distribution cannot be exact.

Two physical conditions for a realistic  $d_s(E)$  are:

1) Minimal energy:  $d_s(E) = 0$  for  $E < E_{\min}$

2) Mean energy finite:  $\langle E \rangle = \int_{-\infty}^{+\infty} d_s(E) E dE = \int_{E_{\min}}^{+\infty} d_s(E) E dE < \infty$

# A very simple numerical example




$$M_0 = 2; E_{\min} = 0.75; \Gamma = 0.4; \Lambda = 3$$

$$d_s(E) = N_0 \frac{\Gamma}{2\pi} \frac{e^{-(E^2 - E_0^2)/\Lambda^2} \theta(E - E_{\min})}{(E - M_0)^2 + \Gamma^2/4}$$

$$d_{BW}(E) = \frac{\Gamma_{BW}}{2\pi} \frac{1}{(E - M_0)^2 + \Gamma_{BW}^2/4}$$

$$\Gamma_{BW}, \text{ such that } d_{BW}(M_0) = d_s(M_0)$$



$$a(t) = \int_{-\infty}^{+\infty} d_s(E) e^{-iEt} dE; \quad p(t) = |a(t)|^2$$

$$p_{BW}(t) = e^{-\Gamma_{BW} t}$$

# The quantum Zeno effect

We perform  $N$  inst. measurements:

the first one at time  $t = t_0$ , the second at time  $t = 2t_0$ , ..., the  $N$ -th at time  $T = Nt_0$ .


$$P_{\text{after } N \text{ measurements}} = p(t_0)^N \approx \left(1 - \frac{t_0^2}{\tau_Z^2}\right)^N = \left(1 - \frac{T^2}{N^2 \tau_Z^2}\right)^N$$

under the assumption that  $t_0$  is small enough.

$$\text{If } N \gg 1 \text{ (at fixed } T\text{): } P_{\text{after } N \text{ measurements}} \approx e^{-\frac{T^2}{N\tau_Z^2}} \approx 1.$$

For large but finite  $N$  :

→ slowing down of the decay.



## **Part 2: Experimental evidence of non-exponential decay**

# Experimental confirmation of non-exponential decays (1)

NATURE | VOL 387 | 5 JUNE 1997


## Experimental evidence for non-exponential decay in quantum tunnelling

**Steven R. Wilkinson, Cyrus F. Bharucha, Martin C. Fischer, Kirk W. Madison, Patrick R. Morrow, Qian Niu, Bala Sundaram\* & Mark G. Raizen**


*Department of Physics, The University of Texas at Austin, Austin, Texas 78712-1081, USA*

An exponential decay law is the universal hallmark of unstable systems and is observed in all fields of science. This law is not, however, fully consistent with quantum mechanics and deviations from exponential decay have been predicted for short as well as long times<sup>1-8</sup>. Such deviations have not hitherto been observed experimentally. Here we present experimental evidence for short-time deviation from exponential decay in a quantum tunnelling experiment. Our system consists of ultra-cold sodium atoms that are trapped in an accelerating periodic optical potential created by a standing wave of light. Atoms can escape the wells by quantum tunnelling, and the number that remain can be measured as a function of interaction time for a fixed value of the well depth and acceleration. We observe that for short times the survival probability is initially constant before developing the characteristics of exponential decay. The conceptual simplicity of the experiment enables a detailed comparison with theoretical predictions.

## Cold Na atoms in a optical potential



$$V(x,t) = V_0 \cos(2k_L x - k_L a t^2)$$




$$x' = x - \frac{1}{2} a t^2$$

$$U(x') = V_0 \cos(2k_L x') + M a x'$$

# Experimental confirmation of non-exponential decays (2)

Measured survival probability  $p(t)$



# Experimental confirmation of non-exponential decays and Zeno /Anti-Zeno effects



VOLUME 87, NUMBER 4

PHYSICAL REVIEW LETTERS

23 JULY 2001

## Observation of the Quantum Zeno and Anti-Zeno Effects in an Unstable System

M. C. Fischer, B. Gutiérrez-Medina, and M. G. Raizen

Department of Physics, The University of Texas at Austin, Austin, Texas 78712-1081

(Received 30 March 2001; published 10 July 2001)

We report the first observation of the quantum Zeno and anti-Zeno effects in an unstable system. Cold sodium atoms are trapped in a far-detuned standing wave of light that is accelerated for a controlled duration. For a large acceleration the atoms can escape the trapping potential via tunneling. Initially the number of trapped atoms shows strong nonexponential decay features, evolving into the characteristic exponential decay behavior. We repeatedly measure the number of atoms remaining trapped during the initial period of nonexponential decay. Depending on the frequency of measurements we observe a decay that is suppressed or enhanced as compared to the unperturbed system.




FIG. 3. Probability of survival in the accelerated potential as a function of duration of the tunneling acceleration. The hollow squares show the noninterrupted sequence, and the solid circles show the sequence with interruptions of  $50 \mu\text{s}$  duration every  $1 \mu\text{s}$ . The error bars denote the error of the mean. The data have been normalized to unity at  $t_{\text{tunnel}} = 0$  in order to compare with the simulations. The solid lines are quantum mechanical simulations of the experimental sequence with no adjustable parameters. For these data the parameters were  $a_{\text{tunnel}} = 15\,000 \text{ m/s}^2$ ,  $a_{\text{interr}} = 2000 \text{ m/s}^2$ ,  $t_{\text{interr}} = 50 \mu\text{s}$ , and  $V_0/h = 91 \text{ kHz}$ , where  $h$  is Planck's constant.

Zeno effekt

Same exp. setup,  
but with measurements in between





FIG. 4. Survival probability as a function of duration of the tunneling acceleration. The hollow squares show the noninterrupted sequence, and the solid circles show the sequence with interruptions of  $40 \mu\text{s}$  duration every  $5 \mu\text{s}$ . The error bars denote the error of the mean. The experimental data points have been connected by solid lines for clarity. For these data the parameters were:  $a_{\text{tunnel}} = 15\,000 \text{ m/s}^2$ ,  $a_{\text{interr}} = 2800 \text{ m/s}^2$ ,  $t_{\text{interr}} = 40 \mu\text{s}$ , and  $V_0/h = 116 \text{ kHz}$ .

Anti-Zeno effect

# GSI oscillations


## Measurement of weak decays of ions.



Decay of H-like Pm into:  
neutrino + Nd

Measurement was:

$$\frac{dN_{\text{decays}}}{dt} \propto -\frac{dp(t)}{dt}$$



Oscillations later confirmed.

arXiv:1309.7294 [nucl-ex]. Explanation still missing!

# Late-time deviations

PRL 96, 163601 (2006)

PHYSICAL REVIEW LETTERS

week ending  
28 APRIL 2006



## Violation of the Exponential-Decay Law at Long Times

C. Rothe, S. I. Hintschich, and A. P. Monkman

*Department of Physics, University of Durham, Durham, DH1 3LE, United Kingdom*

(Received 4 July 2005; published 26 April 2006)

First-principles quantum mechanical calculations show that the exponential-decay law for any metastable state is only an approximation and predict an asymptotically algebraic contribution to the decay for sufficiently long times. In this Letter, we measure the luminescence decays of many dissolved organic materials after pulsed laser excitation over more than 20 lifetimes and obtain the first experimental proof of the turnover into the nonexponential decay regime. As theoretically expected, the strength of the nonexponential contributions scales with the energetic width of the excited state density distribution whereas the slope indicates the broadening mechanism.




FIG. 2 (color). Corresponding double logarithmic fluorescence decays of the emissions shown in Fig. 1. Exponential and power law regions are indicated by solid lines and the emission intensity at time zero has been normalized.

Confirmation of: L. A. Khalfin. 1957. 1957 (Engl. trans. Zh.Eksp.Teor.Fiz.,33,1371)

Francesco Giacosa

## Considerations



- No other short- or long-time deviation from the exp. law was seen in unstable states.
- Verification of the two aforementioned works (Reizen + Rothe) would be needed.
- The measurement of deviations in simple natural systems (elementary particles, nuclei, atoms) would be a great achievement.

# Part 3: from the Lee Hamiltonian to Quantum Field Theory

# Lee Hamiltonian



$$H = H_0 + H_1$$

$$H_0 = M_0 |S\rangle\langle S| + \int_{-\infty}^{+\infty} dk \omega(k) |k\rangle\langle k|$$

$$H_1 = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g \cdot f(k)) (|S\rangle\langle k| + |k\rangle\langle S|)$$

$|S\rangle$  is the initial unstable state, coupled to an infinity of final states  $|k\rangle$ . (Poincare-time is infinite. Irreversible decay). General approach, similar Hamiltonians used in many areas of Physics.

(Ex: Jaynes-Cummings approach)

Example/1: spontaneous emission.  $|S\rangle$  represents an atom in the excited state,  $|k\rangle$  is the ground-state plus photon.

Example/2: pion decay.  $|S\rangle$  represents a neutral pion,  $|k\rangle$  represents two photons (flying back-to-back)

# Propagator and spectral function

$$H = H_0 + H_1 ; H_0 = M_0 |S\rangle\langle S| + \int_{-\infty}^{+\infty} dk \omega(k) |k\rangle\langle k| ; H_1 = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g \cdot f(k)) (|S\rangle\langle k| + |k\rangle\langle S|)$$

$$G_S(E) = \langle S | (E - H + i\epsilon)^{-1} | S \rangle = (E - M_0 + \Pi(E) + i\epsilon)^{-1}$$

$$\Pi(E) = - \int_{-\infty}^{+\infty} \frac{dk}{2\pi} \frac{g^2 f(k)^2}{E - \omega(k) + i\epsilon}$$

$$d_S(E) = \frac{1}{\pi} \text{Im} G_S(E) ;$$

$$a(t) = \langle S | e^{-iHt} | S \rangle = \int_{-\infty}^{+\infty} dE d_S(E) e^{-iEt}$$

It follows:

$$\int_{-\infty}^{+\infty} dE d_S(E) = 1$$

Fermi golden rule:  $\Gamma = \text{Im}[\Pi(M)] / 2$  .


# Exponential limit

$$H = H_0 + H_1 ; H_0 = M_0 |S\rangle\langle S| + \int_{-\infty}^{+\infty} dk \omega(k) |k\rangle\langle k| ; H_1 = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g \cdot f(k)) (|S\rangle\langle k| + |k\rangle\langle S|)$$

$$\omega(k) = k ; f(k) = 1 \Rightarrow \Pi(E) = ig^2 / 2 ; \Gamma = g^2$$

$$d_s(E) = \frac{\Gamma}{2\pi} \frac{1}{(E - M_0)^2 + \Gamma^2 / 4}$$

$$\Rightarrow a(t) = e^{-i(M_0 - i\Gamma/2)t} \Rightarrow p(t) = e^{-\Gamma t}$$




The exponential limit is obtained when the unstable state couples to all the states of the continuum with the same strength

# Non-exponential case (1)

$$H_1 = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g \cdot f(k)) (|S\rangle\langle k| + |k\rangle\langle S|)$$


$$f(k) = \begin{cases} 0 & \text{for } k < E_{\min} \\ 1 & \text{for } E_{\min} \leq k \leq E_{\max} \\ 0 & \text{for } k > E_{\max} \end{cases}$$




$$M_0 = 2; E_{\min} = 0; E_{\max} = 5; g^2 = 0.36 \text{ (all in a.u. of energy)}$$

This is what I have said at the beginning of the talk, but now “well done”

# Non-exponential case (2)



Dashed:  $p_{\text{BW}}(t) = e^{-\Gamma t}$  with  $\Gamma = \text{Im}[\Pi(M)] / 2$



$$h(t) = -\frac{dp(t)}{dt}$$

Dashed:  $h_{\text{BW}}(t) = \Gamma e^{-\Gamma t}$  with  $\Gamma = \text{Im}[\Pi(M)] / 2$

$$\int_0^t h(u)du = 1 - p(t)$$

Namley:  $h(t)dt = p(t) - p(t + dt)$  is the probability that the particles decays between  $t$  and  $t + dt$

# Two-channel case (a)

Found Phys (2012) 42:1262–1299  
DOI 10.1007/s10701-012-9667-3




## Non-exponential Decay in Quantum Field Theory and in Quantum Mechanics: The Case of Two (or More) Decay Channels

Francesco Giacosa

$$H_1 = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g_1 \cdot f_1(k)) (|S\rangle \langle k, 1| + |k, 1\rangle \langle S|) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk (g_2 \cdot f_2(k)) (|S\rangle \langle k, 2| + |k, 2\rangle \langle S|)$$

$$f_i(k) = \begin{cases} 0 & \text{for } k < E_{i,\min} \\ 1 & \text{for } E_{i,\min} \leq k \leq E_{i,\max} \\ 0 & \text{for } k > E_{i,\max} \end{cases}$$



$$M_0 = 2; E_{1,\min} = 0; E_{2,\min} = 0; E_{1,\max} = E_{2,\max} = 5;$$


$$g_1^2 = 0.36; g_2^2 = 0.16 \quad (\text{all in a.u. of energy})$$

Francesco Giacosa

## Two-channel case (b)

$h_1(t)dt =$  probability that the state  $|S\rangle$  decays in the first channel between  $(t,t+dt)$

$h_2(t)dt =$  probability that the state  $|S\rangle$  decays in the second channel between  $(t,t+dt)$



**Measurable effect???**

Details in:

F. G., Non-exponential decay in quantum field theory and in quantum mechanics: the case of two (or more) decay channels, Found. Phys. 42 (2012) 1262 [arXiv:1110.5923].

What about ab initio QFT?

# What about QFT? /1

## Quantum field theory: textbook treatment



$$d\Gamma = \frac{(2\pi)^4}{2M} |\mathcal{M}|^2 \delta(p - k_1 - k_2) \frac{d^3 k_1}{(2\pi)^3 2E_1} \frac{d^3 k_2}{(2\pi)^3 2E_2}$$

see e.g. Peskin-Schroeder or PDG

Care is needed:

- An unstable state is not an asymptotic state
- The formula is valid only for  $\Gamma \ll M$
- Within this treatment the decay is purely exponential
- One needs to go beyond to study non-exp. decays

# What about QFT/2

$$L_{\text{int}} = gS\phi^2$$

$[g] = [\text{Energy}]$ ; QFT super-renorm.

Propagator:



$$\Delta_S(p^2) = \frac{1}{p^2 - M_0^2 + \Pi(p^2) + i\epsilon}$$

Spectral function (or energy distribution):

$$d_S(m) = \frac{2m}{\pi} \text{Im}[\Delta_S(p^2 = m^2)]$$

Normalization follows automatically:

$$\int_0^\infty dm d_S(m) = 1$$



F.G. and G. Pagliara, *On the spectral functions of scalar mesons*,  
Phys. Rev. C 76 (2007) 065204 [arXiv:0707.3594].

# What about QFT/3

## Survival amplitude and example

Survival probability amplitude:

$$a(t) = \int_0^{\infty} dm d_s(m) e^{-imt}$$


Just as in QM: non-trivial result!

No dep. on cutoff for a  
superrenormalizable field theory

In order to show it:

- (i) Analogy to Lee models
- (ii) Work in the Schroedinger picture in QFT

Example:  $p(t)$  for the  $\rho$  meson



F. Giacosa and G. Pagliara, *Deviation from the exponential decay law in relativistic quantum field theory: the example of strongly decaying particles*,  
Mod. Phys. Lett. A **26** (2011) 2247 [arXiv:1005.4817 [hep-ph]].

## What about QFT/4

- a) Relativistic generalization of the Lee model.
- b) Matching. At one loop, QFT and the Lee models are exactly the same.

$$f(\mathbf{k}) = \sqrt{2} \frac{\tilde{\phi}(\mathbf{k})}{(\mathbf{k}^2 + m^2)^{1/4}}$$

where:  $f(\mathbf{k})$  is the coupling function for the Lee model  
 $\tilde{\phi}(\mathbf{k})$  is the QFT regularization function (can be set to 1 in a renormalizable theory)

Details in:

F. G., Non-exponential decay in quantum field theory and in quantum mechanics: the case of two (or more) decay channels, Found. Phys. 42 (2012) 1262 [arXiv:1110.5923].

# What about QFT/5

## Is there a “maximal energy scale“? The case of a renormalizable theory.

PHYSICAL REVIEW D **88**, 025010 (2013)

### Spectral function of a scalar boson coupled to fermions

Francesco Giacosa<sup>1</sup> and Giuseppe Pagliara<sup>2</sup>

<sup>1</sup>*Institut für Theoretische Physik Johann Wolfgang Goethe - Universität, Max von Laue-Strasse 1, D-60438 Frankfurt, Germany*

<sup>2</sup>*Dipartimento di Fisica e Scienze della Terra dell'Università di Ferrara and INFN Sezione di Ferrara,  
Via Saragat 1, I-44100 Ferrara, Italy*


$$\int_0^\Lambda d_H(m) dm = 1$$

no matter how large is  $\Lambda$ ...

but if one tries to do  $\Lambda \rightarrow \infty$  one encounters problems:  
normalization, etc.

$$d_H(m) \propto 1 / (m \cdot \ln^2 m) \quad \text{for large } m$$

Finite outcome: even for a renorm. QFT  
the existence of a maximal energy scale  
(i.e., a minimal length) is needed.



Renormalization: sweep dirt under  
the carpet?

## Part 4: Decay of a moving particle

## Unstable particle with momentum $p$



Up to now: in rest frame of the decaying particle. But what if it moves?  
Let us consider a momentum translation.

$$|S, p\rangle = U_p |S, 0\rangle$$

We **expect** in the exponential limit:

$$P_{nd}(t) = e^{-\frac{\Gamma}{\gamma}t}, \quad \tau = \gamma\Gamma^{-1} \text{ 'dilated lifetime'}$$

Reduction of the  
decay width

$$\frac{\Gamma}{\gamma} \equiv \frac{\Gamma M}{\sqrt{p^2 + M^2}} = \tilde{\Gamma}_p$$

## Finite momentum vs finite velocity



- Subtle but important point: in the long-life limit, a particle with definite momentum has also definite velocity.

$$p = \frac{M}{\sqrt{1 - v^2}}v$$

- In general, however, there is a difference! For an unstable state a boost is not equivalent to a momentum translation.
- Here, we consider a definite momentum

$$|S, p\rangle = \int_0^\infty dm a_S(m) |m, p\rangle$$

# Unstable particle with momentum $p$ : previous works



L. A. Khalfin, Theory of unstable particles and relativity, PDMI Preprint/1997


M. I. Shirokov, Int. J. Theor. Phys. 43 (2004) 1541.

E. V. Stefanovich, Int. Jour. Theor. Phys, 35 12 (1996)

K. Urbanowski, Phys. Lett. B 737 (2014) 346.

S. A. Alavi and C. Giunti, Europhys. Lett. 109 (2015) 6, 6001

My recent paper:



Francesco Giacosa

# Unstable particle with momentum $p$ : unexpected result for the nondecay probability



$$|S, p\rangle = U_p |S, 0\rangle$$

$$|S, p\rangle = \int_0^\infty dm a_S(m) |m, p\rangle$$

The non-decay probability:

$$P_{nd}(t) = e^{-\Gamma_p t}$$

$$\Gamma_p = \sqrt{2} \sqrt{\left[ \left( M^2 - \frac{\Gamma^2}{4} + p^2 \right)^2 + M^2 \Gamma^2 \right]^{1/2} - \left( M^2 - \frac{\Gamma^2}{4} + p^2 \right)}$$

F. G. arXiv:1512.00232 [hep-ph]

$$\Gamma_p \neq \tilde{\Gamma}_p = \Gamma M / \sqrt{p^2 + M^2}$$


But this is not a breaking of relativity!  
It is a different setup.

# Unstable particle with momentum p: deviation

$$\Delta = \frac{\Gamma_p - \tilde{\Gamma}_p}{M}$$

$$\Gamma_p = \sqrt{2} \sqrt{\left[ \left( M^2 - \frac{\Gamma^2}{4} + p^2 \right)^2 + M^2 \Gamma^2 \right]^{1/2} - \left( M^2 - \frac{\Gamma^2}{4} + p^2 \right)}$$

$$\tilde{\Gamma}_p = \frac{\Gamma}{\gamma} \equiv \frac{\Gamma M}{\sqrt{p^2 + M^2}}$$



arXiv:1512.00232 [hep-ph]

$$\frac{p_{\max}}{M} = \sqrt{\frac{2}{3}} \simeq 0.816$$

$$\Delta_{\max} = \frac{\Gamma_{p_{\max}} - \tilde{\Gamma}_{p_{\max}}}{M} \simeq \frac{3}{100} \sqrt{\frac{3}{5}} \left( \frac{\Gamma}{M} \right)^3$$

# QFT text-book



Back to QFT. The S-matrix approach

$$d\Gamma = \frac{(2\pi)^4}{2M} |\mathcal{M}|^2 \delta(p - k_1 - k_2) \frac{d^3 k_1}{(2\pi)^3 2E_1} \frac{d^3 k_2}{(2\pi)^3 2E_2}$$

is again justified for very small decay width!  
Here, the time-dilatation formula holds exactly.

The full QFT proof of the deviation is strictly speaking still missing.

(Technically, the formalism used above is based on so-called Lee Hamiltonians, which are QFT-like, but care is needed).

# Unstable particle with momentum p: some examples of deviations



Muon

$M = 105.65 \text{ MeV}$

$\Gamma = 2.99 \cdot 10^{-16} \text{ MeV}$

$$\Gamma_{p_{\max}} - \tilde{\Gamma}_{p_{\max}} \simeq 5.598 \cdot 10^{-53} \text{ MeV}$$

Neutral pion

$M = 134.98 \text{ MeV}$

$\Gamma = 7.72 \cdot 10^{-6} \text{ MeV}$

$$\Gamma_{p_{\max}} - \tilde{\Gamma}_{p_{\max}} \simeq 5.81 \cdot 10^{-22} \text{ MeV}$$

Rho meson

$M = 775.26 \text{ MeV}$

$\Gamma = 147.8 \text{ MeV}$

$$\Gamma_{p_{\max}} - \tilde{\Gamma}_{p_{\max}} \simeq 0.125 \text{ MeV}$$

Very small deviations!

## Boost: state with definite velocity revisited



$$U_v |S, 0\rangle \equiv |S, v\rangle$$

$$|S, v\rangle = \int_0^\infty dm a_S(m) \sqrt{m} \gamma^{3/2} |m, m\gamma v\rangle$$

The survival (or better, non-decay) probability vanishes at all time!

$$P_{nd}(t) = 0$$

A boosted muon consists of an electron and two neutrinos!

In reality, wave packets smear the effect.

Details in arXiv:1512.00232 [hep-ph]

## Summary...



- The decay is never exponential! This is a fact. This is so both in QM and QFT. Experiments exist, but new ones would be welcome.
- Experimentally seen, but new experiments would be needed
- New interesting effect still to be measured (two-channel case)
- Decay of a moving particle: interesting link between relativity and QM and QFT.
- For a particle with definite momentum  $p$  (for the measuring observer) there is a different formula. Numerically, the Einstein expression is very good but is not exact.
- A boost is a very subtle operation in QM and QFT.

...and outlook



Need of a ab initio Quantum Field Theoretical calculation... this is ongoing now. By using the standard technique (interaction picture, ...).

Thank You

- **When Physicists Attack: Homeless Man Attacks Fellow Transient in Disagreement Over Quantum Physics**
- [1, June 25, 2009 jonathanturley Bizarre, Criminal law, Society](#)
- This week a homeless man in California hit a fellow transient in the face with a skateboard over a disagreement about quantum physics. In San Francisco, Jason Everett Keller, 40, allegedly attacked, Stephan Fava, over a disputed physics question.
- At the time of the attack, Fava was discussing quantum physics with a third homeless man.
- I have been warning for years about the danger of “fighting words” in quantum physics discussions. I confess that I have come close to blows when I hear someone disparage Planck’s Action Constant in a bar.

# The Zeno's paradox in quantum theory

B. Misra and E. C. G. Sudarshan\*

*Center for Particle Theory, University of Texas at Austin, Austin, Texas 78712*

*(Received 24 February 1976)*

Analogy to the arrow

Today we speak of Quantum Zeno effect (and not paradox)

Can the cat be saved? Can the cat save its own life?

There is a difference between an infinitely frequent ideal observations and a continuous observation (that was not yet clear in 1976/1977)

# Experimental confirmation of the quantum Zeno effect - Itano et al (1)

PHYSICAL REVIEW A

VOLUME 41, NUMBER 5

1 MARCH 1990


## Quantum Zeno effect

Wayne M. Itano, D. J. Heinzen, J. J. Bollinger, and D. J. Wineland

*Time and Frequency Division, National Institute of Standards and Technology, Boulder, Colorado 80303*

(Received 12 October 1989)

The quantum Zeno effect is the inhibition of transitions between quantum states by frequent measurements of the state. The inhibition arises because the measurement causes a collapse (reduction) of the wave function. If the time between measurements is short enough, the wave function usually collapses back to the initial state. We have observed this effect in an rf transition between two  ${}^9\text{Be}^+$  ground-state hyperfine levels. The ions were confined in a Penning trap and laser cooled. Short pulses of light, applied at the same time as the rf field, made the measurements. If an ion was in one state, it scattered a few photons; if it was in the other, it scattered no photons. In the latter case the wave-function collapse was due to a null measurement. Good agreement was found with calculations.



(Undisturbed) survival probability

At  $t = 0$ , the electron is in  $|1\rangle$ .

$$p(t) = \cos^2\left(\frac{\Omega t}{2}\right) = 1 - \frac{\Omega^2 t^2}{4} + \dots$$

$$p(T) = 0 \text{ für } T = \pi/\Omega$$

# Experimental confirmation of the quantum Zeno effect - Itano et al (2)





FIG. 2. Diagram of the energy levels of  ${}^9\text{Be}^+$  in a magnetic field  $B$ . The states labeled 1, 2, and 3 correspond to those in Fig. 1.

5000 ions in a Penning trap

Short laser pulses 1-3 work as measurements.




FIG. 3. Graph of the experimental and calculated  $1 \rightarrow 2$  transition probabilities as a function of the number of measurement pulses  $n$ . The decrease of the transition probabilities with increasing  $n$  demonstrates the quantum Zeno effect.

$$p(t) = \cos^2(\Omega t / 2) = 1 - \frac{\Omega^2 t^2}{4} + \dots ; \quad p(T) = 0 \text{ für } T = \pi/\Omega$$

(Transition probability (without measuring) at time  $T$ ):  $1 - p(T) = 1$ .

With  $n$  measurements in between the transition probability decreases!

The electron stays in state 1.

## Other experiments about Zeno/Streed et al

PRL **97**, 260402 (2006)

PHYSICAL REVIEW LETTERS

week ending  
31 DECEMBER 2006

### **Continuous and Pulsed Quantum Zeno Effect**

Erik W. Streed,<sup>1,2</sup> Jongchul Mun,<sup>1</sup> Micah Boyd,<sup>1</sup> Gretchen K. Campbell,<sup>1</sup> Patrick Medley,<sup>1</sup>  
Wolfgang Ketterle,<sup>1</sup> and David E. Pritchard<sup>1</sup>

<sup>1</sup>*Department of Physics, MIT-Harvard Center for Ultracold Atoms, and Research Laboratory of Electronics, MIT,  
Cambridge, Massachusetts 02139, USA*

<sup>2</sup>*Centre for Quantum Dynamics, Griffith University, Nathan, QLD 4111, Australia*  
(Received 14 June 2006; published 27 December 2006)

Use of BEC (with Rb). QZE confirmed.

The intensity of a continuous observation of a quantum state is equivalent to a certain  $t_0$  (Shulman, PRA 57, 1509 (1997) ).

# Other experiments about Zeno/Haroche



PRL **101**, 180402 (2008)

PHYSICAL REVIEW LETTERS

week ending  
31 OCTOBER 2008



## Freezing Coherent Field Growth in a Cavity by the Quantum Zeno Effect

J. Bernu,<sup>1</sup> S. Deléglise,<sup>1</sup> C. Sayrin,<sup>1</sup> S. Kuhr,<sup>1,\*</sup> I. Dotsenko,<sup>1,2</sup> M. Brune,<sup>1,+</sup> J. M. Raimond,<sup>1</sup> and S. Haroche<sup>1,2</sup>

<sup>1</sup>*Laboratoire Kastler Brossel, Ecole Normale Supérieure, CNRS, Université P. et M. Curie,  
24 rue Lhomond, F-75231 Paris Cedex 05, France*

<sup>2</sup>*Collège de France, 11 Place Marcelin Berthelot, F-75231 Paris Cedex 05, France*

(Received 21 July 2008; published 28 October 2008)

Cavity QED: the nr of photons is frozen.

Another verification of QZE.

Direction QFT.

# Quantum Zeno dynamics, Quantum computations, ...



Sudarshan: Seven Science Quests

IOP Publishing

Journal of Physics: Conference Series 196 (2009) 012017

doi:10.1088/1742-6596/196/1/012017

## Quantum Zeno dynamics and quantum Zeno subspaces

Paolo Facchi<sup>1</sup>, Giuseppe Marmo<sup>2</sup>, Saverio Pascazio<sup>3</sup>

<sup>1</sup> Dipartimento di Matematica, Università di Bari

and Istituto Nazionale di Fisica Nucleare, Sezione di Bari, I-70125 Bari, Italy

<sup>2</sup> Dipartimento di Scienze Fisiche, Università di Napoli "Federico II"

and Istituto Nazionale di Fisica Nucleare, Sezione di Napoli, I-80126 Napoli, Italy

<sup>3</sup> Dipartimento di Fisica, Università di Bari

and Istituto Nazionale di Fisica Nucleare, Sezione di Bari, I-70126 Bari, Italy

E-mail: [saverio.pascazio@ba.infn.it](mailto:saverio.pascazio@ba.infn.it)

PRL 108, 080501 (2012)

PHYSICAL REVIEW LETTERS

week ending  
24 FEBRUARY 2012

## Zeno Effect for Quantum Computation and Control

Gerardo A. Paz-Silva,<sup>1,4</sup> A. T. Rezakhani,<sup>1,4,5</sup> Jason M. Dominy,<sup>1,4</sup> and D. A. Lidar<sup>1,2,3,4</sup>

<sup>1</sup>Department of Chemistry, University of Southern California, Los Angeles, California 90089, USA

<sup>2</sup>Department of Electrical Engineering, University of Southern California, Los Angeles, California 90089, USA

<sup>3</sup>Department of Physics, University of Southern California, Los Angeles, California 90089, USA

<sup>4</sup>Center for Quantum Information Science and Technology, University of Southern California, Los Angeles, California 90089, USA

<sup>5</sup>Department of Physics, Sharif University of Technology, Tehran, Iran

(Received 28 April 2011; revised manuscript received 9 February 2012; published 22 February 2012)

C. R. Physique 17 (2016) 685–692



Contents lists available at [ScienceDirect](http://ScienceDirect)

Comptes Rendus Physique

[www.sciencedirect.com](http://www.sciencedirect.com)



Quantum microwaves / Micro-ondes quantiques

Quantum Zeno dynamics in atoms and cavities

*Dynamique de Zénon quantique avec des atomes et des cavités*



Sébastien Gleyzes\*, Jean-Michel Raimond

Laboratoire Kastler Brossel, Collège de France, CNRS, ENS-PSL Research University, UPMC-Sorbonne Universités, 11, place Marcelin-Berthelot, 75005 Paris, France



Francesco Giacosa

# Other experiments about Zeno/Balzer



Same setup as Itano et al. (different ions are used, YB instead of Be),

But now the measurement takes place between 3 and 2.

Results in agreement with Itano, but here the QZE is associated by a series of null-measurements.

## Unstable particle with momentum $p$

We work in the exp. limit

$M$  = rest mass;  $\Gamma$  = decay width in the rest frame.

An unstable particle moves with definite momentum  $p$ .

Which is its decay width? The standard expression is:

$$\tilde{\Gamma}_p = \frac{\Gamma}{\gamma} \equiv \frac{\Gamma M}{\sqrt{p^2 + M^2}}$$

Important but subtle point:

in QM and QFT a state with definite momentum has not definite velocity.

# Non-decay probability

Straightforward calculation

$$\begin{aligned} a(t, p) &= \frac{1}{\delta(p=0)} \langle S, p | e^{-iHt} | S, p \rangle = \int_{-\infty}^{\infty} dm d_S(m) e^{-i\sqrt{m^2+p^2}t} \\ &\simeq \int_{-\infty}^{\infty} dm d_S^{BW}(m) e^{-i\sqrt{m^2+p^2}t} = e^{-i\sqrt{(M-i\Gamma/2)^2+p^2}t} . \end{aligned}$$

One obtains:

$$P_{nd}(t) = |a(t, p)|^2 = e^{-\Gamma_p t}$$

$$\Gamma_p = 2 \operatorname{Im} \left[ \sqrt{(M - i\Gamma/2)^2 + p^2} \right] .$$

This expression does **not** coincide with the usual Einstein expression!

## Decay width of a general state

$$|\Psi\rangle = \int_{-\infty}^{+\infty} dp B(p) |S, p\rangle$$

the quantity  $\langle \Psi | e^{-iHt} | \Psi \rangle$  is *not* what we are looking for.

$$P_{nd}(t) = \int_{-\infty}^{+\infty} dp |\langle S, p | e^{-iHt} | \Psi \rangle|^2$$

$$P_{nd}(t) = \int_{-\infty}^{+\infty} dp |B(p)|^2 e^{-\Gamma_p t}$$

Inclusion of spatial wave function is simple. Generalization straightforward.  
Details in arXiv:1512.00232.

## Boost: state with definite velocity



Point: a velocity translation (i.e. a boost) is not a momentum translation!!!!

$$U_v |S, 0\rangle \equiv |S, v\rangle$$

$$|\langle S, v | e^{-iHt} | S, v \rangle|^2 = e^{-i\gamma\Gamma t}$$

The survival probability shows here an absurd Lorentz contraction!