



Contribution ID: 1548

Type: Talk

From Acceleration Radiation to Black Hole Entropy: A quantum optical perspective

Acceleration radiation is caused by virtual processes in which an atom jumps from the ground state to an excited state, together with the emission of a photon. If an atom is accelerated away from the original point of virtual emission then there is a small probability that the virtual photon will “get away”. When ground-state atoms are accelerated through a high Q microwave cavity, radiation is produced with an intensity which can exceed the intensity of acceleration radiation in free space by many orders of magnitude. The reason is a strong nonadiabatic effect at cavity boundaries [1].

By formulating acceleration radiation as a flat space quantum optics problem one can obtain the entropy of a black hole (BH). Using this approach we find the famous Bekenstein-Hawking BH entropy formula via a simple “laser-like” density matrix analysis. In this way we calculate the entropy of the radiation emitted by a cloud of infalling atoms without introducing the BH temperature [2].

[1]. M.O. Scully, V.V. Kocharovsky, A. Belyanin, E. Fry and F. Capasso, Phys. Rev. Lett. 91, 243004 (2003).

[2]. M.O. Scully, D. Lee, W. Schleich and A.A. Svidzinsky, Black hole acceleration radiation: from a quantum optical perspective, to be published.

Topic:

Mini-workshop: Quantum Foundations and Quantum Information

Summary

Primary author: Prof. SCULLY, Marlan (Texas A&M University, Baylor University, Princeton University)

Presenter: Prof. SCULLY, Marlan (Texas A&M University, Baylor University, Princeton University)

Session Classification: Workshop on Quantum Foundations and Quantum Information

Track Classification: Workshop on Quantum Foundations and Quantum Information