



Tirana workshop,
Physics dept, Tirana University
September 26 and 27th, 2016

The Standard Model and the LHC

D. Denegri,
CEN Saclay/IRFU/SPP and CERN/Ph

introduction on the LHC and CMS
testing the Standard Model at the LHC,
up to the discovery of the Higgs boson
beyond the Standard Model searches at the LHC
what next?




Looking for the Higgs:

The Large Hadron Collider - genesis of the project



The LHC project started at the initiative (and with the daring!!) of C. Rubbia

The Aachen Conference in October 1990 marked the start-up, since then work on the collider and magnets, various detector designs and understanding physics issues went on without let-up




Scientifico-diplomatic trips in 1990/91/92 to Japan, India, Russia, USA, Canada etc

LHC vs SSC: Rubbia's arguments: savings!



- existing LEP tunnel ~1 GCHF
- existing infrastructure at CERN (PS, SPS, etc) ~ 1 GCHF
- "two-in-one" scheme for dipoles saves ~ half the cost of magnet ~ 0.7 to 1 GCHF

thus overall LHC cost ~ 3 GCHF






The LHC - a single set of magnets in the tunnel thanks to the “two-in-one” magnet scheme



the ISR - a typical proton-proton collider;
ISABELLE and the SSC were of same design




The LHC tunnel - a single set of magnets thanks to the “two-in-one” magnet scheme - II



The magnets have to be aligned with $100\mu\text{m}$ precision over the 27 km circumference!

Vacuum in the beam pipe as on the Moon!

Magnetic field lines





the two-in-one magnet
40.000 tons of material at $-271\text{ }^{\circ}\text{C}$
The coldest place in the Univers!



The LHC: required energy and luminosity

- To solve e-w symmetry breaking need to study W_L - W_L scattering at a centre of mass energy of up to ~ 1 TeV



$E_W \sim 500$ GeV

$E_{\text{quark}} \sim 1$ TeV
 $E_{\text{proton}} \sim 6$ TeV

required LHC energy: pp collisions at $\sim 6 + 6$ TeV

- Event Rate = Luminosity x Cross-section x Branching Ratio = $L \times \sigma \times BR$

e.g. $H(1\text{TeV}) \rightarrow ZZ \rightarrow 2e+2\mu$ or $4e$ or 4μ

For 10 events/year = $10^{34} \times 10^7 \times 10^{-37} \times 10^{-3}$!! required $L \sim 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$


- Luminosity, gives the number of proton-proton collisions taking place per sec.

For $L = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ and $\sigma(pp) \sim 100$ mb rate = $10^9/\text{sec}$!!!



The Large Hadron Collider – overview

~ 65% of the 27 km long circumference covered with 1232 2-in-1 superconducting dipoles of 14.3m length operated at 1.9 °K giving a field of $B = 8.3\text{T}$, altogether 1200 tons of superconducting cable and 40.000 tons of material at 1.9 °K superfluid He temperature!



LHC(2012) pp	8.000 GeV	$\sim 7 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$
(2015) PbPb	5.000(NN) GeV	$\sim 10^{27} \text{ cm}^{-2} \text{ s}^{-1}$

in 2016 pp:	13.000 GeV	$\sim 1.2 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
in ~2025:	13.000 GeV	$\sim 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$


$$p = ReB$$





**A typical proton-proton collision at the LHC,
there are ~ one billion such collisions per second
(in reality they are clustered by ~25 every 25 nanoseconds – pile-up)**


Expected Higgs boson production rate is less than one in a billion pp collisions!





LHC operation in 2012 : pile-up, up to ~ 30
(50 nsec bunch spacing),

in 2016 we have ~ 45 pile-ups (25 nsec spacing)




In HL-LHC phase ($\sim 5 \times 10^{34} \text{cm}^{-2} \text{s}^{-1}$) we expect to go to ~150 pile-ups per crossing!



The Large Hadron Collider and experiments

Collision of proton beams...

...observed in giant detectors




- ▶ p (proton)
- ▶ \bar{p} (antiproton)
- ▶ ion
- ▶ ▶ proton/antiproton conversion
- ▶ neutron
- ▶ neutrino

- AD Antiproton Decelerator
- PS Proton Synchrotron
- SPS Super Proton Synchrotron

- LHC Large Hadron Collider
- n-ToF Neutron Time of Flight
- CNGS CERN Neutrinos to Gran Sasso



ATLAS and CMS proton-proton luminosities, from 2011 till mid-July 2016



Notice rate in 2016!



Current (September 1st) LHC schedule for 2016

	July			Aug				Sep						
Wk	27	28	29	30	31	32	33	34	35	36	37	38	39	
Mo	4	11	18	25	1	8	15	22	29	5	12	beta* = 2.5 km data taking	19	26
Tu								MD 2						
We											TS2			
Th				MD 1						Jeune G				
Fr								beta* 2.5 km dev.						
Sa										MD 3				
Su				beta* 2.5 km dev.										

Proton-proton running


heavy ions running (Pb-Pb, p-Pb)

	Oct			Nov				Dec					
Wk	40	41	42	43	44	45	46	47	48	49	50	51	52
Mo	3	10	17	24	31	7	14	21	28	5	12	19	26
Tu	MD 4						ions setup				Extended year end technical stop		
We						TS3							
Th													
Fr					MD 5				Ion run (p-Pb)				
Sa													
Su										Pb MD		Xmas	New Year

End of run
[06:00]



The CMS (Compact Muon Solenoid) detector



CMS designed in ~1991/92, design optimized up to ~1995; Prototypes for sub-detectors tested up to ~2000;

Construction from ~2000 till 2008
Operational in Sept 2008




CMS in transverse view - basic design

C.M.S.
A Compact Solenoidal Detector for L.H.C.

B ~ 2 Tesla

B = 4 Tesla

Total bending
power on a
muon ~ 16Tm



Transverse View

CMS-TS-00079

$$p = reB$$

momentum resolution
from curvature:

$$\Delta p/p \sim \epsilon p / (\sqrt{n} B l^2)$$

ϵ = resolution on point
measurement

n = number of points
measured per track


l = useful track length

B = magnetic field

The goal is to identify and measure all particles produced at the interaction point



LHC experiments are really international endeavors! CMS for example:



Support tube
Support tube

Flux return yoke

Germany, Russia, Chekia, Japan

Hadron calorimeter

USA, Russia, Ukraina, Turkey, Iran,
India, Hungary

Solenoid magnet

France, Italy, Switzerland, Finland,
Croatia, UK, Japan, CERN

Tracking system

Germany, Italy, France, Belgium, USA
Finland, Switzerland, CERN.....

Electromagnetic calorimeter

Russia, China, France, Italy, Japan,
UK, Switzerland, Greece, Taiwan

Muon system

Italy, Germany, Spain, USA, Russia,
Bulgaria, Belgium, Korea, Pakistan, CERN

Support system

China, Pakistan, USA





SM Physics

jets, W, Z, top, Higgs boson



Among first LHC results: Inclusive single jet double-differential cross section at 8 TeV




10-12 orders of variation in magnitude in cross section!!

Total uncertainty 50-10% dominated by JES

Good agreement between data and NLO pQCD with various PDFs




newest studies extend beyond jet- E_t of 2 TeV





3-jet to 2-jet cross section ratio and the evolution of the strong coupling constant $\alpha_s(Q)$

test of QCD asymptotic freedom in SM





$$\alpha_s(M_Z) = 0.1148 \pm 0.0014(\text{exp.}) \pm 0.0018(\text{PDF})_{-0.0000}^{+0.0050}(\text{scale})$$



Jets at 13 TeV, CMS, 2015 data, 1.6 fb⁻¹

2015 25 ns data: extend up to 3 TeV, improve precision in p_T tail, perform α_s fit and study its evolution vs Q






Dijet event in CMS, 13 TeV



CMS Experiment at the LHC, CERN

Data recorded: 2016-May-11 21:40:47.974592 GMT


Run / Event / LS: 273158 / 238962455 / 150







Dijet event in ATLAS at 13 TeV, 2016 data

Large increase in cross section for high mass states going from 8 TeV to 13 TeV






Early 13 TeV data, 800 pb⁻¹ – dijets effective mass






Searches in dijet final states, CMS 13 TeV, 13 fb⁻¹

Low mass Analysis





High mass Analysis





Dijets - search for resonances, 2016 data, $\sim 13\text{fb}^{-1}$




Examples of limits (@95% CL):
 $m(q^*) > 5.6\text{ TeV}$ (ATLAS Run-1: 4.1 TeV)
 $Z' > 1.5\text{ TeV}$

Strongest limit:
 STRING resonances excluded
 up to 7.4 TeV



Microscopic quantum Black Holes - 7 TeV results


- BH production in ADD model
(large flat extra spatial dimensions)
- Democratic and isotropic decay
 - High S_T events (total transverse energy)
 - High total multiplicity (e.g. ≥ 4)



10 Jet event, $S_T = 1.1$ TeV

Limits on Black hole
> 4 – 5 TeV @ 95% C


Microscopic black hole evaporation:
Events with large transverse energy
sum, S_T





ATLAS, searches with jets at 13 TeV, 2016, $\sim 3\text{fb}^{-1}$

Search for resonances decaying to two jets. No significant excess seen.
Limits on quantum black holes, excited quarks, W' , Z' models



Search for thermal black holes
Signal region : 3 to 8 jets
Signal region at high HT
No significant excess, limits on
threshold mass $\sim 9\text{ TeV}$




First $W \rightarrow e\nu$ and $Z \rightarrow e^+e^-$ events in LHC, April 2010




CMS Experiment at LHC, CERN
Run 133874, Event 21466935
Lumi section: 301
Sat Apr 24 2010, 05:19:21 CEST

Electron $p_T = 35.6$ GeV/c
 $ME_T = 36.9$ GeV
 $M_T = 71.1$ GeV/c²




CMS Experiment at LHC, CERN
Run 133877, Event 28405693
Lumi section: 387
Sat Apr 24 2010, 14:00:54 CEST

Electrons $p_T = 34.0, 31.9$ GeV/c
Inv. mass = 91.2 GeV/c²




W and Z spectra in CMS, Nov. 2010



By end-2012 we had
~150.000.000 W and
~15.000.000 Z
decaying leptonically!!



Note progress since
first W, Z!





W, Z production cross sections, testing QCD from the CERN pp-bar collider to the Tevatron and the LHC



W,Z at 13 TeV, first look June 2015







W[±] charge asymmetry, CMS and LHCb, - constraining proton structure functions

W[±] → μ[±] ν charge asymmetry vs η_μ
combining CMS and LHCb data



$$\eta_\mu = -\ln \text{tg}(\theta_\mu/2)$$



$$A(\eta) = \frac{\frac{d\sigma}{d\eta}(W^+ \rightarrow l^+\nu) - \frac{d\sigma}{d\eta}(W^- \rightarrow l^-\bar{\nu})}{\frac{d\sigma}{d\eta}(W^+ \rightarrow l^+\nu) + \frac{d\sigma}{d\eta}(W^- \rightarrow l^-\bar{\nu})}$$






$W(\rightarrow \mu\nu) + \text{jets}$, 13 TeV, CMS

Measurement of the differential cross section $W(\mu\nu)+\text{jets}$
CMS, 2.5fb^{-1} , 13 TeV, compared with NNLO for one inclusive jet
and NLO for all inclusive jet spectra.



Standard Model Production Cross Section Measurements

Status: August 2016








Top quark physics - some aspects



Study of the top quark: one of the main goals of the LHC

LEPTONS		
Electron Neutrino Mass ~0	Muon Neutrino ~0	Tau Neutrino ~0
Electron .511	Muon 105.7	Tau 1.777
QUARKS		
Up Mass: 5	Charm 1.500	Top ~180.000
Down 8	Strange 160	Bottom 4.250



$$g_{Hff} = m_f/v$$



At nominal energy and luminosity LHC produces about 1 top-antitop pair second!

The LHC could be called a 'top factory'


Top quark physics after the Higgs discovery:

- Special role in EWSB mechanism?
- Does it play a role in non-SM physics?



Top at the LHC, production and decays for ex. $top \rightarrow lepton + jets$ final states


tt -bar production diagrams, some....



Top decay in the SM: $t \rightarrow bW$


tt -bar event decay topology in «single lepton mode»:

$tt \rightarrow bWbW \rightarrow blvbqq$






Top pair production cross-sections, up to 13 TeV



LHC and Tevatron results consistent and in agreement with NNLO+NNLL over a large range of center-of-mass energies



tt-bar differential cross section, ATLAS, 13 TeV - testing QCD




Differential tt cross section (l+jets):
ATLAS, 3.2 fb⁻¹, 13TeV, tops resolved and boosted.




Top properties, charge, status in 2013 and final results at 8 TeV

semi-muonic $t\bar{t}$ + photon



$t\bar{t}+\gamma$ $\Delta R(\gamma, \text{jet}) > 0.1$ and $p_T(\gamma) > 2\text{GeV}$

$\sigma(t\bar{t}+\gamma) = 2.4 \pm 0.2(\text{stat.}) \pm 0.6(\text{syst.}) \text{ pb}$
 theory: $1.8 \pm 0.5 \text{ pb}$




$t\bar{t}+\text{jets}$

- Ratio $t\bar{t}\gamma/t\bar{t}$ in fiducial region
- 21% precision




Top mass - from the semileptonic mode, the most appropriate one for mass.....

$$t\bar{t} \rightarrow bW bW \rightarrow bl\nu bqq$$




CMS average : $172.6 \pm 0.1 \pm 0.7$ GeV

Sum of permutation weights / 5 GeV







Top mass in top pairs with J/psi in b-decay chain, 3-lepton final states, at 8 TeV



Using the correlation between the 3-prong mass and the top quark mass
An old idea, first application of
CMSNOTE-
2006-058

measurement statistics limited at present, but ultimately should provide the most precise measurement






CMS, top mass measurements, status September 2016


- We have the best mass measurement in the world (by combining the best analyses per channel), TOP-14-022, Phys. Rev. D 93, 072004 (2016)
- We have several "alternative measurements", not meant to be competitive but complementary



Testing the coherence of the Standard Model - importance of precision measurements of W, top and Higgs masses



W mass correlated to the Higgs and top masses






Electroweak measurements - di-boson channels



ZZ → 4 leptons, 13 TeV, CMS, summer 2016





Measurement of $ZZ \rightarrow ll'l'$

CMS, 2.6fb^{-1} , 13 TeV

$\sigma(ZZ) = 14.6^{+1.9}_{-1.8}(\text{st})^{+0.5}_{-0.3}(\text{sy}) \pm 0.2(\text{th}) \pm 0.4(\text{lu}) \text{ pb}$

NNLO: $15.0^{+0.7}_{-0.6} \pm 0.2 \text{ pb}$

~14% precision, statistically limited






WZ \rightarrow 3 leptons, 13 TeV, CMS, summer 2016

Measurement of $WZ \rightarrow \ell\nu\ell'\ell'$

CMS, 2.3fb^{-1} , 13 TeV


$\sigma(WZ) = 39.9 \pm 3.2(\text{stat})^{+2.9}_{-3.1}(\text{syst}) \pm 0.4(\text{theo}) \pm 1.3(\text{lumi}) \text{ pb}$

NNLO: $50.0^{+1.1}_{-1.0} \text{ pb}$




June 2016

CMS Preliminary




Diboson Cross Section Measurements

Status: August 2016





SM cross section summary, ATLAS, energy dependence 7, 8, 13 TeV






the Higgs.....




Production and detection of the Higgs in CMS - if $m_H \sim 150$ GeV ($H \rightarrow ZZ/ZZ^* \rightarrow 4$ leptons) - expectations from 1992!



expectations from 1992!



Expectations for signal and background if the Higgs has a mass ~ 150 GeV
Similar situation in the mass range $\sim 130 - \sim 400$ GeV

Expected Higgs signal



$M_{\text{Higgs}} = 150$ GeV



Electroweak ZZ background!!



Electroweak ZZ background








$H \rightarrow ZZ \rightarrow \mu\mu\mu\mu$ candidate event in CMS, $\sqrt{s} = 8 \text{ TeV}$ data, 2012






Higgs decay to 4 leptons, ATLAS and CMS, July 4th 2012 - discovery




Notice these modest beginnings! As for W, Z!



H \rightarrow $\gamma\gamma$ candidate in CMS





CMS Experiment at the LHC, CERN
Data recorded: 2012-May-13 20:08:14.621490 GMT
Run/Event: 194108 / 564224000



$$m_{\gamma\gamma}^2 = 2 E_1 E_2 (1 - \cos\theta_{12})$$



H \rightarrow $\gamma\gamma$ in CMS, July 2012



Nobel Prizes and Laureates

Physics Prizes

< 2013 >

▼ About the Nobel Prize in Physics 2013

[Summary](#)[Prize Announcement](#)[Press Release](#)[Advanced Information](#)[Popular Information](#)[Greetings](#)[Award Ceremony Video](#)[Award Ceremony Speech](#)[► François Englert](#)[► Peter Higgs](#)[All Nobel Prizes in Physics](#)[All Nobel Prizes in 2013](#)

The Nobel Prize in Physics 2013

François Englert, Peter Higgs

Share this:     1.9K 

The Nobel Prize in Physics 2013



Photo: A. Mahmoud

François Englert

Prize share: 1/2



Photo: A. Mahmoud

Peter W. Higgs

Prize share: 1/2

The Nobel Prize in Physics 2013 was awarded jointly to François Englert and Peter W. Higgs *"for the theoretical discovery of a mechanism that contributes to our understanding of the origin of mass of subatomic particles, and which recently was confirmed through the discovery of the predicted fundamental particle, by the ATLAS and CMS experiments at CERN's Large Hadron Collider"*




What is the mass?

Measure the mass through high precision channels

$H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ \rightarrow 4l$

19.7 fb⁻¹ (8 TeV) + 5.1 fb⁻¹ (7 TeV)




$$m_H = 125.36 \pm 0.37 \text{ (stat)} \pm 0.18 \text{ (syst)} \text{ GeV} \quad m_H = 125.03 \pm_{0.27}^{0.26} \text{ (stat)} \pm_{0.15}^{0.15} \text{ (syst)} \text{ GeV}$$

for ATLAS and CMS






ATLAS + CMS combination on Higgs mass, final run1 analysis, April 2015



precision of ~2 pro mille!!



H → 4 leptons, Sept-15th 2016, CMS, 13 fb⁻¹





$$m_H = 124.50^{+0.48}_{-0.46} \text{ GeV} = 124.50^{+0.47}_{-0.45}(\text{stat.})^{+0.13}_{-0.11}(\text{sys.}) \text{ GeV}$$

$$\mu = \sigma / \sigma_{SM} = 0.99^{+0.33}_{-0.26}$$



H \rightarrow $\gamma\gamma$, 13 TeV, 13fb⁻¹, ATLAS, August 2016



13 TeV

ATLAS (13.3 fb⁻¹)

(N3LO+XH)

Fiducial σ (fb)

$43.2 \pm 14.9(\text{stat}) \pm 4.9(\text{syst})$


SM prediction (fb)

$62.8^{+3.4}_{-4.4}$




Higgs production mechanisms and decay modes

87.4 % @120 GeV




6.7 %





indirect coupling to gg,
top in the loop

5.4 %




0.5 %




$$g_{HVV} = 2m_V^2/v$$


$$g_{HWW} = g m_W$$



$$g_{Hff} = m_f/v$$







$$g_{HHH} = 3m_H^2/v$$



indirect coupling to $\gamma\gamma$



$t\bar{t}H(H \rightarrow b\bar{b})$ - important channel, measures directly Higgs coupling to top




First result from
 ATLAS,
 $13.2 \text{ fb}^{-1}, 13 \text{ TeV}$.



Higgs couplings - future up to HL-LHC and 3000fb⁻¹

1


Relation Couplage-Masse



$$g_{Hff} = m_f/v$$


$$g_{HHH} = 3m_H^2/v$$

$$g_{HVV} = 2m_V^2/v$$



v

Higgs self-coupling and Higgs potential



With 3000fb⁻¹ data at 13 TeV (by ~ 2025) couplings to 8 particles, including coupling to muons





- Looking further for physics
beyond the Standard Model
- diphotons
 - dilepton pairs, Z' , W' etc
 - SUSY
 - DM.....



Searches for heavy dilepton or diphoton resonances, extra dimensions

Search for excesses in invariant mass spectra:

- Predicted by several BSM models with extended gauge symmetries
- Z' and W' with SM-like couplings
- Kaluza-Klein excitations from RS model of extra dimensions
- E6 models
- Technicolor
-




$$M_{Pl}^2 \sim M_D^{2+n} R^n$$

- Searches for non-resonant excess in kinematic distributions and mass spectra
- Predicted by many Extra Dimension Models
- Universal Extra Dimensions (UED), RS etc





Drell-Yan (e^+e^-) pairs, ATLAS and CMS, 13 TeV search for heavy $Z' \rightarrow l^+l^-$ new gauge boson or KK excitation

Production and Decay of a W'





New limit in SSM:

$$Z'_{SSM} > 4 \text{ TeV}$$





Search for heavy W' in the $W' \rightarrow \text{lepton} + E_t^{\text{missing}}$ channel, 13 TeV data, new gauge boson or KK excitation






New limit in SSM:

SSM $W' > 4.74 \text{ TeV}$




Search for W' in the $W' \rightarrow tb$ channel, 13 TeV data, summer 2016





Models with $e + E_t^{\text{missing}}$, HL-LHC expectations





Several BSM models predict signal in lepton+MET




- W' signal with characteristic Jacobian peak (still visible in electron channel)
- Other models, contact interaction (CI) and DM, xsec not defined by theory but depends on free model parameter L .

W' scaled to theoretical xsec

Distributions scaled to 1 fb.



SUSY - could solve a number of problems at ew scale



Possibly the DM particle....


Stabilizes Higgs boson mass



Supersymmetry - a typical mass spectrum

$m_0 = 90, m_{1/2} = 400, \tan\beta = 10, \mu = 518$

four are here
A, H, H⁺H⁻
direct searches
give no sign yet



produced by strong interactions
direct searches give no sign yet, might be > 2TeV

Stop - produced by strong interactions
could be much lighter than all others sis (>~ 500 GeV)
no sign yet

produced also by ew interactions, no sign yet

Neutralino-1, LSP - DM candidate
 $\tilde{\chi}_1^0$


the 125 GeV one?
possibly.....

...l' objet de toutes nos pensées




Badly unbalanced SUSY-candidate-type event

CMS Experiment at LHC, CERN
Data recorded: Wed Jul 13 09:50:28 2016 CEST
Run/Event: 276776 / 715087594
Lumi section: 433







SUSY searches in CMS, gluino to top, 8 TeV



- Assumptions:
 - $m_{\tilde{t}} > m_{\tilde{g}}$
 - $\text{BR}(\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0) = 100\%$




Gluginos with mass < 1.3 TeV excluded...



SUSY status in 2013 - RPC

Summary of CMS SUSY Results* in SMS framework


SUSY 2013





*Observed limits, theory uncertainties not included
 Only a selection of available mass limits
 Probe *up to* the quoted mass limit



SUSY studies at 13 TeV, stop studies



- Many regions, defined by HT, MT2, Njet(≥ 1), Nb,pTjet
- Final sample, refined treatments, T2tt/T2bb/T2qq interpretation added







SUSY searches, 13 TeV, ATLAS, sbottom, gluinos versus neutralino-1 exclusion plots


Large gains in terms of cross sections for massive particle production in going from 8 to 13 TeV collision energy

For squarks/gluinos of ~ 1.5 TeV the gain is factor of 35!





Limits on gluinos and neutralinos, CMS, mid 2016






Stop perspectives....14 TeV LHC, 300 fb⁻¹

expected in ~2022

CMS Preliminary




If the 125 GeV Higgs is in fact the h from MSSM, then stop is likely in the ~0.5 -1 TeV range (other squarks being ~ > 2-3 TeV)

With ~14 TeV, stop could be discovered up to ~950 GeV with 300 fb⁻¹






Dark Matter searches, direct vs collider searches

Direct searches
elastic scattering
(t-channel)



Collider searches
Pair production
(s-channel)



radiation of a photon - or gluon - in the initial state makes the process visible





Search for mono-jets and mono-photons, 7TeV

Search the gamma+MET and jet+MET data

Look for evidence of Dark Matter production Or extra dimensions

Monojet: ADD limits set > 4 TeV for $n = 2$




Monophoton: > 1.59 (1.66) TeV for $n=3(6)$







DM searches - early limits

Monojet (EXO-11-059) and monophoton (EXO-11-096) analyses probing models with extra dimensions and dark matter Best limits on DM with spin-dependent couplings to date








Dark Matter searches in mono-photons and mono-jets, CMS, 13 TeV, summer 2016 studies



No excess seen!





DM at LHC, mid-2016 status




A coherent picture of DM:

- No significant excess observed so far
- DM mass exclusion up to ~ 550 GeV
- Vector Mediator mass exclusion up to 1.95 TeV

Exotic searches



CMS Preliminary



Many searches met or exceeded the sensitivity of Run 1




Possible very long-term future (~2035/40) - FCC project

An Ultra Large Collider to reach 100 TeV in pp mode, an order of magnitude larger than LHC, with an e^+e^- initial phase at ~350 GeV, potential for e-p and Pb-Pb - the FCC project. Projects at the technology frontier at level of design studies and generating requiring/motivating ambitious R&D efforts


An 80-100 km tunnel encompassing all of the Geneva area....

There are also the ILC, CLIC....projects






FCC-hh main cross sections





New weak gauge interactions, Fcc-hh reach with 10 ab^{-1}





Discovery reach

T.Rizzo, arXiv:1403.5465

Model	1 ab^{-1}	10 ab^{-1}	100 ab^{-1}
SSM	23.8	33.3	41.3
LRM	22.6	31.5	39.5
ψ	20.1	29.1	37.2
χ	22.7	30.6	38.2
η	20.3	29.8	38.0
I	22.4	29.2	36.2



SUSY production at 100 TeV, Fcc-hh mass reach for gluinos, stops, sbottoms





Conclusions

The LHC is an incredible technological and scientific endeavor - on a world-wide scale - and a great success

The experiments ATLAS, CMS, ALICE and LHCb are all operating very successfully. The physics results up to now are magnificent, discovery of the Higgs, beautiful and detailed studies in EWK, QCD and B-physics, rare decay modes, QGP studies etc

In 2016 LHC started operating at 13 TeV; many technical challenges overcome. Near future: clarification of the Higgs (is it THE Higgs boson or a A Higgs from an extentionn of the SM), but the main task is to look for physics beyond the Standard Model, looking for SUSY etc especially so in the HL-LHC phase.

The particle physics community is already working on long-term options/projects to take over in ~ 20-30 years from now