

Event-shape of dileptons plus missing energy at a linear collider as a SUSY/ADD discriminant

Probir Roy


Tata Institute of Fundamental Research, Mumbai, India

Based on work with Partha Konar, hep-ph/0509161, Phys. Lett. B634 (2006) 295

- Proposal
- Signal
- SM background and chosen cuts
- Cross-sections and event-shape variables
- Results
- Discussion

Proposal

New TeV scale physics predicted on the basis of naturalness, gauge hierarchy and dark matter wimp considerations



SUSY

Theory and Phenomenology of Sparticles
Drees, Godbole, Roy
(World Scientific 2004)

ADD

Antoniadis (1990)
Arkani-Hamed, Dimopoulos, Dvali (1998)
Perez-Lorenzana (2005)

Importance of SUSY/ADD discrimination

Focus on lepton sector where LHC is not a powerful probe, e.g. $m_{\tilde{e}} \sim 250$ GeV will be hard

Consider event shapes in $e^+e^- \rightarrow l^+l^- \cancel{E}$ at both ILC ($\sqrt{s} = 500$ GeV) and CLIC ($\sqrt{s} = 3$ TeV).

Signal

SUSY

$$\mathcal{O}(10^2 \text{ GeV})$$

MSSM: free parameters $\overbrace{m_{\tilde{e}_{L,R}}, M_1, M_2, \mu, \tan \beta}$


(only gauge coupling unification assumed at $2 + 10^{16} \text{ GeV}$)

ADD

- All extra dims compactified on a d-torus with radius R_c for each
- SM fields on 3-brane, gravity in the bulk
- $(M_s)^{2+d} = (4\pi)^{d/2} \Gamma(d/2) G_N^{-1} (R_c)^{-d}$. only 2-parameters M_s, d .


Motivation for event shape analysis of $e^+e^- \rightarrow \ell^+\ell^- \cancel{E}$.

- Lepton energy spectrum ISR-sensitive, Battaglia et al JHEP 0507 (2005) 033, hep-ph/0507284.




- Angular distributions also tend to flat over most of the range and somewhat similar.

ADD



SUSY



Method of helicity amplitudes CompHEP Freitas, von Manteuffel, Zerwas
(2004)

(obey ward identities) HELAS PYTHIA Choudhury, Datta, Huitu, Konar,
Moretti

(2003)

$$\begin{aligned} \sigma(e^+e^- \rightarrow \ell^+\ell^- \cancel{E}) &= \sum_n \sigma(e^+e^- \rightarrow \ell^+e^- G_N) \\ &\simeq \int_0^{\sqrt{s}} dm \rho(m) \sigma(m), \\ \rho(m) &= \frac{2R_c^d m^{d-1}}{(4\pi)^{d/2} \Gamma(d/2)} \end{aligned}$$

Leading contribution
 $\sigma(e^+e^- \rightarrow \tilde{\ell}_{L,R}^+ \tilde{\ell}_{L,R}^-) BR(\tilde{\ell}_{L,R}^+ \rightarrow \ell_{L,R}^+ \tilde{\chi}_1^0) BR(\tilde{\ell}_{L,R}^- \rightarrow \ell_{L,R}^- \tilde{\chi}_1^0)$
 off-shell $\tilde{\ell}$ contribution insignificant; so are
 $e^+e^- \rightarrow \tilde{\ell}_{L,R}^+ \tilde{\ell}_{L,R}^- \rightarrow \tilde{\chi}^+ \tilde{\chi}^- \nu_\ell \nu_\ell \rightarrow \ell^+ \ell^- \tilde{\chi}_1^0 \tilde{\chi}_1^0 \nu_e \nu_e \bar{\nu}_\ell \bar{\nu}_\ell$
 BR's controlled by (1) LSP composition (2) channel
 access for $m_{\tilde{\ell}} > M_{\tilde{\chi}^\pm}$.

SM background and cuts

cut $e^+e^- \longrightarrow l^+l^-\nu_e\bar{\nu}_e$ from l^+l^-Z : eliminated by missing mass
 W^+W^- : reconstructible modulo 2-fold ambiguity. Subtract!

Improvement with longitudinally polarized beam

Chosen cuts

- Direction of each $l > 10^\circ$ from beam pipe (control collinear singularities from t-channel γ and beamstrahlung)

- $p_T^\ell > 10 \text{ GeV} \text{ (20 GeV)}$
- $p_T^{\text{miss}} > 15 \text{ GeV} \text{ (25 GeV)}$
- $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} > 0.2$
- $5^\circ < \theta_{\ell+\ell^-} < 175^\circ$
- **Miss > 150 GeV (450 GeV)**

Cross-sections and event-shape variables

$\sigma_{\text{SUSY}}[\text{fb}]$ ILC						$\sigma_{\text{ADD}}[\text{fb}]$ ILC				
$\tan \beta = 10$		$m_{\tilde{\ell}}[\text{GeV}]$				$M_S[\text{TeV}]$				d
$M_2, M_1[\text{GeV}]$	$\mu[\text{GeV}]$	155	205	225	245	.75	1.0	1.5	2.0	
200, 100	-400	427	164	59	7.8	1090	345	68	22	2
300, 150	-400	144	137	75	19	455	108	14	3.3	3
400, 200	-150	92	40	13	0.6	202	36	3.2	0.6	4
400, 200	-100	79	32	6.9	0.3	97	13	0.8	0.1	5

$\sigma_{\text{SUSY}}[\text{fb}]$ CLIC						$\sigma_{\text{ADD}}[\text{fb}]$ CLIC				
$\tan \beta = 10$		$m_{\tilde{\ell}}[\text{GeV}]$				$M_S[\text{TeV}]$				d
$M_2, M_1[\text{GeV}]$	$\mu[\text{GeV}]$	700	800	900	1000	4.5	5.0	5.5	6.0	
200, 100	-500	24	19	15	11	124	81	56	39	2
400, 190	-500	22	18	15	11	58	34	21	14	3
600, 290	-500	21	16	13	10	31	16	9.2	5.5	4
800, 380	-500	21	18	12	8	17	8.3	4.2	2.3	5

SM bkgd \sim 36 fb (72 fb)





$$\text{Thrust } T = \max_{\hat{n}} \frac{\sum_a |\hat{n} - \vec{p}_a|}{\sum_a |\vec{p}_a|};$$

↑
↑
↑
 choice of \hat{n} isotropic spiked





$$\frac{1}{2} \leq T \leq 1$$

For our process, $\lambda_3 = 0$; planar events, shape circular rather than spherical for $S_{\max} = 3/4$
 ISR/FSR effects take S_{\max} beyond 3/4.



Results at $\sqrt{s} = 500$ GeV for ADD (M_s, d) and SUSY ($M_2, M_1, \mu, m_{\tilde{\ell}}$)




Results at $\sqrt{s} = 3$ TeV for ADD (M_s, d) and SUSY ($M_2, M_1, \mu, m_{\tilde{\ell}}$)



Results at $\sqrt{s} = 500$ GeV with PYTHIA for SUSY ($M_2, M_1, \mu, m_{\tilde{\ell}}$)



ADD/SUSY discrimination via sphericity maximum at ILC



Discussion

- Event-shape not ISR/FSR sensitive.
- Sphericity better than thrust in SUSY/ADD discrimination.
- Structure in sphericity distribution for SUSY, structureless monotonic fall-off for ADD.
- Location of sphericity peak a robust measure of $m_{\tilde{\ell}}$; ISR/FSR insensitive; not much variation with \sqrt{s} .