


# A Topologic Approach to Particle Flow "PandoraPFA"

Mark Thomson  
University of Cambridge




## This Talk:

- ★ Philosophy
- ★ The Algorithm
- ★ Some First Results
- ★ Conclusions/Outlook

# 1 Philosophy

- ★ Work from the premise that PFA is not a pure ECAL/HCAL clustering problem
- ★ PFA and calorimeter clustering performed together
- ★ Start by applying loose clustering
- ★ Then join clusters using topology



- ★ Algorithm defined by loose cluster + topological rules

# Goals/Framework

- ★ Try to develop “generic” PFA which will take advantage of a high/very high granularity ECAL
- ★ **Clustering** and **PFA** performed in a single algorithm
- ★ Aim for fairly generic algorithm:
  - very few hard coded numbers
  - use **GEAR** to get basic geometry
- ★ Clustering uses tracking information
- ★ Initial clustering is fairly loose → **ProtoClusters**
- ★ Topological linking of ProtoCluster

## Runs in MARLIN framework using:

- ✦ **GEAR (geometry interface)**
- ✦ **Marlin SimpleDigitisation**
- ✦ **Track finding/fitting : TrackCheater**
- ✦ **PFA Utility classes, e.g. Helix class for track extrap. (Alexei R.)**

## ② The Algorithm

### Overview:

#### ★ Preparation

- ★ Isolation cuts, hit ordering, track quality

#### ★ Initial clustering to form ProtoClusters

- ★ **ProtoClusters** are heavyweight object:

- ★ collection of hits
- ★ know how to grow (configured when created)
- ★ information about shape, direction, isPhoton,...
- ★ +much more (not all used)...

#### ★ Cluster association/merging

- ★ **Tight Topological linking** of clusters
- ★ **Looser merging** of clusters
- ★ **Track-driven merging**


#### ★ PFA

- ★ **Final track-cluster matching**


- This talk gives **flavour** of what's done in each stage skipping details

# Preparation I: Extended Hits

- ★ Create internal **ExtendedCaloHits** from **CaloHits**
- ★ **ExtendedCaloHits** contain extra info:
  - ★ pointer to original hit
  - ★ **pseudoLayer** (see below)
  - ★ measure of isolation for other hits
  - ★ is it MIP like (to ID "tracklike objects")
  - ★ actual layer (decoded from **CellID**)
  - ★ **Pixel Size** (from **GEAR**)
- ★ hits are now self describing




- ★ Arrange hits into **PSEUDOLAYERS** (e.g. Chris Ainsley's **MAGIC**)
  - ★ i.e. order hits in increasing depth within calorimeter
  - ★ **PseudoLayers** follow detector geometry




# Preparation II: Isolation

- ★ Divide hits into isolated and non-isolated
  - ★ Only cluster non-isolated hits
  - ★ "Cleaner"/Faster clustering
  - ★ Significant effect for scintillator HCAL
- 
- ★ Removal of isolated hits degrades HCAL resolution
  - ★ e.g. D10scint:
    - 50 %/ $\sqrt{E}/\text{GeV}$  →
    - 60 %/ $\sqrt{E}/\text{GeV}$



# Preparation III: Tracking




- ★ Use MARLIN TrackCheater
- ★ Tracks formed from MC Hits in TPC/FTD/VTX
- ★ HelixFit (Alexei R)  $\Rightarrow$  track params
- ★ Cuts (primary tracks):
  - ◆  $|d_0| < 5$  mm
  - ◆  $|z_0| < 5$  mm
  - ◆  $>4$  non-Si hits

## + $V_0$ and Kink finding:

- ★ Track resolution better than cluster
- ★ Improves PFA performance by  $\sim 2$  %

# PandoraPFA Clustering II

- ★ Start at inner layers and work outward
- ★ Associate Hits with existing Clusters
- ★ If multiple clusters "want" hit then **Arbitrate**
- ★ Step back **N** layers until associated
- ★ Then try to associate with hits in current layer (M pixel cut)
- ★ If no association made form new Cluster
- ★ + tracks used to seed clusters





# Cluster Association


- ✦ By design clustering errs on side of caution  
i.e. clusters tend to be split
- ✦ Philosophy: easier to put things together than split them up
- ✦ Clusters are then associated together in two stages:
  - 1) Tight cluster association - clear topologies
  - 2) Loose cluster association - catches what's been missed but rather crude




## Photon ID

- ★ Photon ID plays important role
- ★ Simple "cut-based" photon ID applied to all clusters
- ★ Clusters tagged as photons are immune from association procedure - just left alone


Won't merge




Won't merge



Could get merged




# Cluster Association I : track merging




# Cluster Association II : Backscatters

- ★ Forward propagation clustering algorithm has a major drawback: back scattered particles form separate clusters



# Cluster association III : MIP segments


- ★ Look at clusters which are consistent with having tracks segments and project backwards/forward



- ★ Apply tight matching criteria on basis of projected track [NB: + track quality i.e.  $\chi^2$ ]

# Cluster Association Part II


- Have made very clear cluster associations
- Now try “cruder” association strategies
- **BUT first associate tracks to clusters (temporary association)**
- Use track/cluster energies to “veto” associations, e.g.




Provides some protection against silly mistakes

# Sledgehammer Cluster Association

**Proximity**



**Shower Cone**



**+Track-Driven Shower Cone**

Apply looser cuts if have low E cluster associated to high E track

# Performance








Figure of Merit:



- ★ Find smallest region containing 90 % of events
- ★ Determine rms in this region

More robust than fitting double Gaussian

# Preliminary Results : Z → uds events



## \* RMS of Central 90 % of Events


B-Field	$\sigma_E/E = \alpha\sqrt{(E/\text{GeV})}$
2 Tesla	<b>37.8±0.4%</b>
4 Tesla	<b>35.9±0.4 %</b>
6 Tesla	<b>37.4±0.4 %</b>

✦ only weakly depends on B




# Results : Z uds events Angular dependence

★ Plot resolution vs generated polar angle of qq system



★ In barrel : 34-36 %/  $\sqrt{E(\text{GeV})}$

# Radial Dependence



Model	all angles	$\cos\theta < 0.8$
Tesla $r_{\text{tpc}} = 1690, I_{\text{tpc}} = 2730$	$35.3 \pm 0.6$	$33.9 \pm 0.7$
Tesla $r_{\text{tpc}} = 1890, I_{\text{tpc}} = 2930$	$36.6 \pm 0.6$	$35.8 \pm 0.6$
LDC $r_{\text{tpc}} = 1580, I_{\text{tpc}} = 2200$	$37.1 \pm 0.6$	$35.7 \pm 0.6$
LDC $r_{\text{tpc}} = 1380, I_{\text{tpc}} = 2000$	$39.7 \pm 0.6$	$38.9 \pm 0.7$

★ Some evidence that going to small radii gives worse performance  
 ★ BUT... don't take too seriously, Z events + algorithm not perfect

# Outlook

- ★ Looks promising - **good performance for 91.2 GeV Z events**
- ★ **Can be improved:**
  - ✦ still a few features (i.e. does something silly)
  - ✦ some problems with tracking
  - ✦ photon ID is quite basic
  - ✦ + some new ideas (for high density events)
- ★ Code runs within **Marlin** framework and is “nearly” ready for release - aim to optimise on higher energy jets
- ★ + code needs tidying up
  - ✦ started with decent OO structure
  - ✦ then grew organically...
- ★ Aim to have complete algorithm ~ end April
- ★ **Hopefully, soon ready to start full simulation detector optimisation studies**