

Anomalous Higgs Couplings at $e\gamma$ Collider

Mamta

*Department of Physics
S.G.T.B. Khalsa College
University of Delhi
Delhi, India*

collaboration with

Rajyoti Choudhury

March 12, 2006

Introduction

examine the resolving power of an $e\gamma$ collider in context of WWH vertices.

The coupling of Higgs to gauge bosons may be directly related to EWSB.

Standard Model:

$$g_{WWH}^{SM} = e \cot \theta_W M_Z$$

$$g_{ZZH}^{SM} = 2e \frac{M_Z}{\sin 2\theta_W}$$

Coming from the Higgs kinetic term.

If we accept the SM to be an effective low energy description of some other theory, higher dimensional terms are allowed.

$$\mathcal{L}_{eff} = \sum_n \frac{f_n}{\Lambda^2} \mathcal{O}_n$$

where f_n 's are dimensionless “anomalous couplings”, and \mathcal{O}_n are dimension-6 operators constructed from vector bosons and/or Higgs boson fields.

Introduction

Demanding Lorentz invariance and gauge invariance, the most general coupling structure can be expressed as

$$\Gamma_{\mu\nu} = g_V \left[a_V g_{\mu\nu} + \frac{b_V}{m_V^2} (k_{1\mu} k_{2\nu} - g_{\mu\nu} k_1 \cdot k_2) + i \frac{\tilde{b}_V}{m_V^2} \epsilon_{\mu\nu\alpha\beta} k_1^\alpha k_2^\beta \right]$$

where k_i are the momenta of two W's or Z's. In general, each of these couplings can be complex. In the context of SM, at the tree level,

$$a_W^{SM} = a_Z^{SM} = 1 \text{ and } b_W = \tilde{b}_W = 0$$

Shashu, Rohini, Ritesh and Debajyoti [hep-ph/0509070]

Cross-section

the process

$$e^- \gamma \rightarrow \nu_e W^- H$$

followed by the decays

$\rightarrow b \bar{b}$ with branching fraction ~ 0.9

$\rightarrow q \bar{q}$ with branching fraction ~ 0.68




Figure 1: Feynman Diagrams for $e^- \gamma \rightarrow \nu_e W^- h$

Cross-section

$$m_H = 120 \text{ GeV}$$

$$\sqrt{s_{e e}} = 1 \text{ TeV}$$

We assume that

$$a_W = 1 + \Delta a_W$$

We restrict to unpolarised beams

Retaining only the contributions upto the lowest non-trivial order, arising from terms linear in the additional couplings, the cross-section may be written as

$$\sigma = (1 + 2 \Delta a_W) \sigma_0 + \mathcal{R}(b_V) \sigma_1 + \mathcal{I}(b_V) \sigma_2 + \mathcal{R}(\tilde{b}_V) \sigma_3 + \mathcal{I}(\tilde{b}_V) \sigma_4$$

Kinematical Cuts

To ensure detectability of the final products, we require that,

$$-3.0 \leq \eta_j \leq 3.0 \quad \text{for rapidity of each jet}$$

$$p_T \leq 10 \text{ GeV} \quad \text{for each jet}$$

$$\Delta R_{j_1 j_2} \geq 0.7 \quad \text{for each pair of jets}$$

$$\text{where } (\Delta R_{j_1 j_2})^2 \equiv (\Delta\phi)^2 + (\Delta\eta)^2$$

$$p_T^{\text{miss}} \geq 20 \text{ GeV}$$

Apart from the cross-section, the kinematical distributions are different for different splittings.

Couplings Δa_W , $\mathcal{R}(b_W)$ and $\mathcal{I}(\tilde{b}_W)$

Higgs boson of mass 120 GeV and $\sqrt{s_{ee}} = 1$ TeV, the rates in femtobarns with polarised beams are:

$$\sigma = 29.3(1 + 2 \Delta a_W) - 101.6 \mathcal{R}(b_W) + 4.92 \mathcal{I}(\tilde{b}_W)$$

cut	cross-section σ in femto barns
$\mathcal{C}_1: p_T(H) \geq 100$ GeV	$11.92 (1 + 2 \Delta a_W) - 43.4 \mathcal{R}(b_W) - 0.31 \mathcal{I}(\tilde{b}_W)$
$\mathcal{C}_2: p_T(W) \geq 100$ GeV and $ \sin \phi_{HW} \leq 0.4$	$4.27 (1 + 2 \Delta a_W) - 27.5 \mathcal{R}(b_W) + 0.59 \mathcal{I}(\tilde{b}_W)$
$\mathcal{C}_3: p_T(W) \leq 100$ GeV and $p_T^{\text{miss}} \leq 100$ GeV	$18.6 (1 + 2 \Delta a_W) - 41.2 \mathcal{R}(b_W) + 0.41 \mathcal{I}(\tilde{b}_W)$
$\mathcal{C}_4: p_T(H) \leq 100$ GeV	$17.33 (1 + 2 \Delta a_W) - 58.2 \mathcal{R}(b_W) + 5.21 \mathcal{I}(\tilde{b}_W)$
$\mathcal{C}_5: p_T(W) \geq 100$ GeV and $ \sin \phi_{HW} \geq 0.4$	$2.15 (1 + 2 \Delta a_W) - 20.1 \mathcal{R}(b_W) + 3.24 \mathcal{I}(\tilde{b}_W)$
$\mathcal{C}_6: p_T(W) \geq 100$ GeV and $p_T(\text{miss}) \geq 100$ GeV	$1.265 (1 + 2 \Delta a_W) - 13.85 \mathcal{R}(b_W) + 2.40 \mathcal{I}(\tilde{b}_W)$

Couplings Δa_W , $\mathcal{R}(b_W)$ and $\mathcal{I}(\tilde{b}_W)$

Strong correlations between Δa_W and $\mathcal{R}(b_W)$ Neglecting contribution of $\mathcal{I}(\tilde{b}_W)$, after imposition of the cut $\mathcal{C}_i, (i = 1, 2, 3)$, the rate depends on a combination of Δa_W and $\mathcal{R}(b_W)$

$$\sigma(\mathcal{C}_i) = \sigma_{SM}(1 + \eta_i)$$

Thus, for an **integrated luminosity of 500 fb^{-1}** , the lack of deviation from the SM expectation values would (at 3σ level) give limits on η_i as

$$\eta_1 = |2 \Delta a_W - 3.64 \mathcal{R}(b_W)| \leq 0.039$$


$$\eta_2 = |2 \Delta a_W - 6.44 \mathcal{R}(b_W)| \leq 0.064$$

$$\eta_3 = |2 \Delta a_W - 2.22 \mathcal{R}(b_W)| \leq 0.031$$

Individual Limits on the couplings are

Coupling		\mathcal{C}_1	\mathcal{C}_2	\mathcal{C}_3
$ \Delta a_W $	\leq	0.020	0.032	0.015
$ \mathcal{R}(b_W) $	\leq	0.011	0.010	0.014

Couplings Δa_W , $\mathcal{R}(b_W)$ and $\mathcal{I}(\tilde{b}_W)$



Couplings Δa_W , $\mathcal{R}(b_W)$ and $\mathcal{I}(\tilde{b}_W)$




from the cuts \mathcal{C}_4 , \mathcal{C}_5 and \mathcal{C}_6

$$| 12.74 \Delta a_W - 58.2 \mathcal{R}(b_W) + 5.21 \mathcal{I}(\tilde{b}_W) | \leq 0.33$$

$$| 6.3 \Delta a_W - 20.1 \mathcal{R}(b_W) + 3.24 \mathcal{I}(\tilde{b}_W) | \leq 0.20$$

$$| 2.54 \Delta a_W - 13.85 \mathcal{R}(b_W) + 2.4 \mathcal{I}(\tilde{b}_W) | \leq 0.15$$

Couplings Δa_W , $\mathcal{R}(b_W)$ and $\mathcal{I}(\tilde{b}_W)$



Couplings $\mathcal{I}(b_W)$ and $\mathcal{R}(\tilde{b}_W)$

get contribution from these couplings, we restrict the W-bosons to be in the upper hemisphere *i.e.* $\sin \phi_{HW} \geq 0$. This partial cross-section is

$$\begin{aligned} (\sin \phi_{HW} \geq 0) &= 14.7 (1 + 2 \Delta a_W) - 50.79 \mathcal{R}(b_W) + 2.48 \mathcal{I}(\tilde{b}_W) + 7.26 \mathcal{I}(b_W) \\ &+ 26.58 \mathcal{R}(\tilde{b}_W) \end{aligned}$$

get limits on the imaginary couplings, we construct the asymmetry

$$\begin{aligned} \mathcal{A}_1 &= \frac{\sigma_{\sin \phi_{HW} \geq 0} - \sigma_{\sin \phi_{HW} \leq 0}}{\sigma_{\sin \phi_{HW} \geq 0} + \sigma_{\sin \phi_{HW} \leq 0}} \\ &= \frac{2 \left[7.26 \mathcal{I}(b_W) + 26.6 \mathcal{R}(\tilde{b}_W) \right]}{2 \left[14.75 (1 + 2 \Delta a_W) - 50.79 \mathcal{R}(b_W) + 2.48 \mathcal{I}(\tilde{b}_W) \right]} \end{aligned}$$

with a further cut $\cos \theta_H \leq 0$, we get

Couplings $\mathcal{I}(b_W)$ and $\mathcal{R}(\tilde{b}_W)$

$$\begin{aligned}\mathcal{A}2 &= \mathcal{A}1(\cos \theta_H \leq 0) \\ &= \frac{2 \left[0.79 \mathcal{I}(b_W) + 7.35 \mathcal{R}(\tilde{b}_W) \right]}{2 \left[3.36 (1 + 2 \Delta a_W) - 14.27 \mathcal{R}(b_W) + 0.21 \mathcal{I}(\tilde{b}_W) \right]}\end{aligned}$$

Keeping only one of the couplings to be non-zero we obtain the following limits on them for $\mathcal{A}2$:

$$|\mathcal{I}(b_W)| \leq 0.032$$


$$|\mathcal{R}(\tilde{b}_W)| \leq 0.019$$

where the fluctuation in the measurement of asymmetry is taken to be

$$(\delta\mathcal{A})^2 = \frac{1 - \mathcal{A}_{SM}^2}{\sigma_{SM} \mathcal{L}}$$

3 σ level.

Couplings $\mathcal{I}(b_W)$ and $\mathcal{R}(\tilde{b}_W)$



Summary

Contributions proportional to Δa_W and $\mathcal{R}(b_W)$ are intertwined and hence can't be separated.

Similar is the case with couplings $\mathcal{I}(b_W)$ and $\mathcal{R}(\tilde{b}_W)$.

Limits obtained on couplings $\mathcal{I}(b_W)$ and $\mathcal{R}(\tilde{b}_W)$ are weak.

If it were possible to use the CP-conjugate process with $e^+\gamma$ initial state, then it would be possible to construct observables that would isolate the CP-odd couplings and can be used to probe them. The work is under the progress.

Using polarised photons can improve the limits and that work is under progress.