

# Single top-quark production with the matrix-element method in NLO accuracy

Till Martini

Humboldt-Universität zu Berlin – Phänomenologie der Elementarteilchenphysik

“25th International Workshop on Deep Inelastic Scattering and Related Topics” – Birmingham – 4.4.2017

in collaboration with Peter Uwer

based on [\[JHEP 1509 \(2015\) 083\]](#) and ongoing work



Bundesministerium  
für Bildung  
und Forschung



# Matrix Element Method (MEM) in a nutshell [Kondo '88,'91]


For the collision  $A + B \rightarrow a_1(p_1) + a_2(p_2) + \dots + a_n(p_n)$

the differential cross section

$$\frac{d\sigma_n}{d^4 p_1 \dots d^4 p_n} \propto \frac{1}{2s} |\mathcal{M}(p_1, \dots, p_n)|^2 dR_n$$

$$dR_n(p_1, \dots, p_n) = (2\pi)^4 \delta(P - \sum_i p_i) \prod_j \frac{d^3 p_j}{(2\pi)^3 2E_j}$$

is a measure for the probability to observe the final state in the inf. phase space region  $dR_n$  located at  $(p_1, \dots, p_n)$



$$\vec{p} = (p_1, \dots, p_n)$$

# Matrix Element Method (MEM) in a nutshell [Kondo '88,'91]

Collecting generic partonic final state variables in  $\vec{x}$ :

e.g.  $\vec{x} = (E_i, \theta_j, \dots)$ :  $\boxed{\frac{d\sigma_n}{d^4 p_1 \dots d^4 p_n} \rightarrow \frac{d\sigma_n}{d\vec{x}}}$

Experimentally we observe hadronic variables  $\vec{y}$  instead of partonic variables  $\vec{x}$ .

Probability to measure  $\vec{y}$ :

$$\mathcal{P}(\vec{y}) = \frac{1}{\sigma} \int d\vec{x} \frac{d\sigma}{d\vec{x}} W(\vec{x}, \vec{y})$$

Transfer function  $W(\vec{x}, \vec{y})$ :

- ▶ Probability to experimentally observe a hadronic event  $\vec{y}$  given a partonic event  $\vec{x}$
- ▶ Determined by experiments through simulations
- ▶  $\delta$ -functions in the following

# Matrix Element Method (MEM) in a nutshell [Kondo '88,'91]


Extraction of model parameters  $\Omega$  from data by maximizing a likelihood  $\propto$  differential cross section ( $\propto |\mathcal{M}^{\text{LO}}|^2$ ):

**Likelihood** as a function of  $\Omega$  for event sample  $\{\vec{x}_i\}$

$$\mathcal{L}^{\text{LO}}(\Omega) = \prod_i \frac{1}{\sigma^{\text{LO}}(\Omega)} \int d\vec{y} \frac{d\sigma^{\text{LO}}(\Omega)}{d\vec{y}} \underbrace{W(\vec{x}_i, \vec{y})}_{\substack{\text{transfer function,} \\ \text{here: } =\delta(\vec{x}_i - \vec{y})}} = \prod_i \frac{1}{\sigma^{\text{LO}}(\Omega)} \frac{d\sigma^{\text{LO}}(\Omega)}{d\vec{x}_i}$$

Maximizing wrt  $\Omega$  yields estimator  $\hat{\Omega}$ : 
$$\mathcal{L}^{\text{LO}}(\hat{\Omega}) = \sup_{\Omega} \mathcal{L}^{\text{LO}}(\Omega)$$

All information from event used  $\implies$  **most efficient estimator!**



e.g. top mass measurement at Tevatron  
[DO: Nature 429, 638], [CDF: PRD 50, 2966]  
based on  $O(70)$  events!

# MEM beyond LO:

## Experimental analysis so far restricted to LO

Steps towards MEM@NLO:

- ▶ Effects of real radiation [Alwall,Freitas,Mattelaer '11]
- ▶ Final states without strongly interacting particles [Campbell,Giele,Williams '12], [Campbell,Ellis,Giele,Williams '13]
- ▶ Steps towards final states with strongly interacting particles [Campbell,Giele,Williams '13]
- ▶ Complete algorithm [Martini,Uwer '15]
- ▶ Proposed generalisation [Baumeister,Weinzierl '17]

# MEM beyond LO: NLO cross section at parton level

$$\sigma^{\text{NLO}} = \underbrace{\int dR_n \frac{d\sigma^{\text{B}}}{dR_n}}_{\text{Born}} + \underbrace{\int dR_n \frac{d\sigma^{\text{V}}}{dR_n}}_{\text{virtual}} + \underbrace{\int dR_{n+1} \frac{d\sigma^{\text{R}}}{dR_{n+1}}}_{\text{real}}$$

separately IR divergent

Jet Physics: observed final states are jets


partonic events  $\rightarrow$  "Jet events"

New in NLO:

- ▶ Infrared and collinear divergences in virtual and real corrections  $\xrightarrow{\text{KLN theorem}}$  mutual cancelation

- ▶  $n + 1$  particle phase space due to real corrections:

Non-trivial mapping: parton momenta  $\leftrightarrow$  jet momenta



Born and virtual:  $J_i = p_i$  trivial but real:  $J_i = \tilde{J}_i(p_1, \dots, p_{n+1})$

## Likelihood at NLO: 3 major obstacles

$$\mathcal{L}^{\text{NLO}}(\Omega) = \prod_i \frac{1}{\sigma_{n\text{-jet}}^{\text{NLO}}(\Omega)} \left( \frac{d\sigma_{n \rightarrow n\text{-jet}}^{\text{B+V}}(\Omega)}{dJ_1 \dots dJ_n} + \frac{d\sigma_{n+1 \rightarrow n\text{-jet}}^{\text{R}}(\Omega)}{dJ_1 \dots dJ_n} \right) \Big|_{\text{event } i}$$

- ▶ Born+virtual and real contribution separately IR divergent


NEED: Point-wise cancelation in jet phase space

- ▶ Integration over jet cones in real contribution introduces  $\delta$ -functions  $\rightarrow$  numerically problematic!

NEED: factorisation of real phase space

- ▶ Born+virtual matrix elements only defined for 'Born kinematics'

NEED: clustered jets obeying 'Born kinematics'



## Proposal: Replace $2 \rightarrow 1$ clustering by $3 \rightarrow 2$ clustering

$3 \rightarrow 2$  clustering  $p_i, p_j, p_k \rightarrow \tilde{J}_{ij}, \tilde{J}_k$   
can meet **all 3** requirements at the same time



Using  $3 \rightarrow 2$  jet algorithm instead of  $2 \rightarrow 1$   
allows to overcome the 3 obstacles

$3 \rightarrow 2$  clusterings inspired by Catani-Seymour dipole subtraction method [Catani, Seymour '97], [Catani, Dittmaier, Seymour, Trocsanyi '02]

# Jet event weight at NLO

Phase space factorisation allows to define an event weight (differential jet cross section) at NLO

$$\frac{d\sigma_{n\text{-jet}}^{\text{NLO}}(\Omega)}{dJ_1 \dots dJ_n} = \underbrace{\frac{d\sigma^{\text{B+V}}(\Omega)}{dJ_1 \dots dJ_n}}_{\text{still IR divergent}} + \overbrace{\int dR_{\text{unres}}(\Phi)}^{\text{3dim integration}} \underbrace{\frac{d\sigma^{\text{R}}(\Omega)}{dp_1 \dots dp_{n+1}}}_{\text{still IR divergent}}$$

finite through subtraction/slicing

Mutual cancelation of IR-divergences from the **virtual** and the **real** part has to be carried out by a suitable method (e.g. phase space slicing)

# Validation

We consider the s-channel or t-channel production of single top quarks in association with a light jet at the LHC

$$p + p \rightarrow t + j (+X) \quad @ \text{ NLO QCD}$$

optional veto on additional jet emission!


Note: top decay not included, tops are treated as tagged top jets

To Check:

1. Generation of real phase space
2. Generation of unweighted events (at NLO accuracy!)

# Validation 1: Phase space generation


Calculate jet distributions in NLO accuracy using conventional approach (parton level MC + 3  $\rightarrow$  2 jet alg.) and compare with distributions obtained from  $\frac{d\sigma_{n\text{-jet}}^{\text{NLO}}(\Omega)}{dJ_1 \dots dJ_n}$  with factorised jet phase space



## Validation 2: Unweighted events

Fill histograms with unweighted events generated according to

$\rho = \frac{d\sigma_{n\text{-jet}}^{\text{NLO}}(\Omega)}{dJ_1 \dots dJ_n}$  and compare with jet distributions in NLO accuracy obtained from conventional approach (parton MC + 3  $\rightarrow$  2 clus.)



# Application: Matrix Element Method at NLO

(example:  $p + p \rightarrow t + j$  via s-/t-channel @ NLO QCD)

Toy experiment: Generate sample of  $N$  unweighted NLO  $tj$  events

$$\vec{x}_i = (\eta_t, E_j, \eta_j, \phi_j)_i \quad \text{with } \Omega = m_t = m_t^{\text{true}} = 173.2 \text{ GeV}$$


NLO likelihood function for event sample

$$\mathcal{L}^{\text{NLO}}(m_t) = \prod_i^N \mathcal{L}^{\text{NLO}}(\vec{x}_i | m_t) = \left( \frac{1}{\sigma^{\text{NLO}}(m_t)} \right)^N \prod_{i=1}^N \left( \frac{E_j^2 \cosh(\eta_t)}{2 s E_t \cosh^3(\eta_j)} \frac{d\sigma^{\text{NLO}}}{d^4 J_t d^4 J_j}(m_t) \right) \Big|_i$$

Find minimum of negative logarithm of NLO likelihood  
("Log-Likelihood") to obtain estimator  $\hat{m}_t$  for top mass

$$-\log \mathcal{L}^{\text{NLO}}(\hat{m}_t) = \inf_{m_t} \left( -\log \mathcal{L}^{\text{NLO}}(m_t) \right)$$


# MEM at NLO: s-channel, no veto on additional jet



NLO analysis recovers  $m_t^{\text{true}}$  but biased if MEM@LO is used!

- ▶  $\approx 17k$  events allow already precise mass determination
- ▶ Impact of scale variation reduced in NLO analysis
- ▶ Bias in LO estimator covered by scale variation between  $[\frac{\mu_0}{2}, 2\mu_0]$

# MEM at NLO: t-channel, veto on additional jet



NLO analysis recovers  $m_t^{\text{true}}$  but biased if MEM@LO is used!

- ▶ Impact of scale variation significantly reduced in NLO analysis
- ▶ Shift in LO analysis not covered by scale variation  $\Rightarrow$  naive variation between  $[\frac{\mu_0}{2}, 2\mu_0]$  in LO analysis may not give reliable estimate of NLO effects
- ▶ Large shift observed in LO analysis would require significant calibration with related uncertainties

# Conclusion

3 → 2 jet clustering algorithm:

- ▶ Uniquely maps real corrections onto Born kinematics

Allows:

- ▶ Evaluation of event weights for jet events in NLO accuracy
- ▶ Generation of unweighted events at NLO
- ▶ Application of MEM at NLO
  - ▶ Extraction of  $m_t$  with NLO likelihood from NLO single top events: **Perfect agreement with input value!**
  - ▶ Extraction with Born likelihood: **Large deviation from input value possible (not necessarily covered by scale variation)!**
  - ▶ Renormalization scheme well-defined in MEM at NLO (extracted mass = pole mass)

**Outlook:** MEM at NLO for top+Higgs

# BackUp: Ingredients of iterative 2 $\rightarrow$ 1 jet algorithms

## 1. Resolution:


e.g. 
$$y_{ij} = \min(p_i^{\perp 2\rho}, p_j^{\perp 2\rho}) \frac{\Delta_{ij}}{R^2}, \quad \Delta_{ij} = (y_i - y_j)^2 + (\phi_i - \phi_j)^2$$

( $\rho = 1$ : “kt-algorithm”,  $\rho = -1$ : “anti-kt-algorithm”)

## 2. Recombination/Clustering:

If  $y_{ij} < y_{\text{cut}}$ :

$p_i, p_j$  are clustered to jet  $\tilde{J}_{ij}$



e.g. 
$$\tilde{J}_{ij} = p_i + p_j$$

$\rightarrow$  respects 4-momentum conservation,  
violates on-shell condition

or 
$$\tilde{J}_{ij} = \vec{p}_i + \vec{p}_j, \quad \tilde{J}_{ij}^0 = \sqrt{\tilde{J}_{ij}^2 + m_{ij}^2}$$

$\rightarrow$  respects on-shell condition,  
violates energy conservation

(“p-scheme”)

# BackUp: $3 \rightarrow 2$ jet algorithm as an augmented $2 \rightarrow 1$ algorithm

- ▶ Use resolution criterium  $y_{ij}$  of the  $2 \rightarrow 1$  algorithm to pick final state particle to be clustered with final state particle or beam (“**emitter**”)
- ▶ Choose final state particle or beam as “**spectator**”
- ▶ 4 different types of mappings  
(**emitter**,**spectator**) = (**final**,**final**), (**final**,**initial**), (**initial**, **initial**), (**initial**,**final**)
- ▶ Respective clusterings for massless and massive particles already worked out in Catani-Seymour dipole subtraction method [Catani,Seymour '97], [Catani,Dittmaier,Seymour,Trocsanyi '02] (let's use those!)


# BackUp: Modified clustering

[Catani, Seymour '97],

[Catani, Dittmaier, Seymour, Trocsanyi '02]

(example: massless final state clustering with final state spectator)

$$(p_i, p_j, p_k) \rightarrow (\tilde{p}_{ij} = p_i + p_j - \frac{y}{1-y} p_k, \tilde{p}_k = \frac{1}{1-y} p_k) \rightarrow (\tilde{J}_{ij}, \tilde{J}_k)$$



$$\tilde{J}_{ij} + \tilde{J}_k = p_i + p_j + p_k$$

and  $\tilde{J}_{ij}^2 = m_{ij}^2, \tilde{J}_k^2 = m_k^2$

Phase space factorises:  $dR_{n+1}(p_i, p_j, p_k) = dR_n(\tilde{J}_{ij}, \tilde{J}_k) dR_{\text{unres}}(\Phi)$

integrating over  $dR_{\text{unres}}(\Phi = \{\phi, z, y\}) = \frac{\tilde{J}_{ij} \tilde{J}_k}{2(2\pi)^3} d\phi dz dy (1-y)$

generates all  $p_i, p_j, p_k$  with  $(p_i, p_j, p_k) \xrightarrow{!} (\tilde{J}_{ij}, \tilde{J}_k)$

# BackUp: Modified clustering: Generation of real phase space

(e.g. massless final-final clustering à la Catani-Seymour)

**3→2 jet clustering:**  $(p_1, \dots, p_{n+1}) \rightarrow (\tilde{J}_1, \dots, \tilde{J}_n, \Phi)$

$$\tilde{J}_{ij} = p_i + p_j - \frac{y}{1-y} p_k, \quad \tilde{J}_k = \frac{1}{1-y} p_k, \quad \tilde{J}_m = p_m, \quad m \neq i, j, k$$

$$\Phi = \left\{ \phi, z = \frac{p_i \cdot p_k}{p_k \cdot (p_i + p_j)}, y = \frac{p_i \cdot p_j}{p_i \cdot p_j + p_k \cdot (p_i + p_j)} \right\}$$

**Opposite direction required:**  $(\tilde{J}_1, \dots, \tilde{J}_n, \Phi) \rightarrow (p_1, \dots, p_{n+1})$


$$p_i + p_j = \tilde{J}_{ij} + y \tilde{J}_k, \quad p_k = (1-y) \tilde{J}_k, \quad p_m = \tilde{J}_m, \quad m \neq i, j, k$$

$$\Phi = \left\{ \phi, z = \frac{p_i \cdot \tilde{J}_k}{\tilde{J}_{ij} \cdot \tilde{J}_k}, y = \frac{p_i \cdot p_j}{\tilde{J}_{ij} \cdot \tilde{J}_k} \right\}$$

Phase space factorises:  $dR_{n+1}(p_1, \dots, p_{n+1}) = dR_n(\tilde{J}_1, \dots, \tilde{J}_n) dR_{\text{unres}}(\Phi)$

$$\implies dR_{\text{unres}} \longrightarrow \left( p_1(\tilde{J}_1, \dots, \tilde{J}_n, \Phi), \dots, p_{n+1}(\tilde{J}_1, \dots, \tilde{J}_n, \Phi) \right)$$

# BackUp: MEM at NLO: top-quark mass extraction from $e^+e^- \rightarrow t\bar{t}$



NLO analysis recovers  $m_t^{\text{true}}$  but significant difference if MEM@LO is used!