



Search for vector like quarks and heavy resonances decaying to top quarks

Deep Inelastic Scattering conference 2017



Clément Camincher


On behalf of the ATLAS collaboration

5/04/2017

Why using top quarks to probe new physics ?


Top quarks play an important role at the electroweak energy scale

- ▶ Top quarks are very heavy $\approx 173 \text{ GeV}$
- ⇒ Strong Yukawa coupling with the Higgs boson
- ⇒ Major piece in the naturalness issue
- ▶ Many BSM scenarios predict:
 - ▶ New couplings involving tops $\Rightarrow t\bar{t}$ resonances searches
 - ▶ Top partners \Rightarrow VLQ searches




ATLAS a top detector at the LHC

- ▶ At 13TeV, the LHC is a top-quark factory (1 $t\bar{t}$ pair produced every second)
- ▶ ATLAS detector optimized to find top-quarks
 - ▶ Very good electron and muon reconstruction and identification
Present in 39% of top decays
 - ▶ Full azimuthal coverage to measure neutrino transverse component via E_T^{miss}
 - ▶ Close tracker layer (IBL) to select displaced vertexes from b-jets




Top quarks play an important role at the electroweak energy scale

- ▶ Top quarks are very heavy $\approx 173 \text{ GeV}$
- \Rightarrow Strong Yukawa coupling with the Higgs boson
- \Rightarrow Major piece in the naturalness issue
- ▶ Many BSM scenarios predict:
 - ▶ New couplings involving tops $\Rightarrow t\bar{t}$ resonances searches
 - ▶ Top partners \Rightarrow VLQ searches



Concept of the $t\bar{t}$ resonances searches

- ▶ Many BSM scenarios predict new particles (X) decaying into a pair of tops
- ▶ Z' technicolor assisted topcolor
- ▶ Randall Sundrum \rightarrow gluon/graviton: g_{KK}/G_{KK}
- ▶ 2 Higgs Doublet Models \rightarrow heavy pseudoscalar/scalar A/H



Analysis strategy :
Build a model independent analysis


1. Select events
2. Reconstruct $t\bar{t}$ invariant mass
3. Scan $m_{t\bar{t}}$ to find an excess/deficit

- ▶ Lepton + jet channel choice
- ▶ Lepton \Rightarrow multijet background suppression (trigger more efficient on leptons)
- ▶ Jets \Rightarrow keep good statistics level (high branching ratio)

Lepton +jets boosted $t\bar{t}$ search


ATLAS-CONF-2016-014

- ▶ Results using 3.2 fb^{-1} of 13 TeV data
- ▶ Analysis dedicated to probe heavy resonances
- ⇒ Producing boosted top
- ⇒ collimated decay products




▶ Boosted leptonic top selection

- ▶ Select exactly 1 muon or 1 electron
- ▶ b-jet selected as highest p_T jet in $\Delta R(\ell; \text{jet}) < 1.5$
- ▶ Neutrino transverse momentum estimated with E_T^{miss}
- ▶ Neutrino longitudinal momentum estimated constraining the system with the on-shell W mass.



High top p_T




▶ Boosted hadronic top selection


- ▶ 1 top-tagged (80%) fat jet (anti-kt with $R=1.0$)
- ▶ Large jets are trimmed by removing kt sub-jets with p_T fraction below 5%
- ▶ Tagging \Leftrightarrow cut on jet mass and τ_{32} "nsubjettiness" ratio (describing the internal structure of the jet)

Lepton +jets boosted $t\bar{t}$ analysis

- ▶ Dominant background is SM $t\bar{t}$
- ▶ Multijet estimated with data
- ▶ W+jets estimation
 - ▶ Shape from MC
 - ▶ Scaling from data
- ▶ Main systematics from fat jet modeling




Electron selection






Muon selection

- ▶ Fit simultaneously both lepton distributions
- ▶ No deviation found
- ▶ Limit set on a $Z' TC2$
- ▶ Z' with $\Gamma/m = 1.2\%$ excluded in the region $0.7 < m_{Z'} < 2\text{TeV}$



Re-interpretation 8 TeV (20 fb^{-1}) $t\bar{t}$ resonances search

- ▶ 2HDM model predicts Scalar (H) or Pseudoscalar (A) decaying into $t\bar{t}$
- ▶ Large interference (I) with SM $gg \rightarrow t\bar{t}$ production
- ▶ It would require signal (S)+ (I) + background (B) simulation at parton level for each mass point




- ▶ Generator modified to produce only S+I
- ▶ $m_{t\bar{t}}$ Signal modeling at parton level versus Signal + Interference modeling.
- ▶ Validated with S+I+B generation for some parameter space points

Difference treated as systematics uncertainties (0.4%)


$H/A \rightarrow t\bar{t}$ analysis setup and results

Resolved selection




- ▶ Exactly 1 lepton (electron or muon)
- ▶ 4 jets (1 or 2 b-tags)
- ▶ Jet ambiguity solved by minimizing a χ^2 over jet combinations


Invariant mass muon selection




electron selection



Limit on the signal strength μ versus $\tan(\beta)$, $m_{H/A} = 500 \text{ GeV}$




Pseudo scalar



Scalar

Top quarks play an important role at the electroweak energy scale


- ▶ Top quarks are very heavy $\approx 173 \text{ GeV}$
- \Rightarrow Strong Yukawa coupling with the Higgs boson
- \Rightarrow Major piece in the naturalness issue
- ▶ Many BSM scenarios predict:
 - ▶ New couplings involving tops $\Rightarrow t\bar{t}$ resonances searches
 - ▶ Top partners \Rightarrow VLQ searches



Vector Like Quarks (VLQ) searches

- ▶ To solve the hierarchy problem many BSM scenarios predict VLQ
- ▶ They don't receive mass through BEH mechanism
- ▶ VLQ would contribute to Higgs mass loop correction

ATLAS search for VLQ ($T\bar{T}$) pair production



- ▶ T has same electric charge than top-quarks ($Q=+2/3$)
- ▶ Considering only T decays :

$$T \rightarrow Zt, Ht, Wb$$

Strategy of $T\bar{T}$ searches

Three analysis to probe each channel preferentially

Zt+X

ATLAS-CONF-2017-015

Ht+X

ATLAS-CONF-2016-104

Wb+X

ATLAS-CONF-2016-102


Global Strategy

- ▶ Preselection (detail per analysis in backup):
 - ▶ 1 lepton ($W \rightarrow e\nu$) or/and large E_T^{miss} ($Z \rightarrow \nu\nu$)
 - ▶ Several jets ($W \rightarrow qq$, $H \rightarrow bb$, $t \rightarrow qqb$)
 - ▶ 2 or more b-jets ($H \rightarrow bb$, $t \rightarrow Wb$)
- ▶ Background modeling
 - ▶ $t\bar{t}$ (+jets) estimated using MC and constrained via Control Regions (CR)
 - ▶ Multijet contribution estimated with data driven method
- ▶ Signal Regions (SR)
 - ▶ Optimized to probe the desired channel
 - ▶ Additional cuts on high level variables (jet sub-structure, objects combination ...)
 - ▶ Several signal bins to probe several final states

Analysis specificities


Zt+X (1-lepton)

- ▶ Use m_{T2} , m_T^W to reduce background
- ▶ Leading to a single bin SR
- ▶ $t\bar{t}$ and W+jets CR used in the fit




Ht+X (1-lepton and 0-lepton)

- ▶ Several SR defined:
 - ▶ For instance cut on $N_{mass\ tagged\ jet}$
 - ▶ "mass tagged" jet $\Leftrightarrow m(fat\ jet) > 100\text{GeV}$
 - ▶ For each SR, search for bump in: m_{eff}
- $$m_{eff} = p_T^{lep} + p_T^{jets} + E_T^{miss}$$




Wb+X (1-lepton)

- ▶ Defined a boosted and a resolved SR
- ▶ Search for bump in m_T^{lep} distribution for $T \rightarrow Wb \rightarrow l\nu b$




Post-fit distributions


Zt+X



Ht+X




Wb+X




- ▶ Signal enriched regions (SR) fitted simultaneously with **background enriched regions**
- ▶ No deviation found

Observed Mass limit vs Branching ratios


Zt+X



Ht+X



Wb+X




- ▶ Main systematics are jet modeling and heavy-flavours tagging
- ▶ Each analysis well sensitive to the branching ratio (BR) it is dedicated to probe
- ▶ Excluding up to 1.1 TeV for BR=100%
- ▶ Wb+X (BR=100%) limit can also be reinterpreted in terms of $Y_{-4/3}$ production

Search for same sign leptons production

ATLAS-CONF-2016-032

Probe scenarios with at least two leptons of same sign

- ▶ Very rare in SM production
- ▶ Allow to probe vector like top(T) and vector like bottom ($B \rightarrow Wt, Zb, Hb$)
- ▶ Can also probe $T_{5/3} \rightarrow W^+ t \rightarrow W^+ W^+ b$ pair production
- ▶ Selection
 - ▶ ≥ 2 leptons with at least 2 with same charge
 - ▶ Large E_T^{miss}
 - ▶ At least 2 jets with at least 1 b-tagged jet
- ▶ Dominant background
 - ▶ Charge mis-identification estimated in $Z \rightarrow ee$ resonance
 - ▶ Fake lepton estimated with data driven method




No excess found
 \Rightarrow set limits


	Definition	Name
	$e^\pm e^\pm + e^\pm \mu^\pm + \mu^\pm \mu^\pm + eee + ee\mu + e\mu\mu + \mu\mu\mu, N_{\text{jets}} \geq 2$	
$400 < H_T < 700 \text{ GeV}$	$N_b = 1$	SR0
	$N_b = 2$	SR1
	$N_b \geq 3$	SR2
$H_T \geq 700 \text{ GeV}$	$N_b = 1$	$40 < E_T^{\text{miss}} < 100 \text{ GeV}$ SR3
		$E_T^{\text{miss}} \geq 100 \text{ GeV}$ SR4
	$N_b = 2$	$40 < E_T^{\text{miss}} < 100 \text{ GeV}$ SR5
		$E_T^{\text{miss}} \geq 100 \text{ GeV}$ SR6
	$N_b \geq 3$	$E_T^{\text{miss}} > 40 \text{ GeV}$ SR7

Same sign top events limit setting

Observed limits on VLQs mass versus the decay branching ratio




Vector like bottom (B)



Vector like top (T)

- ▶ Main systematics from the fake lepton estimation
- ▶ Limit on VLQ $T_{5/3}$ pair cross section versus its mass
- ▶ The production of $T_{5/3}\bar{T}_{5/3}$ production is excluded for a mass up to 0.99 TeV



Conclusion

$t\bar{t}$ resonance search

- ▶ No evidence so far of $t\bar{t}$ resonance at the TeV scale
- ▶ Only 10% of available data already analysed
- ▶ With 2016 luminosity may probe higher mass regions and lower cross sections
- ▶ In addition the resolved channel will also be added
- ▶ Keep improving the methods to reconstruct top quarks

VLQ search

- ▶ Several channels studied without evidence of VLQ production
- ▶ Previous searches have been significantly improved with Run-2 data
- ▶ VLQ with mass of $\approx 1\text{ TeV}$ are now excluded in some regions
- ▶ The results will be combined

THANKS

THANK YOU
QUESTIONS ?



Details of ttbar analysis

Selection


Var	Boosted electron selection	Boosted Muon selection
Lepton		
N-lepton	==1	
Pt lepton [GeV]	>30	>25GeV
Quality	LHTight	Medium
Isolation	Track based with pt dependant cone size	
η lepton	>2.47 and not in $1.37 < \eta < 1.52$ (crack)	
Overlapp removal	$\Delta R(el; jet) > 0.4$	$\Delta R(\mu; jet) > 0.04 + 10/pt_\mu$
Leptonic b-jet	Highest pt jet within $\Delta R(el; jet)1.5$	
Hadronic fat jet		
N top tagged fat jet	>1	
Top-Taging	80% efficiency using mass and τ_{32}	
Trimming	R sub = 0.2 and f cut = 0.05	
b-tagging		
Jets	Track jets antikt R=0.2	
WP	MV2c20 algorithm 70% efficiency	

W+jets estimation

$$N_{W^+} + N_{W^-} = \left(\frac{r_{MC}+1}{r_{MC}-1} \right) (D_+ - D_-)$$

With $r_{MC} = \frac{N_{MC}^+}{N_{MC}^-}$ and D_\pm the number of event in data with lepton charge of \pm

Systematics contribution in $t\bar{t}$ res. search



H/A analysis precisions

Generator modification:

The modified generator is MadGraph5_aMC@NLO

Systematics

JES/JER (SM $t\bar{t}$)	6%
JES/JER (Signal S+I)	8%
JES/JER (Signal S)	4%
PDF on signal	12.3%
b-tag (bkg)	2%
b-tag (signal)	1%
$t\bar{t}$ cross section	6.5%
$t\bar{t}$ Parton Showering	5%
multijet bkg norm	20%
ST bkg norm	7.7 %
Z+jet bkg norm	48%
Signal remorm	7.5%
Inerference modeling	0.4%

To find better jet lepton combination

⇒ minimize χ^2

$$\chi^2 = \left[\frac{m_{jj} - m_W}{\sigma_W} \right]^2 + \left[\frac{m_{jjb} - m_{jj} - m_{th} - W}{\sigma_{th-W}} \right]^2 + \left[\frac{m_{j\ell\nu} - m_{t\ell}}{\sigma_{t\ell}} \right]^2 + \left[\frac{(p_{T,jjb} - p_{T,j\ell\nu}) - (p_{T,th} - p_{T,t\ell})}{\sigma_{diffp_T}} \right]^2$$

1. term constraint the W jet mass
2. term constraint the top-W mass
3. term constraint the top mass
4. term constraint the pt baance between leptonic and hadronic side

Multijet background estimation

- ▶ Multijet background is dominant at LHC
- ▶ Analysis design to suppress it a maximum
- ▶ Not enough MC to run representative amount of QCD bkg
- ▶ Fake lepton is not well modeled (else it would have been corrected!)
- ▶ Measure QCD background in data directly

Matrix method

- ▶ define Tight and Loose lepton selection
- ▶ Measure $\epsilon (f)$: the fraction of real(fake) lepton selected as tight lepton
- ▶ $\epsilon (f)$ is measured in a signal (background) enriched region

$$\epsilon = N_{tight}^{real} / N_{loose}^{real}$$

$$f = N_{tight}^{fake} / N_{loose}^{fake}$$

The total number of Tight event is

$$N_{tight} = \epsilon \times N_{prompt} + f \times N_{fakes}$$

$$N_{loose} = N_{prompt} + N_{fakes}$$

By inverting the system it yield N_{prompt} and N_{fakes}

Detailed cut of Zt+X

Signal region definition

Variable	SR	TCR	WCR
E_T^{miss}	$> 350 \text{ GeV}$	$> 300 \text{ GeV}$	
m_T^W	$> 170 \text{ GeV}$	$m_T^W \in [30, 90] \text{ GeV}$	
am_{T2}	$> 175 \text{ GeV}$	$> 100 \text{ GeV}$	
$H_{T,\text{sig}}^{\text{miss}}$	> 12	-	
small- R jet p_T	$> 120, 80, 50, 25 \text{ GeV}$	$> 120, 80, 50, 25 \text{ GeV}$	
number of b -tagged jets	≥ 1	≥ 1 $= 0$	
number of large- R jets	≥ 2	≥ 2	
large- R jet mass	$> 80, 60 \text{ GeV}$	$> 80, 60 \text{ GeV}$	
large- R jet p_T	$> 290, 290 \text{ GeV}$ if $E_T^{\text{miss}} < 450 \text{ GeV}$ $> 200, 200 \text{ GeV}$ if $E_T^{\text{miss}} > 450 \text{ GeV}$	$> 200 \text{ GeV}$	

Systematics:

Jet modeling (scale and resolution)	
Also b, c, light-flavor jet tagging.	(1-5%)
MC $t\bar{t}$ uncert in SR	18% (dominant bkg uncert for $t\bar{t}$)
Interference modelling btw $t\bar{t}$ and ST	11% (dominant for ST modelling)
W+jet modeling	6% on backg modelling
PDF	6%

Detailed cut of Ht+X

Preselection

Preselection requirements		
Requirement	1-lepton channel	0-lepton channel
Trigger	Single-lepton trigger	E_T^{miss} trigger
Leptons	=1 isolated e or μ	=0 isolated e or μ
Jets	≥ 5 jets	≥ 6 jets
b -tagging	≥ 2 b -tagged jets	≥ 2 b -tagged jets
E_T^{miss}	$E_T^{\text{miss}} > 20$ GeV	$E_T^{\text{miss}} > 200$ GeV
Other E_T^{miss} -related	$E_T^{\text{miss}} + m_T^W > 60$ GeV	$\Delta\phi_{\text{min}}^{4j} > 0.4$

0-lep SRs

Search regions (≥ 7 jets)			
Mass-tagged jet multiplicity	b -jet multiplicity	m_T^b (min)	Channel name
0	2	-	0J, ≥ 7 , 2b
0	3	-	0J, ≥ 7 , 3b
0	≥ 4	-	0J, ≥ 7 , $\geq 4b$
1	2	-	1J, ≥ 7 , 2b
1	3	< 160 GeV	1J, ≥ 7 , 3b, LM
1	3	> 160 GeV	1J, ≥ 7 , 3b, HM
1	≥ 4	< 160 GeV	1J, ≥ 7 , $\geq 4b$, LM
1	≥ 4	> 160 GeV	1J, ≥ 7 , $\geq 4b$, HM
≥ 2	2	-	$\geq 2J$, ≥ 7 , 2b
≥ 2	3	< 160 GeV	$\geq 2J$, ≥ 7 , 3b, LM
≥ 2	3	> 160 GeV	$\geq 2J$, ≥ 7 , 3b, HM
≥ 2	≥ 4	-	$\geq 2J$, ≥ 7 , $\geq 4b$
Validation regions (6 jets)			
Mass-tagged jet multiplicity	b -jet multiplicity	m_T^b (min)	Channel name
0	2	-	0J, 6j, 2b
0	3	-	0J, 6j, 3b
0	≥ 4	-	0J, 6j, $\geq 4b$
1	2	-	1J, 6j, 2b
1	3	-	1J, 6j, 3b
1	≥ 4	-	1J, 6j, $\geq 4b$
≥ 2	2	-	$\geq 2J$, 6j, 2b
≥ 2	3	-	$\geq 2J$, 6j, 3b
≥ 2	≥ 4	-	$\geq 2J$, 6j, $\geq 4b$

1-lep SRs

Search regions (≥ 6 jets)				
Mass-tagged jet multiplicity	b -jet multiplicity	$m_{\text{tag}}^{\text{min,AR}}$	m_{tag}	Channel name
0	3	-	> 400 GeV	0J, ≥ 6 , 3b
0	≥ 4	-	> 400 GeV	0J, ≥ 6 , $\geq 4b$
1	3	< 100 GeV	> 700 GeV	1J, ≥ 6 , 3b, LM
1	3	> 100 GeV	> 700 GeV	1J, ≥ 6 , 3b, HM
1	≥ 4	< 100 GeV	> 700 GeV	1J, ≥ 6 , $\geq 4b$, LM
1	≥ 4	> 100 GeV	> 700 GeV	1J, ≥ 6 , $\geq 4b$, HM
≥ 2	3	-	-	$\geq 2J$, ≥ 6 , 3b
≥ 2	≥ 4	-	-	$\geq 2J$, ≥ 6 , $\geq 4b$
Validation regions (5 jets)				
Mass-tagged jet multiplicity	b -jet multiplicity	$m_{\text{tag}}^{\text{min,AR}}$	m_{tag}	Channel name
0	3	-	> 400 GeV	0J, 5j, 3b
0	≥ 4	-	> 400 GeV	0J, 5j, $\geq 4b$
1	3	-	> 700 GeV	1J, 5j, 3b
1	≥ 4	-	> 700 GeV	1J, 5j, $\geq 4b$
≥ 2	3	-	-	$\geq 2J$, 5j, 3b
≥ 2	≥ 4	-	-	$\geq 2J$, 5j, $\geq 4b$

Detailed Systematics in Ht+X

Systematics:

Dominant source depend on SR but,
In most sensitive SR syst dominated by :

tt+jet modeling :	
X-Section $t\bar{t}+1c$	up to 6.1% 50%
jet JES/JER	12/16%
ST modelling :	
X-Section Interference	$\approx 5\%$ 24-52%
V+jet	30%
b, c, light jet modelling	
Diboson	48%

Detailed cut of $Wb+X$

Signal regions definitions

Region	S_T	$\Delta R(\text{lep}, \nu)$	Small- R jets	Large- R jets
Boosted signal region	> 1200 GeV	< 0.8	≥ 3	≥ 1
Resolved signal region	> 1200 GeV	< 0.8	≥ 4	$= 0$
$t\bar{t}$ control region 1	> 700 GeV	> 1.0	≥ 3	≥ 1
$t\bar{t}$ control region 2	$750 - 1200$ GeV	< 1.0	≥ 3	≥ 1

Systematics:

Main source from jet and Large R-jet scale and resolution

lumi.	2.9
jet	24% per jet
VV production	6%
VV+jet production	5%
$t\bar{t}$	6%
single top	6%
Multijet bkg estimation	100%

Detailed cut for SS

Systematics sources from background

Source	Signal region							
	SR0	SR1	SR2	SR3	SR4	SR5	SR6	SR7
Cross section	8	11	26	13	9	27	23	57
Jet energy scale	1	1	3	1	1	3	2	4
Jet energy resolution	<1	2	2	2	<1	1	<1	3
<i>b</i> -tagging efficiency	1	2	5	3	1	2	2	7
Luminosity	1	1	1	1	1	1	1	1
Fake/non-prompt leptons	17	7	15	13	26	13	17	17
Charge misID	8	3	7	5	3	6	5	8


Signal regions definition

	Definition	Name	
	$e^{\pm}e^{\pm} + e^{\pm}\mu^{\pm} + \mu^{\pm}\mu^{\pm} + eee + ee\mu + e\mu\mu + \mu\mu\mu, N_{\text{jets}} \geq 2$		
$400 < H_T < 700 \text{ GeV}$	$N_b = 1$	SR0	
	$N_b = 2$	$E_T^{\text{miss}} > 40 \text{ GeV}$	
	$N_b \geq 3$		
$H_T \geq 700 \text{ GeV}$	$N_b = 1$	$40 < E_T^{\text{miss}} < 100 \text{ GeV}$	SR3
		$E_T^{\text{miss}} \geq 100 \text{ GeV}$	SR4
	$N_b = 2$	$40 < E_T^{\text{miss}} < 100 \text{ GeV}$	SR5
		$E_T^{\text{miss}} \geq 100 \text{ GeV}$	SR6
	$N_b \geq 3$	$E_T^{\text{miss}} > 40 \text{ GeV}$	SR7

Tag and Probe

- ▶ negligible for muons
- ▶ Using the pure $Z \rightarrow ee$ resonance
- ▶ Estimate the charge flip ϵ rate in η, pt bins
- ▶ For one el in bin i and one in bin j the expected number of SS event is

$$N_{SS}^{ij} = N_{ij}(\epsilon_i(1 - \epsilon_j) + \epsilon_j(1 - \epsilon_i))$$



Complex variables mentioned in this talk

τ_{32} is the ratio of τ_3/τ_2 and represent oth probability that a Fat jet is rather 3-prong than 2-prong

τ_N is the N-subjetiness and is computed as

$$\tau_N = \frac{1}{d_0} \sum_k p_{Tk} \times R_k^{min}, \text{ with } d_0 = \sum_k p_{Tk} \times R$$

where k run over all the clusters and R is the characteristic size of sub-kt jets

m_{T2} target dilepton $t\bar{t}$ events. For such topology it has an end-point at $m_{top} = 175\text{GeV}$.
Derived from m_{T2} variables

$$m_T^W \text{ is teh w trasverse mass } m_T^W = \sqrt{2p_T^{lep} E_T^{miss} [1 - \cos(\Delta\phi)]}$$