


Hard Two-Photon Contribution to Elastic Lepton-Proton Scattering Determined by the OLYMPUS Experiment

Dmitry Khaneft for the OLYMPUS collaboration


Johannes Gutenberg University of Mainz
Helmholtz-Institute Mainz

Deep Inelastic Scattering
Birmingham, April 4, 2017




Helmholtz-Institut Mainz

One Value - Two Methods - Two Results



The (Probable) Cause


Unaccounted hard two-photon exchange



How to measure

$$R_{2\gamma} = \frac{\sigma_{e+p}}{\sigma_{e-p}} = \frac{|\mathcal{M}_{1\gamma}|^2 + 2\mathcal{R}(\mathcal{M}_{1\gamma}\mathcal{M}_{2\gamma})}{|\mathcal{M}_{1\gamma}|^2 - 2\mathcal{R}(\mathcal{M}_{1\gamma}\mathcal{M}_{2\gamma})}$$

Situation Until Recently



Modern TPE Experiments

CLAS

- 0.5-3.5 GeV e^\pm beams
- PRL 114, 062003 (2015)


VEPP-3

- 1/1.6 GeV e^\pm beams
- PRL 114, 062005 (2015)

OLYMPUS

- 2 GeV e^\pm beams
- PRL 118, 092501 (2017)


Kinematic Reach of Two-Photon Experiments




OLYMPUS detector

Target chamber

- Hydrogen gas target
- Cooled to ~ 70 K



OLYMPUS detector




Toroid magnet

- Current 5000 A
- Field 0.28 T


OLYMPUS detector

Drift chambers

- $20^\circ < \theta < 80^\circ$
- $-15^\circ < \phi < 15^\circ$




OLYMPUS detector



Time of F light

- Main trigger
- Particle ID

OLYMPUS detector



Luminosity monitors

- Slow control
- 12° detector
- Symmetric Møller/Bhabha monitor

Luminosity Measurements

Slow Control:

- On-line luminosity measurement using running conditions
- Absolute 5%, relative 2%

12° MWPC/GEM monitor:

- Elastic lepton-proton scattering
- Absolute 2.4%, relative 0.46%

Symmetric Møller/Bhabha monitor:

- Multi-interaction lepton-lepton and lepton-proton events (MIE)
- 0.1% statistical, 0.27% systematic

Luminosity Measurements

MIE method was chosen as the most accurate

TPE at 1.29° is negligible

- $\langle Q^2 \rangle = 0.002 \text{ (GeV/c)}^2$
- $\langle \epsilon \rangle = 0.99975$

TPE was also measured at 12° using MPWC/GEM

- $\langle Q^2 \rangle = 0.165 \text{ (GeV/c)}^2$
- $\langle \epsilon \rangle = 0.98$

Timeline

2007

- Letter of Intent

2008

- Proposal approved

2009

- Technical design report

2010

- Funding approved
- BLAST was shipped to DESY

2011

- Tests
- Commissioning

2012

- Run I
- DORIS improvements
- Run II


2013

- Cosmic runs
- Detector survey
- Magnetic field map
- Beam position monitor calibration

2013-2016


- Data analysis

Collected Luminosity



Analysis Chain

- Experimental and simulated data have the same format
- Experimental and simulated data are analyzed with the same code



Analysis Procedure

Four* analyses with the same:

- experimental data samples
- simulated data samples
- Q^2 and ϵ binning
- yield normalization the SYMB-MIE luminosity


and different:

- particle identification
- cuts' size

Results are a combination of the four analyses

*Axel Schmidt, Jan Bernauer, Brian Henderson, Rebecca Russell

Radiative Corrections




Hard Two-Photon Contribution to Elastic Lepton-Proton Scattering Determined by the OLYMPUS Experiment


B. S. Henderson,¹ L. D. Ice,² D. Khanef, ³ C. O'Connor,¹ R. Russell,¹ A. Schmidt,¹ J. C. Bernauer,^{1,*} M. Kohl,^{4,†} N. Akopov,⁵ R. Alarcon,² O. Ates,⁴ A. Avetisyan,⁵ R. Beck,⁶ S. Belostotski,⁷ J. Bessuille,¹ F. Brinker,⁸ J. R. Calarco,⁹ V. Carassiti,¹⁰ E. Cisbani,¹¹ G. Ciullo,¹⁰ M. Contalbrigo,¹⁰ R. De Leo,¹² J. Diefenbach,⁴ T. W. Donnelly,¹ K. Dow,¹ G. Elbakian,⁵ P. D. Eversheim,⁶ S. Frullani,¹¹ Ch. Funke,⁶ G. Gavrilov,⁷ B. Gläser,³ N. Görrissen,⁸ D. K. Hasell,¹ J. Hauschildt,⁸ Ph. Hoffmeister,⁶ Y. Holler,⁸ E. Ihloff,¹ A. Izotov,⁷ R. Kaiser,¹³ G. Karyan,^{8,‡} J. Kelsey,¹ A. Kiselev,⁷ P. Klassen,⁶ A. Krivshich,⁷ I. Lehmann,¹³ P. Lenisa,¹⁰ D. Lenz,⁸ S. Lumsden,¹³ Y. Ma,³ F. Maas,³ H. Marukyan,⁵ O. Miklukho,⁷ R. G. Milner,¹ A. Movsisyan,^{5,§} M. Murray,¹³ Y. Naryshkin,⁷ R. Perez Benito,³ R. Perrino,¹² R. P. Redwine,¹ D. Rodríguez Piñeiro,³ G. Rosner,¹³ U. Schneekloth,⁸ B. Seitz,¹³ M. Statera,¹⁰ A. Thiel,⁶ H. Vardanyan,⁵ D. Veretennikov,⁷ C. Vidal,¹ A. Winnebeck,¹ and V. Yeganov⁵

(OLYMPUS Collaboration)

First OLYMPUS Results



OLYMPUS - CLAS - VEPP-3 Comparison



Conclusion

- All three modern experiments show similar results
- Two-photon effects are small at lower Q^2
- More experimental data are needed at higher energies

