

TMDs at small x


Jian Zhou
ShanDong University

DIS Apr. 2017, Birmingham

Outline:

- Joint TMD and small x evolution
- Spin dependent TMDs at small x
 - 1: unpolarized target
 - 2: transversely polarized target
- Summary

Gluon initiated Drell-Yan process



➤ $M^2 \gg p_T^2$, **TMD factorization**, $\ln \frac{M^2}{p_T^2}$ resummed by Collins-Soper equation

1982-1983, Collins, Soper

➤ $S \gg M^2$, **Kt factorization**, $\ln \frac{S}{M^2}$ resummed by BFKL equation

1991, Catani, Ciafaloni and Hautmann
1991, Collins and R. K. Ellis

The overlap region $S \gg M^2 \gg p_T^2$

An explicit NLO cross section calculation shows that both the large logarithms appear.

2013, Mueller, Xiao, Yuan

Such joint resummation has been also discussed in other literatures.

2015, Balitsky and Tarasov; 2015 Marzani

In collinear calculation $\ln \frac{M^2}{\mu^2}$ absorbed into PDF.

One natural question:


$$\int \frac{d\xi^- d^2\xi_\perp}{(2\pi)^3 P^+} e^{ixP^+\xi^- - ik_\perp \cdot \xi_\perp} \langle P | F^{+i}(\xi^-, \xi_\perp) \mathcal{L}_\xi^\dagger \mathcal{L}_0 F^{+i}(0) | P \rangle$$

Does it accommodate both type large logarithms?

$$\ln \frac{S}{M^2} \quad \ln \frac{M^2}{p_T^2}$$

small x gluon TMD at LO


In a simple quark model, at tree level:



$$xG(x, l_{\perp}, x\zeta)_{LO}|_{x \rightarrow 0} = \frac{\alpha_s C_F}{\pi^2} \frac{1}{l_{\perp}^2}$$

- both large logarithms are absent at LO;
how is it dressed by quantum corrections at NLO?

NLO Sample diagrams:



➤ Calculation formulated in the Ji-Ma-Yuan scheme.

In the leading logarithm approximations,

Small x gluon TMD at NLO reads,

$$\begin{aligned} xG(x, l_{\perp}, x\zeta)_{NLO} &= xG(x, l_{\perp}, x\zeta)_{LO} \\ &+ \frac{\alpha_s N_c}{\pi^2} \ln \frac{1}{x} \int \frac{d^2 k_{\perp}}{k_{\perp}^2} \left[xG_{LO}(x, k_{\perp} + l_{\perp}, x\zeta) - \frac{l_{\perp}^2}{2(l_{\perp} + k_{\perp})^2} xG_{LO}(x, l_{\perp}, x\zeta) \right] \\ &+ \frac{\alpha_s N_c}{2\pi} \left[\ln \frac{x^2 \zeta^2}{l_{\perp}^2} - \frac{1}{2} \ln \frac{x^2 \zeta^2}{\mu^2} - \left(\ln \frac{x^2 \zeta^2}{l_{\perp}^2} \right)^2 \right] xG_{LO}(x, l_{\perp}, x\zeta) \\ &+ \frac{\alpha_s N_c}{\pi^2} \int \frac{d^2 k_{\perp}}{2 [k_{\perp}^2 + l_{\perp}^4 / x^2 \zeta^2]} \ln \frac{k_{\perp}^2 (k_{\perp}^2 + x^2 \zeta^2)}{(k_{\perp}^2 + l_{\perp}^2)^2} xG_{LO}(x, k_{\perp} + l_{\perp}, x\zeta) \end{aligned}$$

2016, ZJ

$$\ln \frac{1}{x} \longrightarrow \ln \frac{S}{M^2}$$

$$\ln \frac{x^2 \zeta^2}{k_T^2} \longrightarrow \ln \frac{M^2}{p_T^2}$$

The resulting gluon TMD indeed simultaneously satisfies the both

BFKL equation:

$$\frac{\partial [xG(x, l_{\perp}, x\zeta)]}{\partial \ln(1/x)} = \frac{\alpha_s N_c}{\pi^2} \int \frac{d^2 k_{\perp}}{k_{\perp}^2} \left\{ xG(x, k_{\perp} + l_{\perp}, x\zeta) - \frac{l_{\perp}^2}{2(l_{\perp} + k_{\perp})^2} xG(x, l_{\perp}, x\zeta) \right\}$$

CS equation:









$$\frac{\partial [G(x, b_{\perp}, x\zeta)]}{\partial \ln \zeta} = -\frac{\alpha_s N_c}{\pi} \ln \left[\frac{x^2 \zeta^2 b_{\perp}^2}{4} e^{2\gamma_E - \frac{1}{2}} \right] G(x, b_{\perp}, x\zeta)$$

More formal treatment

Computing small x gluon TMDs in CGC with Collins 2011 scheme

Small x TMDs in CGC at NLO

Sample diagrams



Collinear approach *vs* CGC I

- TMDs in collinear approach
collinear divergence DGLAP
- TMDs in CGC,
rapidity divergence BK or JIMWLK

Collins-Soper light cone divergence appears in both collinear approach and CGC calculation

Match small x TMDs onto two point functions instead of PDFs.

Collinear approach vs CGC II

$$\tilde{f}_g^{(sub.)}(x, r_\perp, \zeta_c) = e^{-S_{pert}^g(Q, r_\perp)} \sum_i C_{g/i}(\mu_r/\mu) \otimes f_i(x, \mu)$$

Sudakov factor

Hard coefficient

Colliner PDF

$$xG^{(1)}(x, r_\perp, \zeta_c) = -\frac{2}{\alpha_S} \mathcal{H}^{WW}(\alpha_s(Q)) e^{-S_{sud}(Q^2, r_\perp^2)} \mathcal{F}_{Y=\ln 1/x}^{WW}(x_\perp, y_\perp)$$

Hard coefficient

Sudakov factor

Two point function

Xiao, Yuan and ZJ, 2017.

Two step evolution: $S \longrightarrow M^2 \longrightarrow k_T^2$

Spin dependent TMDs at small x


1: Inside an unpolarized target

Linearly polarized gluon TMD

$$\int \frac{dr^- d^2 r_\perp}{(2\pi)^3 P^+} e^{-ix_1 P^+ r^- + i\vec{k}_{1\perp} \cdot \vec{r}_\perp} \langle A | F^{+i}(r^- + y^-, r_\perp + y_\perp) L^\dagger L F^{+j}(y^-, y_\perp) | A \rangle$$

$$= \frac{\delta_\perp^{ij}}{2} x_1 G(x_1, k_{1\perp}) + \left(\hat{k}_{1\perp}^i \hat{k}_{1\perp}^j - \frac{1}{2} \delta_\perp^{ij} \right) x_1 h_1^\perp{}^g(x_1, k_{1\perp}), \quad \text{Mulders, Rodrigues, 2001}$$

Unpolarized gluon TMD
computed in the MV model





Kovchegov, 96
J. Marian, Kovner, McLerran & Weigert, 97

Gluon TMDs in the MV model

The linearly polarized gluon TMDs in the MV model,


Metz & ZJ, 2011

Weizsäcker-Williams(WW) distribution:



$$xh_{1,WW}^{\perp g}(x, k_{\perp}) = \frac{N_c^2 - 1}{8\pi^3} S_{\perp} \int d\xi_{\perp} \frac{K_2(k_{\perp} \xi_{\perp})}{\frac{1}{4\mu_A} \xi_{\perp} Q_s^2} \left(1 - e^{-\frac{\xi_{\perp}^2 Q_s^2}{4}} \right)$$

Dipole distribution:



$$xh_{1,DP}^{\perp g}(x, k_{\perp}) = xG_{DP}^g(x, k_{\perp}) = \frac{k_{\perp}^2 N_c}{2\pi^2 \alpha_s} S_{\perp} \int \frac{d^2 \xi_{\perp}}{(2\pi)^2} e^{-i\vec{k}_{\perp} \cdot \vec{\xi}_{\perp}} e^{-\frac{Q_{sq}^2 \xi_{\perp}^2}{4}}$$

positivity bound saturated for any value of k_t

TMD evolution effect




FIG. 2: The ratio $R = h_1^{\perp g} / f_1^g$ as function of k_{\perp} , at $x = 0.01$ for $\mu = 6, 15$ and 90 GeV.

D. Boer, P. Mulders, JZ, Y. J. Zhou, 2017

2: Inside a transversely polarized target

Three T-odd gluon TMDs

Identify 6 leading power gluon TMDs for a transversely polarized target (8 in total). Among them, 3 gluon TMDs are T-odd distributions.

$$\begin{aligned}
 & \frac{1}{xP^+} \int \frac{dy^- d^2 y_T}{(2\pi)^3} e^{ik \cdot y} \langle P, S_T | 2\text{Tr} [F_{+T}^\mu(0) U F_{+T}^\nu(y) U'] | P, S_T \rangle \\
 &= \delta_T^{\mu\nu} f_1^g + \left(\frac{2k_T^\mu k_T^\nu}{k_\perp^2} - \delta_T^{\mu\nu} \right) h_1^{\perp g} - \delta_T^{\mu\nu} \frac{\epsilon_{T\alpha\beta} k_T^\alpha S_T^\beta}{M} f_{1T}^{\perp g} \\
 & \quad - i\epsilon_T^{\mu\nu} \frac{k_T \cdot S_T}{M} g_{1T}^g - \frac{\tilde{k}_T^{\{\mu} k_T^{\nu\}}}{k_\perp^2} \frac{k_T \cdot S_T}{M} h_{1T}^{\perp g} - \frac{\tilde{k}_T^{\{\mu} S_T^{\nu\}} + \tilde{S}_T^{\{\mu} k_T^{\nu\}}}{2M} h_{1T}^g
 \end{aligned}$$

Mulders, Rodrigues, 2001

Are the T-odd gluon TMDs relevant at small x?

T-odd gluon TMDs & the odderon

Starting point,

$$\Gamma_{T\text{-odd}}^{\mu\nu} = \frac{1}{xP^+} \int \frac{dy^- d^2 y_T}{(2\pi)^3} e^{ik \cdot y} \langle P, S_T | 2 \text{Tr} [F_{+T}^\mu(0) U^{[-]\dagger} F_{+T}^\nu(y) U^{[+]}] | P, S_T \rangle$$

Using time reversal invariance and parity symmetry, at small x one obtains,

$$\Gamma_{T\text{-odd}}^{\mu\nu} = \frac{k_T^\mu k_T^\nu}{g^2 V x P^+} \int \frac{d^2 y_{1T} d^2 y_{2T}}{(2\pi)^3} e^{ik_T \cdot y_T} \langle P, S_T | \text{Tr} [U^{[\square]}(y_T) - U^{[\square]\dagger}(y_T)] | P, S_T \rangle$$

Schematically,

$$\Gamma_{T\text{-odd}}^{\mu\nu} \propto \frac{1}{2} k_T^\mu k_T^\nu \left\{ \begin{array}{c} \overrightarrow{\hspace{10em}} \\ \overleftarrow{\hspace{10em}} \end{array} \right\} - \left\{ \begin{array}{c} \overleftarrow{\hspace{10em}} \\ \overrightarrow{\hspace{10em}} \end{array} \right\}$$

$\hat{D}(R_\perp, r_\perp) = \frac{1}{N_c} \text{Tr} \left[U(R_\perp + \frac{r_\perp}{2}) U^\dagger(R_\perp - \frac{r_\perp}{2}) \right]$

Nothing but an odderon operator in CGC

$$\hat{O}(R_\perp, r_\perp) = \frac{1}{2i} \left[\hat{D}(R_\perp, r_\perp) - \hat{D}(R_\perp, -r_\perp) \right]$$

Kovchegov, Szymanowski & Wallon 2004
Hatta, Iancu, Itakura & McLerran 2005

Spin dependent odderon

$$\begin{aligned}
 \langle \hat{O}(r_{\perp}) \rangle &= -\frac{c_0 \alpha_s^3 \pi}{8M_p R_0^2} e^{-\frac{1}{4} r_{\perp}^2 Q_s^2} r_{\perp}^2 \epsilon_{\perp}^{ij} S_{\perp i} r_{\perp j} \int dx_q d^2 z_{\perp} \sum_{u,d} \mathcal{E}(x_q, z_{\perp}^2) \\
 &= -\frac{c_0 \alpha_s^3 \pi}{8M_p R_0^2} e^{-\frac{1}{4} r_{\perp}^2 Q_s^2} r_{\perp}^2 \epsilon_{\perp}^{ij} S_{\perp i} r_{\perp j} \left(\kappa_p^u + \kappa_p^d \right)
 \end{aligned}$$

ZJ 2013

In the momentum space:

$$\frac{\epsilon_T^{\alpha\beta} S_{T\alpha} k_{T\beta}}{M} O_{1T,x}^{\perp}(k_{\perp}^2) \propto \left\{ \begin{array}{c} \text{--->---} \\ \text{<---} \end{array} \right\} - \left\{ \begin{array}{c} \text{<---} \\ \text{--->---} \end{array} \right\}$$

$$\Gamma_{T\text{-odd}}^{\mu\nu} \propto \frac{1}{2} k_T^{\mu} k_T^{\nu} \left\{ \begin{array}{c} \text{--->---} \\ \text{<---} \end{array} \right\} - \left\{ \begin{array}{c} \text{<---} \\ \text{--->---} \end{array} \right\}$$

Two different parametrizations

Equating two parametrizations:

$$\frac{k_T^\mu k_T^\nu N_c \epsilon_T^{\alpha\beta} S_{T\alpha} k_{T\beta}}{2\pi^2 \alpha_s x M} O_{1T,x}^\perp(k_\perp^2) = -\delta_T^{\mu\nu} \frac{\epsilon_{T\alpha\beta} k_T^\alpha S_T^\beta}{M} f_{1T}^{\perp g}$$

$$- \frac{\tilde{k}_T^{\{\mu} k_T^{\nu\}} k_T \cdot S_T}{k_\perp^2 M} h_{1T}^{\perp g} - \frac{\tilde{k}_T^{\{\mu} S_T^{\nu\}} + \tilde{S}_T^{\{\mu} k_T^{\nu\}}}{2M} h_{1T}^g$$



Simple algebra leads to

$$x f_{1T}^{\perp g} = x h_{1T}^g = x h_{1T}^{\perp g} = \frac{k_\perp^2 N_c}{4\pi^2 \alpha_s} O_{1T,x}^\perp(k_\perp^2)$$

Boer, Echevarria, Mulders, ZJ; 2016

All of three dipole type T-odd gluon TMDs become identical at small x!

Summary

At small x:

- A joint resummation can be achieved
- Very rich polarization dependent phenomenology

**It would be interesting to test these theoretical ideas
at RHIC and the future EIC**

A general remark:

empolying powerful small x machinery to TMD/spin physics often leads to fruitful results.

Thank you for your attention.