

Measurement of Longitudinal Single-Spin Asymmetry for W Boson Production in Polarized Proton+Proton Collisions at STAR

Qinghua Xu (Shandong University)
for the STAR Collaboration




DIS2017, Birmingham
April 3-7, 2017



Flavor separation of nucleon spin


- Sea quark polarization not well constrained from DIS yet:



D. De Florian, R. Sassot, M. Stratmann, W. Vogelsang, PRD80(2009)

Probing sea quark polarization via W production

- Quark polarimetry with W's in p+p collision (example of W^+):



- Spin asymmetry measurements:


$$A_L^{W^+} = \frac{\sigma_+ - \sigma_-}{\sigma_+ + \sigma_-} = \frac{-\Delta u(x_1)\bar{d}(x_2) + \Delta \bar{d}(x_1)u(x_2)}{u(x_1)\bar{d}(x_2) + \bar{d}(x_1)u(x_2)} = \begin{cases} -\frac{\Delta u(x_1)}{u(x_1)}, & y_{W^+} \gg 0 \\ \frac{\Delta \bar{d}(x_1)}{\bar{d}(x_1)}, & y_{W^+} \ll 0 \end{cases}$$

$$A_L^{W^-} = \begin{cases} -\frac{\Delta d(x_1)}{d(x_1)}, & y_{W^-} \gg 0 \\ \frac{\Delta \bar{u}(x_1)}{\bar{u}(x_1)}, & y_{W^-} \ll 0 \end{cases}$$

Probing sea quark polarization via W production

- Ws naturally separate quark flavors
 - > backward/forward region probe sea & valence quarks
- Ws are 100% **parity-violated**
 - > select only one helicity of the coupled (anti)quarks
- Ws are clean theoretically
 - > no fragmentation function involved
- Complementary to SIDIS: high Q^2 , test universality of pdf


$$A_L^W = \frac{\sigma_+ - \sigma_-}{\sigma_+ + \sigma_-}$$





Expectation of $W A_L$ at RHIC

- Large parity-violating asymmetries expected.
- Simplified interpretation at forward and backward rapidity:


$$A_L^{W^-} \propto \frac{-\Delta d(x_1)\bar{u}(x_2) + \Delta\bar{u}(x_1)d(x_2)}{d(x_1)\bar{u}(x_2) + \bar{u}(x_1)d(x_2)}$$



$$A_L^{W^+} \propto \frac{-\Delta u(x_1)\bar{d}(x_2) + \Delta\bar{d}(x_1)u(x_2)}{u(x_1)\bar{d}(x_2) + \bar{d}(x_1)u(x_2)}$$



RHIC- a polarized proton+proton collider




- Polarization direction changes from bunch to bunch
- Spin rotators provide choice of spin orientation

RHIC performance with p+p collisions

- p+p collisions at 500/510 GeV with long. polarization in 2009, 2011, 2012 and 2013.
- STAR data sample for W_{A_L} analysis:

STAR Longitudinal pp 500/510			
Run	L (pb ⁻¹)	P	P ² L (pb ⁻¹)
2009	12	38%	1.7
2011	9.4	49%	2.3
2012	77	56%	24
2013	246.2	56%	77.2



STAR - Solenoid Tracker At RHIC

Magnet

- 0.5 T Solenoid

Triggering & Luminosity Monitor


- Beam-Beam Counters
 - $3.4 < |\eta| < 5.0$
- Zero Degree Calorimeters
- Vertex Position Detector

Central Tracking

- Large-volume TPC
 - $|\eta| < 1.3$

Calorimetry

- Barrel EMC (Pb/Scintillator)
 - $|\eta| < 1.0$
- Endcap EMC (Pb/Scintillator)
 - $1.0 < \eta < 2.0$
- Forward Meson Spectrometer
 - $2.5 < \eta < 4.0$




(- those marked red are relevant to W analysis)

W selection via $W \rightarrow e\nu$ at STAR


$W \rightarrow e + \nu$ Candidate Event:

- Isolated track pointing to isolated EM cluster in calorimeter
- Large “missing energy” opposite the electron candidate




QCD Background Event

- Several tracks pointing to energy deposit in several towers
- p_T sum is balanced by di-jet, no large “missing energy”



W selection at STAR : Jacobian peak



- Isolation ratio $E_{2 \times 2} / E_{4 \times 4} > 95\%$




- Isolation ratio $E_T^e / E_T^{\Delta R < 0.7} > 88\%$

$$\vec{p}_T^{bal} = \vec{p}_T^e + \sum_{\Delta R > 0.7} \vec{p}_T^{jets}$$


- Signed P_T -balance = $\frac{\vec{p}_T^e \cdot \vec{p}_T^{bal}}{|\vec{p}_T^e|} > 14 \text{ GeV}$
- away $E_T < 11 \text{ GeV}$

Signal of Jacobian peak with E_T dis. after selection :

-STAR 2013 with BEMC ($|\eta| < 1$)



W selection ($|\eta| < 1$) : BG Estimation



- Primary Background

Data-driven QCD : BG Events which satisfy $e^+/-$ candidate isolation cuts

Second EEMC : due to “jet” escape without East EEMC based on real West EEMC


- Weak decay Background

From $Z \rightarrow ee$, and $W \rightarrow \tau\nu$, determined from MC



-STAR 2013

W selection at forward region with EEMC

Signal of Jacobian peak after similar selections at $1 < \eta < 2$:




Background estimation at $1 < \eta < 2$:



STAR mid-rapidity $W A_L$ –2011+2012

- First multiple-eta-bin A_L results from 2011+2012 data:




- A_L of W^- shows indication that data are larger than the DSSV predictions
- A_L of W^+ is consistent with theoretical predictions with DSSV pdf.
- Indication of symmetry breaking of polarized sea.

STAR, PRL113(2014)72301

Global Analysis with STAR W A_L 2012


STAR, PRL113(2014)72301




STAR 2012 W results provide significant constraints on $\Delta\bar{u}$, $\Delta\bar{d}$.

arXiv:1304.0079

$\Delta\bar{u}$



$\Delta\bar{d}$






Global Analysis with STAR W A_L results


- Big impact seen in NNPDFpol1.1 global analysis after including STAR A_L data.

NNPDF1.1, Nucl.Phys. B887,276 (2014)

- Polarized sea asymmetry:




W A_L results – STAR 2013




- STAR 2013 W A_L results:
 - Jinlong Zhang @ INPC 2016
 - Devika Gunarathne @ SPIN2016

W A_L results – STAR 2013




- STAR 2013 W A_L results:
 - Jinlong Zhang @ INPC 2016
 - Devika Gunarathne @ SPIN2016
- Impact in reweighting NNPDFpol1.1




E. Nocera, arXiv:1702.05077

W A_L results – STAR 2013




- New preliminary A_L results at near-forward rapidity !
- STAR 2013 results are the most precise measurements of W A_L so far.

W A_L results – STAR 2013



- New preliminary A_L results at near-forward rapidity !
- STAR 2013 results are the most precise measurements of W A_L so far.
- Consistent with 2011+2012 published results, with 40% uncertainty reduced.
- Further confirmed the preference of positive $\Delta\bar{u}$.
- Can provide further constraints for $\Delta\bar{u}$, $\Delta\bar{d}$.


W A_L results – STAR 2013



- STAR, PRL113,072301(2014)
 - PHENIX, PRD93,051103(2016)

- New preliminary A_L results at near-forward rapidity !
- STAR 2013 results are the most precise measurements of $W A_L$ so far.
- Consistent with 2011+2012 published results, with 40% uncertainty reduced.
- Further confirmed the preference of positive $\Delta\bar{u}$.
- Can provide further constraints for $\Delta\bar{u}$, $\Delta\bar{d}$.

Impact of 2013 W A_L results



NNPDFpol1.1 reweighting


$\Delta\bar{u}$

$$\langle \Delta\bar{u} \rangle^{[0.01,0.2]} = 0.01 \pm 0.04$$

$$\langle \Delta\bar{u} \rangle_{rw}^{[0.01,0.2]} = 0.02 \pm 0.03$$

$Q^2=10\text{GeV}^2$

NNPDFpol1.1
 NPDPol1.1 (rw)




$\Delta\bar{d}$

$$\langle \Delta\bar{d} \rangle^{[0.01,0.2]} = -0.04 \pm 0.04$$

$$\langle \Delta\bar{d} \rangle_{rw}^{[0.01,0.2]} = -0.05 \pm 0.03$$

$(Q^2=10\text{GeV}^2)$



-Provided further constraints for $\Delta\bar{u}$, $\Delta\bar{d}$!

E. Nocera, arXiv:1702.05077

Summary

- ❑ Sea quark polarization plays an important role in fully understanding the nucleon spin structure.
- ❑ Unique clean probe of sea quark polarization via W production at RHIC:
 - RHIC $W A_L$ results provided important constraints on $\Delta\bar{u}, \Delta\bar{d}$.
First clear evidence of flavor asymmetry for polarized sea.
- ❑ New $W A_L$ results from STAR 2013 data set:
 - ✓ Near forward rapidity measurement just released.
 - ✓ The most precise measurements of $W A_L$.
 - ✓ Provide further constraints on sea quark helicity distributions.

Thanks!
