

Searches for Sterile Neutrinos at Future electron-proton Colliders


Oliver Fischer

University of Basel, Switzerland

April the 5th 2017, Birmingham

Based on S. Antusch, E. Cazzato, OF, 1612.02728

Motivation for sterile neutrinos



- ▶ Neutrino oscillations are evidence for new physics.
⇒ *At least* two light neutrinos are massive.
- ▶ Sterile neutrinos for type I seesaw mechanism.

The Seesaw Mechanism

- ▶ Naïve (1 ν_L , 1 ν_R) version: $m_\nu = \frac{1}{2} \frac{v_{\text{EW}}^2 |y_\nu|^2}{M_R}$
- ▶ More realistic example, the (2 ν_L , 2 ν_R) version:

$$Y_\nu = \begin{pmatrix} \mathcal{O}(y_\nu) & 0 \\ 0 & \mathcal{O}(y_\nu) \end{pmatrix}, \quad M_N = \begin{pmatrix} M_R & 0 \\ 0 & M_R(1 + \varepsilon) \end{pmatrix}$$

$$\Rightarrow m_{\nu_i} = \frac{v_{\text{EW}}^2 \mathcal{O}(y_\nu^2)}{M_R} (1 + \varepsilon)$$

\Rightarrow Knowledge of m_{ν_i} implies a relation between y_ν and M_R .

Lowscale Seesaw


- ▶ Specific structures of the Yukawa and mass matrices can be realised by symmetries (no fine tuning).
- ▶ A $(2 \nu_L, 2 \nu_R)$ example:

$$Y_\nu = \begin{pmatrix} \mathcal{O}(y_\nu) & 0 \\ \mathcal{O}(y_\nu) & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & M_R \\ M_R & \epsilon \end{pmatrix}$$

$$\Rightarrow m_{\nu_i} = 0 + \epsilon \frac{v_{EW}^2 \mathcal{O}(y_\nu^2)}{M_R^2}$$

- \Rightarrow In general: no fixed relation between y_ν and M_R .
- \Rightarrow Large y_ν are compatible with neutrino oscillations.

The Big Picture





Sterile neutrinos: attractive BSM features

Baryon asymmetry:

- ▶ Oscillations between sterile and active neutrinos
- ▶ Novel *Higgs-decay* mechanism

Hambye, Teresi; PRL 117 (2016)



Dark Matter:

- ▶ DM stability from lepton symmetry
- ▶ Freeze-in mechanism
- ▶ Testable LFV signatures

Heurtier, Teresi; Phys. Rev. D 94 (2016) no.12, 125022

Present indirect constraints from precision data


- ▶ Analysis of non-unitarity of the PMNS matrix.
- ▶ 34 precision observables:
Electroweak Precision Observables (EWPO), lepton universality, charged lepton flavour violation, CKM unitarity
- ▶ Highest posterior density intervals at 90% Bayesian C.L.:

-0.0021	$\leq \varepsilon_{ee} \leq$	-0.0002	$ \varepsilon_{e\mu} $	$<$	1.0×10^{-5}
-0.0004	$\leq \varepsilon_{\mu\mu} \leq$	0	$ \varepsilon_{e\tau} $	$<$	2.1×10^{-3}
-0.0053	$\leq \varepsilon_{\tau\tau} \leq$	0	$ \varepsilon_{\mu\tau} $	$<$	8.0×10^{-4}

Antusch, OF; JHEP 1410 (2014) 094

- ★ Non-unitarity parameters: $\varepsilon_{\alpha\alpha} = -\theta_{\alpha}^* \theta_{\alpha}$.
- ★ Weak statistical preference for non-zero mixing for ε_{ee} .

Present constraints including direct searches



- ▶ Z pole search: limits from Z branching ratios .


Abreu *et al.* Z.Phys. C74 (1997) 57-71

- ▶ Higgs decays: Best constraints from $h \rightarrow \gamma\gamma$.

- ▶ Direct Search: $\delta\sigma_{SM}^{WW} = 0.011_{stat} + 0.007_{syst}$


OPAL collaboration, Abbiendi *et al.* (2007)

Heavy neutrino production at electron-proton colliders




- ▶ Leading order production of heavy neutrino mass eigenstate.
- ▶ $W_t^{(q)}$: dominant at lower center-of-mass energies.
- ▶ $W_t^{(\gamma)}$: relevant for larger masses.

Production cross sections



LHeC



FCC-eh

For 60 GeV as benchmark for the electron beam E_e :

- ▶ $\sigma_{\nu N}$ increase of $\sim 30\%$ for $E_e \rightarrow 100$ GeV.
- ▶ Increased by $\sim 80\%$ when including polarisation.
- ▶ Consider 1 ab^{-1} (for FCC-eh and LHeC).

Signal channels from $\mathbf{W}_t^{(q)}$

Name	Final State	$ \theta_\alpha $ Dependency	LFV
lepton-trijet	$jjj\ell_\alpha^-$	$\frac{ \theta_e\theta_\alpha ^2}{\theta^2}$	✓
jet-dilepton	$j\ell_\alpha^-\ell_\beta^+\nu$	$\frac{ \theta_e\theta_\alpha ^2}{\theta^2}^{(*)}$	✓
trijet	$jjj\nu$	$ \theta_e ^2$	×
monojet	$j\nu\nu\nu$	$ \theta_e ^2$	×

- ▶ LFV (and LNV) signature for $\alpha \neq e$, $\beta \neq \alpha$, and $\gamma \neq \alpha, \beta$
- ▶ Unambiguous lepton-number-violating final states, e.g. e^+jjj .

Signal channels from $W_t^{(\gamma)}$


Name	Final State	$ \theta_\alpha $ Dependency	LFV
lepton-quadrivet	$jjjj\ell_\alpha^-$	$\frac{ \theta_e\theta_\alpha ^2}{\theta^2}$	✓
dilepton-dijet	$\ell_\alpha^- \ell_\beta^+ \nu jj$	$\frac{ \theta_e\theta_\alpha ^2}{\theta^2}^{(*)}$	✓
trilepton	$\ell_\alpha^- \ell_\beta^- \ell_\gamma^+ \nu \nu$	$\frac{ \theta_e\theta_\alpha ^2}{\theta^2}^{(*)}$	✓
quadrivet	$jjjj\nu$	$ \theta_e ^2$	×
electron-di-b-jet	$e^- bb\nu\nu$	$ \theta_e ^2$	×
dijet	$jj\nu\nu\nu$	$ \theta_e ^2$	×
monolepton	$\ell_\alpha^- \nu\nu\nu\nu$	$ \theta_e ^2$	×

- ▶ Additional signatures of LFV/LNV.
- ▶ Cross section suppressed by the small PDF of the photon.
- ▶ More efficient at higher center-of-mass energies.

Caveat:

- ★ For the following “first look” we confined ourselves to the parton level.
- ★ The analysis on the reconstructed level is about to begin.

First look: lepton-flavour-conserving signatures



— $e^-jjj: |\Theta|^2=|\theta_e|^4/|\theta|^2$

— $\nu jjj: |\Theta|^2=|\theta_e|^2$

— $j\nu\nu\nu: |\Theta|^2=|\theta_e|^2$


— $e^-\nu\nu bb: |\Theta|^2=|\theta_e|^2$

Dashed lines denote the LHeC

[Antusch, Cazzato, OF; \[1612.02728\]](#)


- ▶ Sensitive to $|\theta_e|$ or $|\theta_e| \times \text{Br}(N \rightarrow e^- W^+)$.
- ▶ Here: “Conservative” sensitivity for $|\theta_\alpha| \ll |\theta_e|$.
- ▶ For $|\theta_\alpha| \sim |\theta_e|$ LFV is *expected*.

Lepton-flavour-violating signatures



- ▶ Very sensitive tests of combinations $|\theta_e\theta_\alpha|$.
- ▶ Upper bounds:
 $|\theta_e\theta_\mu|$ from $\mu \rightarrow e\gamma$ (MEG); $|\theta_e\theta_\tau|$ from precision data.
- ▶ Requires $|\theta_\alpha| \gtrsim |\theta_e|$ for sizeable branching ratios.

Synergy and Complementarity with other colliders



Antusch, Cazzato, OF; [1612.02728]

Conclusions

- ▶ Sterile neutrinos are well motivated extensions of the SM.
 - ▶ Symmetry protected seesaw scenarios allow for electroweak scale sterile neutrino masses and $\mathcal{O}(1)$ active-sterile mixings.
 - ▶ Present constraints: active-sterile mixing $|\theta|^2 \leq 10^{-3}$.
 - ▶ Electron-proton colliders produce heavy neutrinos via $|\theta_e|$.
 - ▶ Most sensitive searches for sterile neutrinos with masses
 - below m_W via displaced vertices (not discussed here)
 - above $\mathcal{O}(100)$ GeV via lepton-flavour violating signatures.
 - ▶ Lepton-number violating signatures not expected.
- ⇒ Direct searches for sterile neutrinos at ep colliders have great prospects at testing sterile neutrinos with masses up to 3 TeV!

Thank you for your attention.

Backup I - EWPO

Experimental results and SM predictions for the EWPO, and the modification*, to first order in the “non-unitarity” parameters

$$\varepsilon_{\alpha\alpha} = \theta_{\alpha}^* \theta_{\beta}. \quad (\text{formulae for } M \gg m_Z)$$

Prediction in MUV	SM Prediction	Experiment
$[R_{\ell}]_{\text{SM}} (1 - 0.15(\varepsilon_{ee} + \varepsilon_{\mu\mu}))$	20.744(11)	20.767(25)
$[R_b]_{\text{SM}} (1 + 0.03(\varepsilon_{ee} + \varepsilon_{\mu\mu}))$	0.21577(4)	0.21629(66)
$[R_c]_{\text{SM}} (1 - 0.06(\varepsilon_{ee} + \varepsilon_{\mu\mu}))$	0.17226(6)	0.1721(30)
$[\sigma_{had}^0]_{\text{SM}} (1 - 0.25(\varepsilon_{ee} + \varepsilon_{\mu\mu}) - 0.27\varepsilon_{\tau})/\text{nb}$	41.470(15)	41.541(37)
$[R_{inv}]_{\text{SM}} (1 + 0.75(\varepsilon_{ee} + \varepsilon_{\mu\mu}) + 0.67\varepsilon_{\tau})$	5.9723(10)	5.942(16)
$[M_W]_{\text{SM}} (1 - 0.11(\varepsilon_{ee} + \varepsilon_{\mu\mu}))/\text{GeV}$	80.359(11)	80.385(15)
$[\Gamma_{\text{lept}}]_{\text{SM}} (1 - 0.59(\varepsilon_{ee} + \varepsilon_{\mu\mu}))/\text{MeV}$	83.966(12)	83.984(86)
$[(s_{W,\text{eff}}^{\ell,\text{lep}})^2]_{\text{SM}} (1 + 0.71(\varepsilon_{ee} + \varepsilon_{\mu\mu}))$	0.23150(1)	0.23113(21)
$[(s_{W,\text{eff}}^{\ell,\text{had}})^2]_{\text{SM}} (1 + 0.71(\varepsilon_{ee} + \varepsilon_{\mu\mu}))$	0.23150(1)	0.23222(27)

* Minimal Unitarity Violation scheme: [Antusch et al.; JHEP 0610 \(2006\) 084.](#)

Backup II - lepton universality

Modification due to sterile neutrinos (formulae for $M \gg m_Z$):

$$R_{\alpha\beta} = \sqrt{\frac{(NN^\dagger)_{\alpha\alpha}}{(NN^\dagger)_{\beta\beta}}} \simeq 1 + \frac{1}{2} (\varepsilon_{\alpha\alpha} - \varepsilon_{\beta\beta}) .$$

	Process	Bound		Process	Bound
$R_{\mu e}^\ell$	$\frac{\Gamma(\tau \rightarrow \nu_\tau \mu \bar{\nu}_\mu)}{\Gamma(\tau \rightarrow \nu_\tau e \bar{\nu}_e)}$	1.0018(14)	$R_{\mu e}^\pi$	$\frac{\Gamma(\pi \rightarrow \mu \bar{\nu}_\mu)}{\Gamma(\pi \rightarrow e \bar{\nu}_e)}$	1.0021(16)
$R_{\tau\mu}^\ell$	$\frac{\Gamma(\tau \rightarrow \nu_\tau e \bar{\nu}_e)}{\Gamma(\mu \rightarrow \nu_\mu e \bar{\nu}_e)}$	1.0006(21)	$R_{\tau\mu}^\pi$	$\frac{\Gamma(\tau \rightarrow \nu_\tau \pi)}{\Gamma(\pi \rightarrow \mu \bar{\nu}_\mu)}$	0.9956(31)
$R_{e\mu}^W$	$\frac{\Gamma(W \rightarrow e \bar{\nu}_e)}{\Gamma(W \rightarrow \mu \bar{\nu}_\mu)}$	1.0085(93)	$R_{\tau\mu}^K$	$\frac{\Gamma(\tau \rightarrow K \nu_\tau)}{\Gamma(K \rightarrow \mu \bar{\nu}_\mu)}$	0.9852(72)
$R_{\tau\mu}^W$	$\frac{\Gamma(W \rightarrow \tau \bar{\nu}_\tau)}{\Gamma(W \rightarrow \mu \bar{\nu}_e)}$	1.032(11)	$R_{\tau e}^K$	$\frac{\Gamma(\tau \rightarrow K \nu_\tau)}{\Gamma(K \rightarrow e \bar{\nu}_e)}$	1.018(42)

Backup III - CKM unitarity constraint

Current world averages: $V_{ud} = 0.97427(15)$, $V_{ub} = 0.00351(15)$

$$|V_{ij}^{th}|^2 = |V_{ij}^{exp}|^2(1 + f^{\text{process}}(\varepsilon_{\alpha\alpha})) ,$$

$$|V_{ud}^{th}|^2 = |V_{ud}^{exp,\beta}|^2(NN^\dagger)_{\mu\mu} .$$

For the kaon decay processes we have:

$$|V_{us}^{th}|^2 = |V_{us}^{exp,K \rightarrow e}|^2(NN^\dagger)_{\mu\mu} ,$$

$$|V_{us}^{th}|^2 = |V_{us}^{exp,K \rightarrow \mu}|^2(NN^\dagger)_{ee} .$$

Process	$V_{us}f_+(0)$
$K_L \rightarrow \pi e \nu$	0.2163(6)
$K_L \rightarrow \pi \mu \nu$	0.2166(6)
$K_S \rightarrow \pi e \nu$	0.2155(13)
$K^\pm \rightarrow \pi e \nu$	0.2160(11)
$K^\pm \rightarrow \pi \mu \nu$	0.2158(14)
Average	0.2163(5)

Processes involving tau leptons:

Process	$f^{\text{process}}(\varepsilon)$	$ V_{us} $
$\frac{B(\tau \rightarrow K \nu)}{B(\tau \rightarrow \pi \nu)}$	$\varepsilon_{\mu\mu}$	0.2262(13)
$\tau \rightarrow K \nu$	$\varepsilon_{ee} + \varepsilon_{\mu\mu} - \varepsilon_{\tau\tau}$	0.2214(22)
$\tau \rightarrow \ell, \tau \rightarrow s$	$0.2\varepsilon_{ee} - 0.9\varepsilon_{\mu\mu} - 0.2\varepsilon_{\tau\tau}$	0.2173(22)

Backup IV - lepton flavour violation

- Present experimental limits at 90% C.L.:

Process	MUV Prediction	Bound	Constraint on $ \varepsilon_{\alpha\beta} $
$\mu \rightarrow e\gamma$	$2.4 \times 10^{-3} \varepsilon_{\mu e} ^2$	5.7×10^{-13}	$\varepsilon_{\mu e} < 1.5 \times 10^{-5}$
$\tau \rightarrow e\gamma$	$4.3 \times 10^{-4} \varepsilon_{\tau e} ^2$	1.5×10^{-8}	$\varepsilon_{\tau e} < 5.9 \times 10^{-3}$
$\tau \rightarrow \mu\gamma$	$4.1 \times 10^{-4} \varepsilon_{\tau\mu} ^2$	1.8×10^{-8}	$\varepsilon_{\tau\mu} < 6.6 \times 10^{-3}$

- Estimated sensitivities of planned experiments at 90% C.L.:

Process	MUV Prediction	Bound	Sensitivity
$Br_{\tau e}$	$4.3 \times 10^{-4} \varepsilon_{\tau e} ^2$	10^{-9}	$\varepsilon_{\tau e} \geq 1.5 \times 10^{-3}$
$Br_{\tau\mu}$	$4.1 \times 10^{-4} \varepsilon_{\tau\mu} ^2$	10^{-9}	$\varepsilon_{\tau\mu} \geq 1.6 \times 10^{-3}$
$Br_{\mu eee}$	$1.8 \times 10^{-5} \varepsilon_{\mu e} ^2$	10^{-16}	$\varepsilon_{\mu e} \geq 2.4 \times 10^{-6}$
$R_{\mu e}^{Ti}$	$1.5 \times 10^{-5} \varepsilon_{\mu e} ^2$	2×10^{-18}	$\varepsilon_{\mu e} \geq 3.6 \times 10^{-7}$

$\Rightarrow R_{\mu e}^{Ti}$ yields a sensitivity to m_{ν_R} up to 0.3 PeV.

Backup V - Symmetry Protected Seesaw Scenario

Benchmark model, defined in Antusch, OF; JHEP **1505** (2015) 053

- ▶ Collider phenomenology dominated by two sterile neutrinos N_i with protective symmetry, such that

$$\mathcal{L}_N = -\frac{1}{2}\overline{N_R^1}M(N_R^2)^c - y_{\nu\alpha}\overline{N_R^1}\tilde{\phi}^\dagger L^\alpha + \text{H.c.}$$

- ▶ Further “decoupled” sterile neutrinos may exist.
- ▶ Active-sterile mixing parameters:

$$\theta_\alpha = y_{\nu\alpha}\frac{v_{\text{EW}}}{\sqrt{2}}, \theta^2 \equiv \sum_\alpha |\theta_\alpha|^2$$

- ▶ The leptonic mixing matrix to leading order in θ_α :

$$\mathcal{U} = \begin{pmatrix} \mathcal{N}_{e1} & \mathcal{N}_{e2} & \mathcal{N}_{e3} & -\frac{i}{\sqrt{2}}\theta_e & \frac{1}{\sqrt{2}}\theta_e \\ \mathcal{N}_{\mu 1} & \mathcal{N}_{\mu 2} & \mathcal{N}_{\mu 3} & -\frac{i}{\sqrt{2}}\theta_\mu & \frac{1}{\sqrt{2}}\theta_\mu \\ \mathcal{N}_{\tau 1} & \mathcal{N}_{\tau 2} & \mathcal{N}_{\tau 3} & -\frac{i}{\sqrt{2}}\theta_\tau & \frac{1}{\sqrt{2}}\theta_\tau \\ 0 & 0 & 0 & \frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\theta_e^* & -\theta_\mu^* & -\theta_\tau^* & -\frac{i}{\sqrt{2}}\left(1 - \frac{\theta^2}{2}\right) & \frac{1}{\sqrt{2}}\left(1 - \frac{\theta^2}{2}\right) \end{pmatrix}$$

Backup VI - Heavy neutrino interactions

- ▶ **Charged current (CC):**

$$j_{\mu}^{\pm} = \frac{g}{2} \theta_{\alpha} \bar{\ell}_{\alpha} \gamma_{\mu} (-iN_1 + N_2)$$

- ▶ **Neutral current (NC):**


$$j_{\mu}^0 = \frac{g}{2 c_W} [\theta^2 \bar{N}_2 \gamma_{\mu} N_2 + (\bar{\nu}_i \gamma_{\mu} \xi_{\alpha 1} N_1 + \bar{\nu}_i \gamma_{\mu} \xi_{\alpha 2} N_2 + \text{H.c.})]$$

- ▶ Higgs boson **Yukawa** interaction:

$$\mathcal{L}_{\text{Yukawa}} = \sum_{i=1}^3 \xi_{\alpha 2} \frac{\sqrt{2} M}{v_{\text{EW}}} \nu_i \phi^0 (\bar{N}_1 + \bar{N}_2)$$

- ▶ With the mixing parameters: $\xi_{\alpha 1} = (-i) \mathcal{N}_{\alpha\beta}^* \frac{\theta_{\beta}}{\sqrt{2}}$, $\xi_{\alpha 2} = i \xi_{\alpha 1}$

ILC direct searches



Antusch, Cazzato, OF; *in preparation*

- ▶ Operation scenario G-20, with 4 ab^{-1} at $\sqrt{s} = 500 \text{ GeV}$.
- ▶ Using 1 ab^{-1} at $\sqrt{s} = 1.0 \text{ TeV}$.
- ▶ Displaced vertex searches possible for $M < m_W$.