## **Tutorial** 1

24 January 2017, Archamps

hector.garcia.morales@cern.ch, andrea.latina@cern.ch, dario.pellegrini@cern.ch, guido.sterbini@cern.ch

## TUTORIAL 1: FIRST PART

My first MADX job.

Check that your console is working as expected with the following example.

- 1. Open an editor and write your first MADX input file (just 1 line like "stop;" or "exit;" or "quit;").
- 2. Run it. If all is fine, nothing interesting should happen.

## TUTORIAL 1: SECOND PART

## My first accelerator.

- 1. Make a simple FODO cell of  $L_{cell} = 100$  m. Each quad is  $L_{quad} = 5$  m long. Put the start of the first quadrupole at the start of the sequence. Each quad has a focal length of f = 200 m ( $K1 \times L_{quad} = 1/f$  in thin lens approximation).
- 2. Define a proton beam at  $E_{tot} = 2$  GeV. Activate the sequence, try to find the periodic solution and plot the  $\beta$ -functions. If you found  $\beta_{max} \approx 460$  m you succeeded.
- 3. Using the plot you obtained can you estimate the phase advance of the cell? Compare with the tunes obtained from the TWISS.
- 4. Try with  $E_{tot} = 0.7$  GeV: what is the MADX error message?
- 5. Try with f = 20 m: what is the MADX error message?