

Nucleon TMDs from Lattice QCD

Boram Yoon ^{a*}, Tanmoy Bhattacharya^a, Michael Engelhardt^b, Jeremy Green^c,
Rajan Gupta^a, Philipp Hägler^d, Bernhard Musch^d, John Negele^e, Andrew Pochinsky^e,
Sergey Syritsyn^f

^aLos Alamos National Laboratory, MS B283, P.O. Box 1663, Los Alamos, NM 87545, USA

^bDepartment of Physics, New Mexico State University, Las Cruces, NM 88003-8001, USA

^cInstitut für Kernphysik, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany

^dInstitut für Theoretische Physik, Universität Regensburg, 93040 Regensburg, Germany

^eCenter for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

^fTheory Center, Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA

Abstract

We present a lattice QCD calculation of transverse momentum dependent parton distribution functions (TMDs) of protons using staple-shaped Wilson lines. For time-reversal odd observables, we calculate the generalized Sivers and Boer-Mulders transverse momentum shifts in SIDIS and DY cases, and for T-even observables we calculate the transversity related to the tensor charge and the generalized worm-gear shift. The calculation is done on two different $n_f = 2 + 1$ ensembles: domain-wall fermion (DWF) with lattice spacing 0.084 fm and pion mass of 297 MeV, and clover fermion with lattice spacing 0.114 fm and pion mass of 317 MeV. The results from those two different discretizations are consistent with each other.

*Speaker. E-mail: boram@lanl.gov