



Contribution ID: 115

Type: oral presentation

## **Skewness of Event-by-event Elliptic Flow** Fluctuations in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with CMS

Thursday 13 July 2017 09:40 (20 minutes)

Event-by-event elliptic flow harmonic distributions  $p(v_2)$  are measured in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV using the CMS detector for the integrated  $p_T$  range  $0.3 < p_T < 3.0$  GeV/c and pseudorapidity range |eta| < 1.0. In order to gain insight on the nature of the initial geometry fluctuations, cumulant flow harmonics are calculated from the moments of the  $v_2$  probability distribution  $p(v_2)$ . A fine-level splitting between the higher-order cumulants is observed. The skewness with respect to the reaction plane is estimated from the cumulants and found to be negative, which suggests a non-Gaussian nature for the initial-state fluctuations, as predicted by hydrodynamic models. These observations suggest a non-Gaussian nature of the initial-state fluctuations. Furthermore, assuming that the flow harmonics are linearly proportional to the initial-state eccentricities, the  $p(v_2)$  distributions are fitted using an elliptic power law parameterization to study the initial-state geometry fluctuations.

## List of tracks

Hydrodynamics

Primary author: THE CMS COLLABORATION

Presenter: NAZAROVA, Elizaveta (M.V. Lomonosov Moscow State University (RU)) Session Classification: Parallel Hydrodynamics 1

Track Classification: Hydrodynamics