

Blois Workshop 2017
Prague, June 26th-30th 2017

$\Psi(2S)/J/\psi$ ratio at HERA


Marta Ruspa

Univ. Piemonte Orientale & INFN-Torino, Italy





Inclusive and exclusive diffraction


Inclusive



Exclusive




$$e p \rightarrow e' X p$$



$$e p \rightarrow e' J/\psi p \quad J/\psi \rightarrow \mu^+ \mu^-$$

Inclusive and exclusive diffraction

Inclusive




Q^2 = virtuality of photon =
= (4-momentum exchanged at e vertex)²

W = invariant mass of γ^* -p system

t = (4-momentum exchanged at p vertex)²
typically: $|t| < 1 \text{ GeV}^2$

- Single diffraction/elastic: N=proton
- Double diffraction: proton-dissociative system N

Exclusive



M_X = invariant mass of γ^* -IP system

x_{IP} = fraction of proton's momentum
carried by IP


β = Bjorken's variable for the IP
= fraction of IP momentum
carried by struck quark
= x/x_{IP}

- Scattered electron

- VM decay products and nothing
else in the central detector

- Proton undetected

Motivation



- $\Psi(2S)$ wave function different from J/ψ wave function
- Ratio $R = \frac{\sigma_{\gamma p \rightarrow \Psi(2S)p}}{\sigma_{\gamma p \rightarrow J/\psi p}}$ sensitive to radial wave function of charmonium

pQCD models predict $R \approx 0.17$ (photoproduction) and rise of R with Q^2

Samples

$$\begin{aligned}\Psi(2S) &\rightarrow J/\psi \pi^+ \pi^- & J/\psi &\rightarrow \mu^+ \mu^- \\ \Psi(2S) &\rightarrow \mu^+ \mu^- \\ J/\psi &\rightarrow \mu^+ \mu^-\end{aligned}$$



- Data sample: **all ZEUS data (1996-2007)**
integrated luminosity 468 pb⁻¹
- Monte Carlo samples:

- **signal**

DIFFVM exclusive VM production

- **background**

GRAPE Bethe-Heitler elastic and proton dissociative dimuon production



Event selection

- Scattered electron detected
- Scattered proton undetected
- Two reconstructed tracks identified as muons and nothing else in the detector above noise level
- Two reconstructed tracks identified as muons, two pion tracks and nothing else in the detector above noise level

$$\begin{aligned} 30 &\leq W \leq 210 \text{ GeV} \\ 2 &\leq Q^2 \leq 80 \text{ GeV}^2 \\ |t| &\leq 1 \text{ GeV}^2 \end{aligned}$$

$$\begin{aligned} \Psi(2S) &\rightarrow \mu^+ \mu^- \\ J/\psi &\rightarrow \mu^+ \mu^- \end{aligned}$$

$$\Psi(2S) \rightarrow J/\psi \pi^+ \pi^- \quad J/\psi \rightarrow \mu^+ \mu^-$$


→ Proton dissociative events removed above masses $\sim M_N 4 \text{ GeV}$
Assuming cross section ratio does not vary with M_N , **results not affected by proton dissociation background**

Signal extraction

ZEUS

$$\Psi(2S) \rightarrow \mu^+ \mu^-$$

$$J/\psi \rightarrow \mu^+ \mu^-$$




Signal extraction

ZEUS

$$\Psi(2S) \rightarrow \mu^+ \mu^-$$

$$J/\psi \rightarrow \mu^+ \mu^-$$



Dimuon **background fit** to straight line for
 $2 < M_{\mu^+ \mu^-} < 2.62 \text{ GeV}$ and $4.05 < M_{\mu^+ \mu^-} < 5 \text{ GeV}$


Function used for the fit and $M_{\mu^+ \mu^-}$ window varied as systematic checks

Signal extraction

ZEUS

$$\Psi(2S) \rightarrow \mu^+ \mu^-$$

$$J/\psi \rightarrow \mu^+ \mu^-$$



Number of events above background in the ranges

$$3.59 < M_{\mu^+ \mu^-} < 3.79 \text{ GeV} \rightarrow N_{\Psi(2S)}$$


$$3.02 < M_{\mu^+ \mu^-} < 3.17 \text{ GeV} \rightarrow N_{J/\psi}$$

Dimuon **background fit** to straight line for
 $2 < M_{\mu^+ \mu^-} < 2.62 \text{ GeV}$ and $4.05 < M_{\mu^+ \mu^-} < 5 \text{ GeV}$

Function used for the fit and $M_{\mu^+ \mu^-}$ window varied as systematic checks

Signal extraction


$$\Psi(2S) \rightarrow J/\psi \pi^+ \pi^- \quad J/\psi \rightarrow \mu^+ \mu^-$$




- ZEUS 468 pb⁻¹

$$\Delta M = M_{\mu\mu\pi\pi} - M_{\mu\mu}$$

Signal extraction



ZEUS






- ZEUS 468 pb⁻¹

$$\Delta M = M_{\mu\mu\pi\pi} - M_{\mu\mu}$$

Signal extraction

$$\Psi(2S) \rightarrow J/\psi \pi^+ \pi^- \quad J/\psi \rightarrow \mu^+ \mu^-$$

ZEUS



- ZEUS 468 pb⁻¹

$$\Delta M = M_{\mu\mu\pi\pi} - M_{\mu\mu}$$

No background (upper limit of 3 events at 90% C.L. estimated)

$$0.5 < \Delta M < 0.7 \text{ GeV} \rightarrow N_{\Psi(2S)}$$

Cut applied above:

$$3.02 < M_{\mu\mu} < 3.17 \text{ GeV}$$

Measured ratios

$$R_{J/\psi\pi\pi} = \frac{\sigma_{\psi(2S)}}{\sigma_{J/\psi(1S)}} = \frac{N_{\psi(2S)}}{N_{J/\psi(1S)}} \cdot \frac{Acc_{J/\psi(1S) \rightarrow \mu^+\mu^-}}{Acc_{\psi(2S) \rightarrow J/\psi\pi^+\pi^-}} \cdot \frac{1}{BR_{\psi(2S) \rightarrow J/\psi\pi^+\pi^-}}$$

$$R_{\mu\mu} = \frac{\sigma_{\psi(2S)}}{\sigma_{J/\psi(1S)}} = \frac{N_{\psi(2S)}}{N_{J/\psi(1S)}} \cdot \frac{Acc_{J/\psi(1S) \rightarrow \mu^+\mu^-}}{Acc_{\psi(2S) \rightarrow \mu^+\mu^-}} \cdot \frac{BR_{J/\psi(1S) \rightarrow \mu^+\mu^-}}{BR_{\psi(2S) \rightarrow \mu^+\mu^-}}$$

R_{comb} = combination of $R_{J/\psi\pi\pi}$ and $R_{\mu\mu}$

$$Acc_i = \frac{N_i^{reco}}{N_i^{true}}$$

$$BR[\psi(2S) \rightarrow J/\psi \pi\pi] = (33.6 \pm 0.4)\%$$

$$BR[\psi(2S) \rightarrow \mu\mu] = (7.7 \pm 0.8) \times 10^{-3}\%$$

$$BR[J/\psi \rightarrow \mu\mu] = (5.93 \pm 0.06)\%$$

Measured ratios

$R_{J/\psi\pi\pi}$	$0.26 \pm 0.03_{-0.01}^{+0.01}$
$R_{\mu\mu}$	$0.24 \pm 0.05_{-0.03}^{+0.02}$
R_{comb}	$0.26 \pm 0.02_{-0.01}^{+0.01}$
$R_{\psi(2S)}$	$1.1 \pm 0.2_{-0.1}^{+0.2}$

$$30 \leq W \leq 210 \text{ GeV}$$

$$2 \leq Q^2 \leq 80 \text{ GeV}^2$$


$$|t| \leq 1 \text{ GeV}^2$$

$$R_{\psi(2S)} = R_{J/\psi\pi\pi} / R_{\mu\mu}$$


Q^2 (GeV ²)	$R_{J/\psi\pi\pi}$	$R_{\mu\mu}$	R_{comb}	$R_{\psi(2S)}$
2 – 5	$0.21 \pm 0.07_{-0.03}^{+0.04}$	$0.10 \pm 0.09_{-0.09}^{+0.09}$	$0.17 \pm 0.05_{-0.02}^{+0.05}$	–
5 – 8	$0.19 \pm 0.05_{-0.02}^{+0.02}$	$0.13 \pm 0.06_{-0.03}^{+0.12}$	$0.17 \pm 0.04_{-0.02}^{+0.05}$	$1.5 \pm 0.8_{-0.7}^{+0.4}$
8 – 12	$0.27 \pm 0.05_{-0.01}^{+0.06}$	$0.29 \pm 0.08_{-0.08}^{+0.03}$	$0.28 \pm 0.05_{-0.03}^{+0.03}$	$0.9 \pm 0.3_{-0.1}^{+0.4}$
12 – 24	$0.27 \pm 0.05_{-0.03}^{+0.04}$	$0.24 \pm 0.08_{-0.08}^{+0.01}$	$0.26 \pm 0.05_{-0.03}^{+0.01}$	$1.1 \pm 0.4_{-0.1}^{+0.6}$
24 – 80	$0.56 \pm 0.13_{-0.09}^{+0.04}$	$0.42 \pm 0.17_{-0.04}^{+0.12}$	$0.51 \pm 0.10_{-0.04}^{+0.04}$	$1.3 \pm 0.6_{-0.6}^{+0.3}$
W (GeV)	$R_{J/\psi\pi\pi}$	$R_{\mu\mu}$	R_{comb}	$R_{\psi(2S)}$
30 – 70	$0.24 \pm 0.07_{-0.13}^{+0.01}$	$0.24 \pm 0.10_{-0.14}^{+0.03}$	$0.24 \pm 0.06_{-0.13}^{+0.01}$	$1.0 \pm 0.5_{-0.2}^{+0.5}$
70 – 95	$0.30 \pm 0.06_{-0.04}^{+0.01}$	$0.31 \pm 0.09_{-0.03}^{+0.09}$	$0.30 \pm 0.05_{-0.03}^{+0.02}$	$1.0 \pm 0.3_{-0.2}^{+0.1}$
95 – 120	$0.28 \pm 0.06_{-0.01}^{+0.05}$	$0.24 \pm 0.08_{-0.05}^{+0.04}$	$0.27 \pm 0.05_{-0.01}^{+0.03}$	$1.2 \pm 0.5_{-0.2}^{+0.5}$
120 – 210	$0.22 \pm 0.05_{-0.01}^{+0.07}$	$0.17 \pm 0.07_{-0.05}^{+0.02}$	$0.21 \pm 0.04_{-0.01}^{+0.03}$	$1.3 \pm 0.6_{-0.2}^{+0.7}$
$ t $ (GeV ²)	$R_{J/\psi\pi\pi}$	$R_{\mu\mu}$	R_{comb}	$R_{\psi(2S)}$
0 – 0.1	$0.23 \pm 0.05_{-0.02}^{+0.02}$	$0.23 \pm 0.09_{-0.05}^{+0.04}$	$0.23 \pm 0.04_{-0.02}^{+0.01}$	$1.0 \pm 0.4_{-0.2}^{+0.3}$
0.1 – 0.2	$0.22 \pm 0.06_{-0.03}^{+0.02}$	$0.23 \pm 0.09_{-0.06}^{+0.02}$	$0.22 \pm 0.05_{-0.02}^{+0.02}$	$0.9 \pm 0.4_{-0.2}^{+0.5}$
0.2 – 0.4	$0.27 \pm 0.06_{-0.01}^{+0.06}$	$0.18 \pm 0.07_{-0.06}^{+0.05}$	$0.24 \pm 0.04_{-0.02}^{+0.03}$	$1.5 \pm 0.6_{-0.2}^{+0.5}$
0.4 – 1	$0.32 \pm 0.06_{-0.03}^{+0.05}$	$0.30 \pm 0.08_{-0.05}^{+0.02}$	$0.32 \pm 0.05_{-0.02}^{+0.01}$	$1.1 \pm 0.3_{-0.1}^{+0.3}$

Ratios vs Q^2 , W and t


ZEUS



ZEUS




ZEUS



→ Independent of W and t

→ Increasing with Q^2

Comparison with models and with H1



HIKT, Hüfner et al.: dipole model, dipole-proton constrained by inclusive DIS data
 AR, Armesto and Rezaeian: impact parameter dependent CGC and IP-Sat model
 KMW, Kowalski Motyka Watt: QCD description and universality of quarkonia production
 FFJS, Fazio et al.: two component Pomeron model
 KNNPZZ, Nemchik et al.: dipole cross section derived from BFKL generalised eq.
 LM, Lappi and Mäntysaari : dipole picture in IP-Sat model

HIKT calculation

[J. Hüfner et al., Phys. Rev. D 62, 094022 (2000)]






FIGURE 2. Integrated cross section for elastic photoproduction with real photons ($Q^2 = 0$) calculate with GBW and KST dipole cross sections and for four potentials to generate J/ψ wave function. Experimental data points from the H1 [20], E401 [21], E516 [22] and ZEUS [23] experiments.

- Two parameterization of the $c\bar{c}$ -dipole cross section (GBW and KST)
- Four phenomenological potentials of the wave functions:
 - BT, LOG with $m_c \approx 1.5 \text{ GeV}$
 - COR and POW with $m_c \approx 1.8 \text{ GeV}$

→ BT predictions larger than the data




Comparison with models and with H1



HIKT, Hüfner et al.: dipole model, dipole-proton constrained by inclusive DIS data
 AR, Armesto and Rezaeian: impact parameter dependent CGC and IP-Sat model
 KMW, Kowalski Motyka Watt: QCD description and universality of quarkonia production
 FFJS, Fazio et al.: two component Pomeron model
 KNNPZZ, Nemchik et al.: dipole cross section derived from BFKL generalised eq.
 LM, Lappi and Mäntysaari : dipole picture in IP-Sat model

AR calculation


[N. Armesto and A. H. Reazeian, Phys. Rev. D 90, 054003 (2014)]



- Impact-parameter-dependent Color Glass Condensate model (b-CGC) or Saturation model (IP-Sat) for the calculation of the $c\bar{c}$ -dipole cross section

→ IP-Sat prediction about 30% lower and gives a better description of the data


Comparison with models and with H1



HIKT, Hüfner et al.: dipole model, dipole-proton constrained by inclusive DIS data
 AR, Armesto and Rezaeian: impact parameter dependent CGC and IP-Sat model
 KMW, Kowalski Motyka Watt: QCD description and universality of quarkonia production
 FFJS, Fazio et al.: two component Pomeron model
 KNNPZZ, Nemchik et al.: dipole cross section derived from BFKL generalised eq.
 LM, Lappi and Mäntysaari : dipole picture in IP-Sat model

KMW calculation


[H. Kowalski et al., Phys. Rev. D 74, 074016 (2006)]



- Assumes universality of production of vector quarkonia states.
Parameter δ depends on the choice of the charmonium wave function

→ $\delta = 0$ provides a better description of the data

Comparison with models and with H1



HIKT, Hüfner et al.: dipole model, dipole-proton constrained by inclusive DIS data
 AR, Armesto and Rezaeian: impact parameter dependent CGC and IP-Sat model
 KMW, Kowalski Motyka Watt: QCD description and universality of quarkonia production
 FFJS, Fazio et al.: two component Pomeron model
 KNNPZZ, Nemchik et al.: dipole cross section derived from BFKL generalised eq.
 LM, Lappi and Mäntysaari : dipole picture in IP-Sat model

Summary

- $J/\psi(2S)/J/\psi$ measured by ZEUS with full HERA statistics
- $J/\psi(2S)/J/\psi$ rises with Q^2 and is constant in W and $|t|$
- Discrimination of different models possible


DESY 16-008, Nucl. Phys. B 909 (2016) 934

Backup

FFJS calculation

[S. Fazio et al., Phys. Rev. D 90, 016007 (2014)]


- Two-component Pomeron model



KNNPZZ calculation

- [B. Kopeliovich et al., Phys. Rev D 44, 3466 (1991),
B. Kopeliovich et Al., Phys. Lett. B 324, 469 (1994)
J. Nemchik et al., Phys. Lett. B 341, 228 (1994)
J. Nemchik et al., J. Exp. Theor. Phys. 86, 1054 (1998)]


- Generalized BFKL equation for $ccba$ - dipole cross section






LM calculation

[T. Lappi and H. Mäntysaari, Phys. Rev. C 83, 065202 (2011),
T. Lappi and H. Mäntysaari, PoS (DIS2014), 069 (2014)]

- BFKL equation + IP-Sat






ZEUS



$J/\psi(1S) \rightarrow \mu^+ \mu^-$
 • ZEUS 468 pb⁻¹
 ■ DIFFVIM + BH
 ▨ BH

Monte Carlo reweighted in t , Q^2 and angular distributions

ZEUS






$\Psi(2S) \rightarrow \mu^+ \mu^-$
 • ZEUS 468 pb⁻¹
 ■ DIFFVIM + BH
 ▨ BH

Monte Carlo reweighted in t , Q^2 and angular distributions

Data/MC

$\Psi(2S) \rightarrow J/\psi \pi^+ \pi^- \quad J/\psi \rightarrow \mu^+ \mu^-$

ZEUS




$\Psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-$

• ZEUS 468 pb⁻¹

■ DIFFVM

Monte Carlo reweighted in t , Q^2 and angular distributions

Comparison with H1 earlier measurement



H1 collaboration:

Eur.Phys.J.C10:373-393,1999

→ Much larger luminosity in ZEUS measurement (HERA I + HERA II)