

# Triple parton scattering effects in $D$ -meson production at the LHC

Rafał Maciuta

Institute of Nuclear Physics PAN, Kraków, Poland

The 17th Conference on Elastic and Diffractive Scattering, EDS Blois 2017

26th – 30th June 2017, Prague, Czech Republic




# Multiple-Parton Interactions (MPI) at the LHC

## Our previous studies:

Hard **Double-Parton Scattering (DPS)** effects in proton-proton collisions:

- $pp \rightarrow 4\text{jets } X$  (needs dedicated experimental analyses: ATLAS, CMS)  
Phys.Lett. B749 (2015) 57-62; Phys.Rev. D94 (2016) no.1, 014019
- $pp \rightarrow c\bar{c} + 2\text{jets } X$  (experimentally available at ATLAS) soon on arXiv
- $pp \rightarrow c\bar{c}c\bar{c} X$  (predictions confirmed by the LHCb double charm data)  
Phys.Rev. D87 (2013) no.7, 074039; Phys.Lett. B758 (2016) 458-464
- $pp \rightarrow c\bar{c}b\bar{b} X$  (experimentally available at LHCb) soon on arXiv
- $pp \rightarrow b\bar{b}b\bar{b} X$  (experimentally available at LHCb) soon on arXiv

In this talk: **Triple-parton scattering (TPS)** in  $pp \rightarrow c\bar{c}c\bar{c}c\bar{c} X$





Can we observe some evidence of TPS effects at the LHC in the case of triple charm production?



# Multiple charm quark production at the LHC

## Double-parton scattering in double $c\bar{c}$ pair production



Luszczak, Maciula, Szczurek, Phys.Rev. D85 (2012) 094034)



Total cross sections for SPS single  $c\bar{c}$  and DPS double  $c\bar{c}$  become comparable at LHC energies

## Triple-parton scattering in triple $c\bar{c}$ pair production

d'Enterria, Snigirev, Phys. Rev. Lett. 118, 122001 (2017)



Total cross section for TPS triple  $c\bar{c}$  at LHC energies become sizeable ( $\sim 100 \mu\text{b}$ )

Can we observe the TPS effects experimentally?




# The idea behind

**Our goal:** Reliable predictions for  $D^0 D^0 D^0$  production for the LHCb experiment


- we try to stay in contact with experimental data for SPS single  $D^0$  and DPS double  $D^0$  production to minimize the theoretical uncertainties.

SPS single  $D^0$




⇒

DPS double  $D^0$



⇒

TPS triple  $D^0$



- theoretical uncertainties in the SPS get amplified by factors of 2 and 3 for the DPS and TPS
- use the model calculation that provides a good (of the same quality) parametrizations of both the single and the double charm LHCb data

The LHCb single and double charm data well described within the  $k_T$ -factorization approach with standard set of pQCD calculation parameters

# Theoretical framework

## $k_T$ -factorization approach $\rightarrow k_{1,t}, k_{2,t} \neq 0$

e.g. Collins-Ellis, Nucl. Phys. B360 (1991) 3; Catani-Ciafaloni-Hautmann, Nucl. Phys. B366 (1991) 135;


- exact kinematics from the very beginning and additional hard dynamics coming from transverse momenta of incident partons
- very efficient approach for correlation studies

multi-differential cross section:


$$\frac{d\sigma}{dy_1 dy_2 d^2 p_{1,t} d^2 p_{2,t}} = \int \frac{d^2 k_{1,t}}{\pi} \frac{d^2 k_{2,t}}{\pi} \frac{1}{16\pi^2 (x_1 x_2 s)^2} \overline{|\mathcal{M}_{g^*g^* \rightarrow Q\bar{Q}}|^2} \times \delta^2(\vec{k}_{1,t} + \vec{k}_{2,t} - \vec{p}_{1,t} - \vec{p}_{2,t}) \mathcal{F}_g(x_1, k_{1,t}^2) \mathcal{F}_g(x_2, k_{2,t}^2)$$

- leading-order **off-shell** matrix elements  $\overline{|\mathcal{M}_{g^*g^* \rightarrow Q\bar{Q}}|^2}$
- $\mathcal{F}_g(x, k_T^2)$  - **transverse momentum dependent** gluon distributions
- part of **higher-order real corrections** effectively included


pair creation  
with gluon emission




flavour excitation



gluon splitting



# Unintegrated gluon distribution functions (uGDFs)



Most popular models:

- Kwieciński, Jung (CCFM, wide range of  $x$ )
- Kimber-Martin-Ryskin (DGLAP-BFKL, wide range of  $x$ )
- Kwieciński-Martin-Staśto (BFKL-DGLAP, small  $x$ -values)
- Kutak-Staśto (BK, saturation, only small  $x$ -values)

We use: **Kimber-Martin-Ryskin (KMR) approach:**

$$f_g(x, k_T^2, \mu^2) \equiv \frac{\partial}{\partial \log k_T^2} \left[ g(x, k_T^2) T_g(k_T^2, \mu^2) \right] = T_g(k_T^2, \mu^2) \frac{\alpha_s(k_T^2)}{2\pi} \sum_b \int_x^1 dz P_{gb}(z) b\left(\frac{x}{z}, k_T^2\right)$$

$$f_g(x, k_T^2, \mu^2) = T_g(k_T^2, \mu^2) \frac{\alpha_s(k_T^2)}{2\pi} \times \int_x^1 dz \left[ \sum_q P_{gq}(z) \frac{x}{z} q\left(\frac{x}{z}, k_T^2\right) + P_{gg}(z) \frac{x}{z} g\left(\frac{x}{z}, k_T^2\right) \Theta\left(\frac{\mu}{\mu + k_T} - z\right) \right]$$

$$\mathcal{F}_g(x, k_T^2, \mu^2) \equiv \frac{1}{k_T^2} f_g(x, k_T^2, \mu^2) .$$



- calculated from collinear PDFs (most up-to-date PDF sets can be used)  
we take: MMHT2014lo gluon PDF
- unintegrated quarks available (important for reliable predictions e.g. for jets)
- unique feature: possible additional hard emission from the uPDF  
(part of higher-order real corrections)



Inclusive  $D^0$  production

$$\sqrt{s} = 7, 13 \text{ TeV}$$

The LHCb fiducial volume under interest:  $3 < p_T^D < 12 \text{ GeV}$ ,  $2 < y^D < 4$




- a very good description of the LHCb inclusive charm data with the central value of the standard uncertainty bands at both energies
- the usual variation of these parameters is not necessary in order to stay in touch with the data
- The  $\pm 40\%$  uncertainty is related with factorization/renormalization scales and charm quark mass and will be propagated to the double and triple meson case

Then such a parametrization of the data can be used for the predictions for **DPS double** and **TPS triple  $D^0$**  production



# Double-parton scattering (DPS) mechanism




**DPS in general form** for  $pp \rightarrow c \bar{c} c \bar{c} X$ :

$$d\sigma^{DPS} = \frac{1}{2} \cdot \Gamma_{gg}(b, x_1, x_2; \mu_1^2, \mu_2^2) \Gamma_{gg}(b, x'_1, x'_2; \mu_1^2, \mu_2^2) \\ \times d\sigma_{ij \rightarrow k}(x'_1, x_1, \mu_1^2) \cdot d\sigma_{gg \rightarrow c\bar{c}}(x_2, x'_2, \mu_2^2) dx_1 dx_2 dx'_1 dx'_2 d^2b$$

**dPDF** - emission of one parton with assumption that second parton is also emitted

$$\Gamma_{gg}(b, x_1, x_2; \mu_1^2, \mu_2^2) = F_g(x_1, \mu_1^2) F_g(x_2, \mu_2^2) F(b; x_1, x_2, \mu_1^2, \mu_2^2)$$

- longitudinal and transverse correlations between two partons
- spin, flavor and color correlations
- well established theory: e.g. Diehl, Ostermeier, Schafer, JHEP 03, 089 (2012)  
but not yet available for phenomenological studies



## Factorized ansatz (pocket-formula)

In a simple probabilistic picture process initiated by:

**two simultaneous hard parton-parton scatterings in one proton-proton interaction**

$$\sigma^{DPS} = \frac{1}{2\sigma_{eff}} \cdot \sigma^{SPS}(g g \rightarrow c \bar{c}) \cdot \sigma^{SPS}(g g \rightarrow c \bar{c})$$

**two subprocesses are not correlated and do not interfere**

- $\sigma_{eff} \Rightarrow$  model parameter  $\Rightarrow$  normalization of  $\sigma^{DPS}$






# Double-parton scattering (DPS) mechanism

## Factorized ansatz (pocket-formula)

- a good approximation for **small-x partons**
- **color/flavor correlations suppressed** in evolution (Kasemets et al., Phys. Rev. D91, 014015 (2015))
- **spin (polarization) correlations very small** (Echevarria et al. JHEP 04, 034 (2015))


## Separation of longitudinal and transverse degrees of freedom

- **dPDFs in multiplicative form:**  $\Gamma_{ij}(b; x_1, x_2, \mu_1^2, \mu_2^2) = F_i(x_1, \mu_1^2)F_j(x_2, \mu_2^2)F(b)$
- only transverse correlations taken into account
- $\sigma_{eff} = \left[ \int d^2b (F(b))^2 \right]^{-1}$ ,  $F(b)$  - overlap of the matter distribution in transverse plane where  $b$  is a distance between both partons
- nonperturbative quantity with dimension of cross section, connected to transverse size of proton



- extracted from several experimental analyses
- in principle may not be universal
- detailed studies: Seymour, Siódmok, JHEP 10, 113 (2013)
- LHCb double charm data:  $\sigma_{eff} = 21_{-6}^{+7}$  mb
- ATLAS 4jets data:  $\sigma_{eff} = 14.9$  mb
- **world average:**  $\sigma_{eff} \approx 15$  mb (large uncertainties)



SPS vs. DPS: Inclusive  $c\bar{c}c\bar{c}$ LHCb at  $\sqrt{s} = 7$  TeV

## CHARM MESON-MESON pair production:

DD pairs – both mesons containing c-quarks



- impossible to be produced within standard SPS single  $c\bar{c}$  mechanism
- SPS double charm very small

First measurement by LHCb: J. High Energy Phys. 06, 141 (2012)

Cross section much larger than the SPS predictions

⇒ clear evidence for DPS?

Mode	$\sigma$ [nb]
$D^0D^0$	$690 \pm 40 \pm 70$
$D^0\bar{D}^0$	$6230 \pm 120 \pm 630$




Łuszczak, Maciuta, Szczurek, Phys.Rev. D85 (2012) 094034

Maciuta, Szczurek, Phys.Rev. D87 (2013) no.7, 074039

Hameren, Maciuta, Szczurek, Phys.Rev. D89 (2014) no.9, 094019



# Triple charm in triple-parton scattering (TPS)



## First theoretical analysis for charm quarks:

d'Enterria, Snigirev, *Phys.Rev.Lett.* 118, no. 12, 122001 (2017)

- a generic expressions to compute TPS cross sections
- total charm quark cross sections in NNLO collinear approach

## Our calculations:

Maciula, Szczurek, *arXiv:1703.07163 (hep-ph)*

- analysis for triple  $D$  meson production
- $k_T$ -factorization approach
- differential distributions


## TPS cross section in a general form:

$$\begin{aligned}
 \sigma_{pp \rightarrow c\bar{c}c\bar{c}c\bar{c}}^{\text{TPS}} &= \left(\frac{1}{3!}\right) \int \Gamma_p^{ggg}(x_1, x_2, x_3; \bar{b}_1, \bar{b}_2, \bar{b}_3; \mu_1^2, \mu_2^2, \mu_3^2) \\
 &\times \hat{\sigma}_{c\bar{c}}^{gg}(x_1, x'_1, \mu_1^2) \hat{\sigma}_{c\bar{c}}^{gg}(x_2, x'_2, \mu_2^2) \hat{\sigma}_{c\bar{c}}^{gg}(x_3, x'_3, \mu_3^2) \\
 &\times \Gamma_p^{ggg}(x'_1, x'_2, x'_3; \bar{b}_1 - \bar{b}, \bar{b}_2 - \bar{b}, \bar{b}_3 - \bar{b}; \mu_1^2, \mu_2^2, \mu_3^2) \\
 &\times dx_1 dx_2 dx_3 dx'_1 dx'_2 dx'_3 d^2b_1 d^2b_2 d^2b_3 d^2b
 \end{aligned}$$

- triple-parton distribution functions (triple PDFs)
- adoption to real calculations much more limited than in the case of dPDFs



# Triple charm in triple-parton scattering (TPS)



## First theoretical analysis for charm quarks:

d'Enterria, Snigirev, *Phys.Rev.Lett.* 118, no. 12, 122001 (2017)

- a generic expressions to compute TPS cross sections
- total charm quark cross sections in NNLO collinear approach

## Our calculations:

Maciula, Szczurek, *arXiv:1703.07163 (hep-ph)*

- analysis for triple  $D$  meson production
- $k_T$ -factorization approach
- differential distributions

## Factorized ansatz for TPS (pocket-formula)

In a simple probabilistic picture process initiated by:

three simultaneous hard parton-parton scatterings in one proton-proton interaction

$$\sigma_{pp \rightarrow c\bar{c}c\bar{c}c\bar{c}}^{\text{TPS}} = \left( \frac{1}{3!} \right) \frac{\sigma_{pp \rightarrow c\bar{c}}^{\text{SPS}} \cdot \sigma_{pp \rightarrow c\bar{c}}^{\text{SPS}} \cdot \sigma_{pp \rightarrow c\bar{c}}^{\text{SPS}}}{\sigma_{\text{eff,TPS}}^2}$$



three subprocesses are not correlated and do not interfere



- $\sigma_{\text{eff,TPS}} = k \times \sigma_{\text{eff,DPS}}$ , with  $k = 0.82 \pm 0.11$
- $\sigma_{\text{eff,DPS}} = 21 \text{ mb}$  (extracted from LHCb data)  $\Rightarrow \sigma_{\text{eff,TPS}} = 17 \text{ mb}$



Inclusive  $D^0 D^0 D^0$  $\sqrt{s} = 7, 13 \text{ TeV}$ 

The transverse momentum distribution of one of the  $D^0$  meson  
from the  $D^0 D^0$  pair or from the  $D^0 D^0 D^0$  triplet



Inclusive  $D^0 D^0 D^0$  $\sqrt{s} = 7, 13 \text{ TeV}$ The invariant mass distribution of  $D^0 D^0$  pair and of  $D^0 D^0 D^0$  triplet

Inclusive  $D^0 D^0 D^0$ 

$$\sqrt{s} = 7, 13 \text{ TeV}$$

**The integrated cross sections** for double and triple  $D^0$  meson production (in nb) within the LHCb acceptance:  $2 < y_{D^0} < 4$  and  $3 < p_T^{D^0} < 12 \text{ GeV}$  calculated in the  $k_T$ -factorization approach.

Final state	$\sqrt{s} = 7 \text{ TeV}$	$\sqrt{s} = 13 \text{ TeV}$
DPS: $\sigma(D^0 D^0 + X)$	784.74	2992.91
TPS: $\sigma(D^0 D^0 D^0 + X)$	2.38	17.71

**Number of events** for different values of the feasible integrated luminosity in the LHCb experiment for the calculated cross sections

$\sqrt{s}$	Integrated Luminosity	DPS ( $D^0 D^0$ )	TPS ( $D^0 D^0 D^0$ )
7 TeV	355 $\text{pb}^{-1}$	$0.43 \times 10^6$	51
	1106 $\text{pb}^{-1}$	$1.34 \times 10^6$	159
13 TeV	1665 $\text{pb}^{-1}$	$7.70 \times 10^6$	1789
	5000 $\text{pb}^{-1}$	$23.11 \times 10^6$	5374

- a few thousands of events of triple  $D^0$  production at  $\sqrt{s} = 13 \text{ TeV}$



# Conclusions

We have presented first estimation of the triple-parton scattering production of triple open charm meson at the LHC:

$$pp \rightarrow D^0 D^0 D^0 X$$

- expected cross section: tens of nanobarns (with precision of factor  $\approx 3$ )
- a few thousands of events of triple  $D^0$  production can be observed at  $\sqrt{s} = 13$  TeV within the LHCb detector acceptance

Thank You for attention!

