ACCELERATOR PHYSICS AND TECHNOLOGY – EPISODE I

Ezio Todesco
CERN, Technology Department
Magnet Superconductors and Cryostat Group

FOREWORD

- The science of superconducting magnets is a exciting, fancy and dirty mixture of physics, engineering, and chemistry
  - Chemistry and material science: the quest for superconducting materials with better performances
  - Quantum physics: the key mechanisms of superconductivity
  - Classical electrodynamics: magnet design
  - Mechanical engineering: support structures
  - Electrical engineering: powering of the magnets and their protection
  - Cryogenics: keep them cool …

- The cost optimization also plays a relevant role
  - Keep them cheap …

FOREWORD

- An example of the variety of the issues to be taken into account
  - The field of the LHC dipoles (8.3 T) is related to the critical field of Niobium-Titanium (Nb-Ti), which is determined by the microscopic quantum properties of the material



Quantized fluxoids penetrating a superconductor used in accelerator magnets

A 15m truck unloading a 27 tons LHC dipole

  - The length of the LHC dipoles (15 m) has been determined by the maximal dimensions of (regular) trucks allowed on European roads
- This makes the subject complex, challenging and complete for the formation of a (young) physicist or engineer

FOREWORD

- The size of our objects
  - Length of an high energy physics accelerator: ~Km

RHIC ring at BNL, Long Island, US

Main ring at Fermilab, Chicago, US

FOREWORD

- The size of our objects
  - Length of an accelerator magnet: ~10 m
  - Diameter of an accelerator magnet: ~m
  - Beam pipe size of an accelerator magnet: ~cm

https://www.youtube.com/watch?v=KKFnsFFdPh8

Unloading a 27 tons dipole



Dipoles in the LHC tunnel, Geneva, CH



A stack of LHC dipoles, CERN, Geneva, CH

CONTENTS

The synchrotron and its magnets

How to generate magnetic fields

What superconductivity gives

REMINDER: THE SYNCHROTRON AND ITS MAGNETS

- Electro-magnetic field accelerates particles
- Magnetic field steers the particles in a closed (~circular) orbit

  to drive particles through the same accelerating structure several times



  - Most of the accelerator bends, a small part increases the energy
  - As the particle is accelerated, its energy increases and the magnetic field is increased ("synchro") to keep the particles on the same orbit

- What are the limitations to increase the energy ?
  - Proton machines: the maximum field of the dipoles (LHC, Tevatron, SPS …)
  - Electron machines: the synchrotron radiation due to bending trajectories (LEP)

REMINDER: THE SYNCHROTRON AND ITS MAGNETS

- The arcs: bending the beam → energy







A schematic view of a synchrotron

Dipoles for bending Quadrupoles for focusing

Sextupoles, octupoles … for correcting

- Long straight sections (LSS) → luminosity
  - Interaction regions (IR) housing the experiments
    - Solenoids (detector magnets) acting as spectrometers
    - Quadrupole triplet to squeeze the beams in collision
  - Regions for other services
    - Beam injection and dump (dipole kickers)
    - Accelerating structure (RF cavities) and beam cleaning (collimators)



The lay-out of the LHC

REMINDER: THE SYNCHROTRON AND ITS MAGNETS

- Why do we need many km to get a few TeV?
  - Dynamics ruled by Lorentz force

$$\vec{F} = e\vec{v} \times \vec{B}$$

$$\vec{p} = m\gamma\vec{v}$$

$$\gamma = \frac{1}{\sqrt{1-\frac{v^2}{c^2}}}$$

$$\left|\frac{d\vec{v}}{dt}\right| = \frac{v^2}{\rho}$$





Hendrik Antoon Lorentz, Dutch

(18 July 1853 – 4 February 1928), painted by Menso Kamerlingh Onnes, brother of Heinke, who discovered superconductivity

$$F = evB$$

$$\vec{F} = \frac{d}{dt}p = m\frac{d}{dt}(\gamma v) \sim m\gamma\frac{d}{dt}v$$

$$eB = m\gamma\frac{v}{\rho} = \frac{p}{\rho}$$

$$F = m\gamma\left|\frac{d\vec{v}}{dt}\right| = m\gamma\frac{v^2}{\rho}$$

$$p = eB\rho$$

$$E[GeV] = 0.3 \times B[T] \times \rho[m]$$

REMINDER: THE SYNCHROTRON AND ITS MAGNETS

- In many textbooks the gamma is attached to the mass to create the concept of relativistic mass

$$\vec{p} = m\gamma\vec{v} \qquad \gamma = \frac{1}{\sqrt{1-\frac{v^2}{c^2}}} \qquad m_R = m\gamma$$

  - So momentum equation is as in Newton
  - So we «understand» that speed of light cannot be reached because particle mass go to infinity
- It is a rather misleading concept
  - We lose mass invariance …
  - Einstein did not like it
  - I would suggest avoid using it

It is not good to introduce the concept of the mass of a moving body for which no clear definition can be given. It is better to introduce no other mass concept than the 'rest mass' *m*. Instead of introducing *M* it is better to mention the expression for the momentum and energy of a body in motion.
— *Albert Einstein in letter to Lincoln Barnett, 19 June 1948 (quote from L. B. Okun (1989), p. 42*

TERMINATOR-3 INTERLUDE

We analyse the accelerator shown in Terminator-3 [Warner Bros., Columbia Pictures, 2003]

- Estimation of the magnetic field

$$E[GeV] = 0.3 \times B[T] \times \rho[m]$$

  - Energy = 5760 GeV
  - Radius ~30 m
  - Field = 5760/0.3/30 ~ 640 T (a lot !)

- Is it possible to have 640 T magnets ??
  - Or is it science-fiction?



5.76 TeV nominal energy

A 200 m ring ?

REMINDER: THE SYNCHROTRON AND ITS MAGNETS

- Relation momentum-magnetic field-orbit radius
  - Having 8 T magnets, we need 3 Km curvature radius to have 7 TeV
  - If we would have 800 T magnets, 30 m would be enough …
  - We will show why 8 T is the present limit for accelerator magnets



$$E[GeV] = 0.3 \times B[T] \times \rho[m]$$



CONTENTS

GENERATION OF MAGNETIC FIELDS: BIOT-SAVART LAW

- A magnetic field is generated by two mechanisms
  - An electrical charge in movement (macroscopic current)
  - Coherent alignment of atomic magnetic momentum (ferromagnetic domains)



Félix Savart, French
(June 30, 1791-March 16, 1841)

- Biot-Savart law: magnetic field generated by a current line is

$$|B| = \frac{I\mu_0}{2\pi\rho}$$

  - Proportional to current
  - Inversely proportional to distance
  - Perpendicular to current direction and distance



Jean-Baptiste Biot, French
(April 21, 1774 – February 3, 1862)

GENERATION OF MAGETIC FIELDS: FIELD OF A WINDING



- Magnetic field generated by a winding
  - We compute the central field given by a sector dipole with uniform current density j

$$|B| = \frac{I\mu_0}{2\pi\rho} \qquad I \rightarrow j\rho d\rho d\theta$$

$$B = -4\frac{j\mu_0}{2\pi}\int_0^{\alpha}\int_r^{r+w}\frac{\cos\theta}{\rho}\rho d\rho d\theta = -\frac{2j\mu_0}{\pi}w\sin\alpha$$



    - Setting α=60° one gets a more uniform field

  - $B \propto$ current density (obvious)
  - $B \propto$ coil width w (less obvious)
  - B is independent of the aperture r (much less obvious)

$$B[T] \approx 7\times 10^{-4}\, j[\mathrm{A/mm}^2]\, w[\mathrm{mm}]$$

GENERATION OF MAGETIC FIELDS: SUPERCONDUCTORS VERSUS NORMAL CONDUCTORS

- Magnetic field generated by a winding of width w

$$B[T] \approx 7 \times 10^{-4}\ j[\mathrm{A/mm}^2]w[\mathrm{mm}]$$



  - The current density in copper for typical wires used in transmission lines is ~ 5 [A/mm^2]
  - Using special techniques for cooling one can arrive up to ~ 100 [A/mm^2]
  - Superconductors allow current densities in the sc material of ~1000 [A/mm^2]
    - Example: LHC dipoles have j_{sc}=1500 A/mm^2
      j=360 A/mm^2 , (~ ¼ of the cable made by sc !)
      Coil width w~30 mm, B~8 T



  - There is still a factor 10, and moreover the normal conducting consumes a lot of power …

GENERATION OF MAGETIC FIELDS: IRON DOMINATED ELECTROMAGNETS

- Normal conducting magnets for accelerators are made with a copper winding around a ferromagnetic core that greatly enhances the field
  - This is a very effective and cheap design



- The shape of the pole gives the field homogeneity
  - The limit is given by the iron saturation, i.e. 2 T
  - This limit is due to the atomic properties, i.e. it looks like a hard limit

- Therefore, superconducting magnets today give a factor ~4 larger field than normal conducting – not so bad anyway …
    - LHC with 2 T magnets would be 100 Km long, and it would not fit between the lake and the Jura …



TESLA INTERLUDE



Nikolai Tesla (10 July 1856 - 7 January 1943)

- Born at midnight during an electrical storm in Smiljan near Gospić (now Croatia)
- Son of an orthodox priest
- A national hero in Serbia – but also in the other republics of ex-Yugoslavia

Career

- Polytechnic in Gratz (Austria) and Prague
- Emigrated in the States in 1884
- Electrical engineer
- Inventor of the alternating current induction motor (1887)
- Author of 250 patents





A rather strange character, a lot of legends on him …
Check on the web ! (wikipedia, etc …)

CONTENTS

SUPERCONDUCTIVITY

- 104 years ago, in 1911, Kamerlingh Onnes discovers the superconductivity of mercury
  - Below 4.2 K, mercury has a non measurable electric resistance – not very small, but zero !
  - This discovery has been made possible thanks to his efforts to liquifying Helium, a major technological advancement needed for the discovery
  - 4.2 K is called the critical temperature: below it the material is superconductor

Heinke Kamerlingh Onnes

(18 July 1853 – 4 February 1928)

Nobel prize 1913

- Superconductivity has been discovered in other elements, with critical temperatures ranging from a few K (low temp. sc) to up to 150 K (high temperature sc)

- The behaviour has been modeled later in terms of quantum mechanics
  - Electron form pairs (Cooper pairs) that act as a boson, and "freely" move in the superconductor without resistance
  - Several Nobel prizes have been awarded in this field …



SUPERCONDUCTIVITY

- 1950: Ginzburg and Landau propose a macroscopic theory (GL) for superconductivity
  - Nobel prize in 2003 to Ginzburg, Abrikosov, Leggett

Ginzburg and Landau (circa 1947)

- 1957: Bardeen, Cooper, and Schrieffer publish microscopic theory (BCS) of Cooper-pair formation in low-temperature superconductors
  - Nobel prize in 1972

Bardeen, Cooper and Schrieffer

- 1986: Bednorz and Muller discover superconductivity at high temperatures in layered materials having copper oxide planes
  - Nobel prize in 1986 (a fast one …)



George Bednorz and Alexander Muller

SUPERCONDUCTIVITY

- The quest for the Holy Graal of superconductivity at higher temperatures
  - LTS: Low Temperature Superconductors (below 30 K)
  - HTS: High Temperature Superconductors (above 30 K)
- Two main application: power lines and magnets – radically different
  - Power lines: no field or absent field, possibly high T to simplify cooling
  - Magnets: have "enough" current density able to stay in large field, working at low T is not a problem





Courtesy from J. Schwartz, CERN academic training 2012

https://indico.cern.ch/conferenceDisplay.py?confId=158073

SUPERCONDUCTIVITY

For making magnets, our Holy Graal is having ability to survive magnetic fields

- Type I superconductors: they expel magnetic field (example: Hg)
  - They cannot be used for building magnets
- Type II superconductors: they do not expel magnetic field (example: Nb-Ti)
  - The magnetic field penetrates locally in very tiny quantized vortex

$$\phi_0 = \frac{h}{2e}$$

  - The current acts on the fluxoids with a Lorentz force that must be balanced, otherwise they start to move, dissipate, and the superconductivity is lost
  - The more current density, the less magnetic field, and viceversa → concept of critical surface



Artist view of flux penetration in a type II superconductor



First image of flux penetration, U. Essmann and H. Trauble Max-Planck Institute, Stuttgart Physics Letters 24A, 526 (1967)

SUPERCONDUCTIVITY

- The magnetic field penetrates locally in very tiny quantized vortex
- The current acts on the fluxoids with a Lorentz force that must be balanced, otherwise they start to move, dissipate, and the superconductivity is lost



Artist view of flux penetration in a type II superconductor and resulting Lorentz force

- The sc material is built to have a strong pinning force to counteract fluxoid motion
  - Pinning centers are generated with imperfections in the lattice
  - This is sometimes done with doping
  - It is a very delicate and fascinating cooking …



Optimal doping of HTS with NbO_3 to improve critical current [B. Li, et al., Physica D (2012) in press]

SUPERCONDUCTIVITY

- The material is superconductor as long as *B*, *j*, and temperature stay below the critical surface
  - The maximum current density ~ 10 000 A/mm^2, but this at zero field and zero temperature
  - In a magnet, the winding has a current density to create a magnetic field → the magnetic field is also in the winding → this reduces the current density
- Operational temperature
  - The lowest the better … but not at 0 K !
    - Specific heats go to zero
  - Many machines run at 4.2 K (liquid He)
  - LHC has been the first accelerator to operate at 1.9 K (after Tore Supra tokamak)

    ☺ Superfluid helium ! (second purely quantum effect on which LHC technology relies daily)



Critical surface for Nb-Ti

Tore Supra Tokamak

SUPERCONDUCTIVITY

- Critical current density vs. field for different materials (semilog scale) at 4.2 K
  - To remember: more critical current density, less field



Critical current density in the superconductor versus field for different materials at 4.2 K [P. J. Lee, et al]

SUMMARY

- Principles of magnets
  - Why superconducting magnets are very effective
  - The mechanisms behind superconductivity
- Superconductivity is based on couples and relies on defects
  - And gives many Nobel prizes …
- Some features of the design
- Coming soon
  - Why 8 T is the present limit for Nb-Ti
  - Why Ms. Terminator sticks on the T3 accelerator dipoles
  - Going to larger fields: other materials

REFERENCES

- Books
  - M. N. Wilson, “Superconducting magnets”, Oxford University Press, London (1976)
  - K. H. Mess, P. Schmuser, S. Wolff, “Superconducting accelerator magnets”, World Scientific, Singapore (1996).
  - A. Devred, “Practical low temperature superconductors for electromagnets”, CERN Yellow report 2004-006.
  - For superconductivity, check the last chapter of 3rd volume of Feynmann lectures!
- Review paper
  - L. Bottura, L. Rossi, “Superconducting magnets for particle accelerators”, *Rev. Sci. Accel. Tech.* **5** 30003 (2012)
  - A. Tollestrup, E. Todesco, `The development of superconducting magnets for use in particle accelerators: from Tevatron to the LHC', *Rev. Sci. Accel. Tech.* **1** 185-210 (2008)

ACKNOWLEDGEMENTS

- T. Taylor, L. Rossi, P. Lebrun, L. Bottura who gave the lectures in 2004-6, 2010-11, from which I took material and ideas
- P. Ferracin and S. Prestemon for the material prepared for the US Particle Accelerator School
- www.wikipedia.org for most of the pictures of the scientists
- Google Earth for the images of accelerators in the world
- The Nikolai Tesla museum of Belgrade, for brochures, images, and information, and the anonymous guard I met in August 2002
- Warner Bros. and Columbia Pictures for some images of Terminator-3: the rise of machines, by J. Mostow