

4-dimensional angular analysis of $\bar{B} \rightarrow D^ \ell^- \bar{\nu}_\ell$
with hadronic tagging at BABAR*

[arXiv:1903.10002]

Biplab Dey

(on behalf of the *BaBar* Collaboration)




EPS-HEP 2019, Ghent
July 12th, 2019

Motivation: $\bar{B} \rightarrow D^* \ell^- \bar{\nu}_\ell$ angular analysis

- Along with semi-tauonic case, $\ell \in \{e, \mu\}$ hold puzzles as well.
- $|V_{cb}|$ tension, higher order HQET corrections for the **form-factors**, ...
- Experimentally very **clean**, and fully **exclusive** final-state with hadronic tagging.
- With the narrow D^* vector meson, excellent system to probe NP on the leptonic side.

The 4-d kinematic variables



- 4-body decay topology
- $\sqrt{q^2}$: di-lepton mass.
- 3 angles: $\Omega \in \{\theta_\ell, \theta_V, \chi\}$
- Spin-1 D^* retains full **spin info** of the recoiling W^* in $b \rightarrow cW^{*-}$, unlike spin-0 D , where this info is reduced \Rightarrow **richer pheno!**

- Two **complementary approaches** employed in the analysis:
 - Angular observables extracted using **moments method** in $10 q^2$ bins.
 - **Unbinned** z -expansion **FF fit** to the form-factors (today).
- In either case, we perform a **full 4-d analysis**.

The generic 4-d pdf [\[arXiv:1505.02873\]](https://arxiv.org/abs/1505.02873)

- Differential rate (4-d fit pdf):

$$\frac{d\Gamma}{dq^2 d\Omega} \propto \sum_{i=1}^{14} f_i(\Omega) \Gamma_i(q^2)$$

- Transversity q^2 amplitudes:

$$H_0(q^2) \equiv h_0$$

$$H_{\{\parallel, \perp\}}(q^2) \equiv h_{\{\parallel, \perp\}} \underbrace{e^{i\delta_{\{\parallel, \perp\}}}}_{\text{NP phase}}$$

- Orthonormal angular basis:

- $Y_l^m \equiv Y_l^m(\theta_\ell, \chi)$
- $P_l^m \equiv \sqrt{2\pi} Y_l^m(\theta_V, 0)$

i	$f_i(\Omega)$	$\Gamma_i^{\text{tr}}(q^2)/(\mathbf{k}q^2)$
1	$P_0^0 Y_0^0$	$h_0^2 + h_{\parallel}^2 + h_{\perp}^2$
2	$P_2^0 Y_0^0$	$-\frac{1}{\sqrt{5}}(h_{\parallel}^2 + h_{\perp}^2) + \frac{2}{\sqrt{5}}h_0^2$
3	$P_0^0 Y_2^0$	$\frac{1}{2\sqrt{5}}[(h_{\parallel}^2 + h_{\perp}^2) - 2h_0^2]$
4	$P_2^0 Y_2^0$	$-\frac{1}{10}(h_{\parallel}^2 + h_{\perp}^2) - \frac{2}{5}h_0^2$
5	$P_2^1 \sqrt{2} \text{Re}(Y_2^1)$	$-\frac{3}{5}h_{\parallel} h_0 \cos \delta_{\parallel}$
6	$P_2^1 \sqrt{2} \text{Im}(Y_2^1)$	$\frac{3}{5}h_{\perp} h_0 \sin \delta_{\perp}$
7	$P_0^0 \sqrt{2} \text{Re}(Y_2^2)$	$-\frac{3}{2\sqrt{15}}(h_{\parallel}^2 - h_{\perp}^2)$
8	$P_2^0 \sqrt{2} \text{Re}(Y_2^2)$	$\frac{\sqrt{3}}{10}(h_{\parallel}^2 - h_{\perp}^2)$
9	$P_0^0 \sqrt{2} \text{Im}(Y_2^2)$	$\sqrt{\frac{3}{5}}h_{\perp} h_{\parallel} \sin(\delta_{\perp} - \delta_{\parallel})$
10	$P_2^0 \sqrt{2} \text{Im}(Y_2^2)$	$-\frac{\sqrt{3}}{5}h_{\perp} h_{\parallel} \sin(\delta_{\perp} - \delta_{\parallel})$
11	$P_0^0 Y_1^0$	$-\sqrt{3}h_{\perp} h_{\parallel} \cos(\delta_{\perp} - \delta_{\parallel})$
12	$P_2^0 Y_1^0$	$\frac{3}{\sqrt{15}}h_{\perp} h_{\parallel} \cos(\delta_{\perp} - \delta_{\parallel})$
13	$P_2^1 \sqrt{2} \text{Re}(Y_1^1)$	$\frac{3}{\sqrt{5}}h_{\perp} h_0 \cos \delta_{\perp}$
14	$P_2^1 \sqrt{2} \text{Im}(Y_1^1)$	$-\frac{3}{\sqrt{5}}h_{\parallel} h_0 \sin \delta_{\parallel}$


The data sample: hadronic tagging

- We use 426 fb^{-1} full *BABAR* data: $\Upsilon(4S) \rightarrow B_{\text{tag}} \bar{B}_{\text{sig}} (\rightarrow D^* \ell^- \bar{\nu}_\ell)$

- Hadronic B_{tag} reconstruction: 2968 modes compared to 651 in *BABAR-09* $D\ell\bar{\nu}_\ell$ tagged analysis.

- Lower purity but **higher efficiency**.

- Skim-level tag-side requirement:
 $m_{\text{ES}} > 5.27 \text{ GeV}$, $|\Delta E| < 72 \text{ MeV}$.



- Similar to $\bar{B} \rightarrow D^* \tau^- \bar{\nu}_\tau$ *BABAR* paper, except no requirement on purity of tag-side modes, since our **signal-side** is very **clean**.
- Still cut-based 😞, unlike Belle II

The TreeFit to full topology


- Only $D^* \rightarrow D\pi_{\text{slow}}$ considered. $|\vec{p}_{\pi_{\text{slow}}}| < 400$ MeV in lab frame.
- $\Delta m \equiv (m_{D^*} - m_D)$ within 4σ of PDG expectation.
- Almost **no *uds*** continuum background for the charm system.
- **TreeFit** is critical to this analysis:
 - **Mass-constrain** the following: $\{B_{\text{tag}}, B_{\text{sig}}, D, D^*, \nu_{\text{miss}}\}$
 - $\Upsilon(4S)$ is **beam-spot** constrained. D^* **vertex**-constrained after B flight.
- Final **background** ($\sim 3\%$) very **small**. Estimated from generic $B\bar{B}$ MC and assigned as **systematic**.

The discriminating variable U

- Signal variable in B_{sig} rest frame for missing neutrino (from TreeFit w/o ν mass-constraint):

$$U = E_{\text{miss}} - |\vec{p}_{\text{miss}}| = E_{\nu} - |\vec{p}_{\nu}|$$


- Global comparison:



- Good comparisons found in other variables.
- $N_{\text{tot}} = 6112$, $N_{\text{bkgd}} \sim 180$

Heavy Quark Effective Theory for $\bar{B} \rightarrow D^*$

- w : relativistic γ factor of D^* in B RF
- At $\frac{\Lambda_{\text{QCD}}}{m_{b,c}} \rightarrow 0$ limit, only w matters.
- HQS: spin-flavor symm. among $\{b^\uparrow, b^\downarrow, c^\uparrow, c^\downarrow\}$
- Non-pert. QCD effects pushed into a single universal FF, $\zeta(w)$.
- In time scales $\ll \Lambda_{\text{QCD}}^{-1}$ Dirac structure in the weak current irrelevant



$$\frac{\langle D^*(v', \varepsilon) | V^\mu | \bar{B}(v) \rangle}{\sqrt{m_B m_{D^*}}} = i h_V(w) \varepsilon^{\mu\nu\alpha\beta} \varepsilon_\nu^* v'_\alpha v_\beta$$

$$A_1 = \frac{w+1}{2} r' h_{A_1}$$

$$\frac{\langle D^*(v', \varepsilon) | A^\mu | \bar{B}(v) \rangle}{\sqrt{m_B m_{D^*}}} = h_{A_1}(w)(w+1)\varepsilon^{*\mu} - h_{A_2}(w)(\varepsilon^* \cdot v)v^\mu - h_{A_3}(w)(\varepsilon^* \cdot v)v'^\mu$$

$$A_2 = \frac{r h_{A_2} + h_{A_3}}{r'} \equiv \frac{R_2 h_{A_1}}{r'}$$

$$V = \frac{h_V}{r'} \equiv \frac{R_1 h_{A_1}}{r'}$$

- HQS limit: $\{h_V, h_{A_1}, h_{A_3}\} \rightarrow \zeta(w)$ and $h_{A_2} \rightarrow 0$. $r' \equiv \frac{2\sqrt{m_{D^*} m_B}}{(m_{D^*} + m_B)}$

FF parameterization: BGL basis (Boyd *et al.*)

- Three versions: BGL95, BGL97 and BGL17.
- BGL FF's: $\{f_0, F_1, g\}$ related to conventional $\{A_1, A_2, V\}$ FF's and the helicity amplitudes $\{H_{\pm}, H_0\}$.

$$f_0 = (m_B + m_{D^*})A_1$$

$$F_1 = \frac{1}{2m_{D^*}} \left[(m_B^2 - q^2 - m_{D^*}^2)(m_B + m_{D^*})A_1 - \frac{4m_B^2 |\vec{\mathbf{k}}|^2}{(m_{D^*} + m_B)} A_2 \right]$$

$$= \sqrt{q^2} H_0$$

$$g = \frac{2V}{m_B + m_{D^*}}$$

$$H_{\pm} = \left(f_0 \mp m_B |\vec{\mathbf{k}}| g \right)$$

The BGL z -expansion

- z is a conformal map from $t \equiv q^2$ to unit circle.

$$z(t, t_0) = \frac{\sqrt{t_+ - t} - \sqrt{t_+ - t_0}}{\sqrt{t_+ - t} + \sqrt{t_+ - t_0}}$$

$$t_{\pm} = (m_B \pm m_{D^*})^2$$

- Choose t_0 so that z is small in the physical region $t \in [0, t_-]$

- Expansion of each FF from **dispersion** relations + **analyticity**

$$F_i(z) = \frac{1}{P_i(z)\phi_i(z)} \sum_{n=0}^N a_n^i z^n$$

- Blaschke-factor $P(z)$ from $B_c^{(*)}$ poles in non-physical $\sqrt{q^2} > m(BD^*)$ region.
- QCD ϕ_i outer functions.

- NB: $P_i(z)$ and $\phi_i(z)$ are non-trivial and carry significant part of the heavy-lifting in the q^2 shape. We follow Gambino'17.

FF parameterization: CLN [Caprini'97]

- CLN: similar expansion as BGL, but model-dependent **extrapolation** to $w \rightarrow 1$. Compact practical form with only **3 fit parameters**:


$$h_{A_1}(w) = h_{A_1}(1)[1 - 8\rho_{D^*}^2 z(w) + (53\rho_{D^*}^2 - 15)z(w)^2 - (231\rho_{D^*}^2 - 91)z(w)^3]$$

$$R_1(w) = R_1(1) - 0.12(w - 1) + 0.05(w - 1)^2$$

$$R_2(w) = R_2(1) + 0.11(w - 1) - 0.06(w - 1)^2$$

- **Curvature** and **slope** related for h_{A_1} . Too constrained?
- The expansions are derived from HQET+QCDSR, but experimentalists ignore the **uncertainties**.
- 20 years old. Calculations/uncertainties being carefully revisited by several groups (Gambino/Ligeti) now.

The $|V_{ub}|$ - $|V_{cb}|$ inclusive-exclusive tension saga



- $|V_{ub}|, |V_{cb}|$ tension: inclusive *persistently* $>$ exclusive at $> 3\sigma$.
- 2017 Grinstein/Gambino: fits to four 1-d projections (prelim. Belle)
- Claim: BGL form-factor parameterization resolves the $|V_{cb}|$ tension.
- Generated a lot of excitement...

BABAR BGL fits setup

- Linear BGL expansion till $N = 1$. ($N = 2$ terms not statistically significant but creates problems w/ unitarity)
- Two relations used to connect coefficients:
 - **Lattice** (MILC) constrains h_{A1} at zero-recoil and $f(q_{\max}^2) = 2\sqrt{m_B m_D^*} h_{A1}$
 - Also at zero-recoil, $F_1(q_{\max}^2) = (m_B - m_D^*) f(q_{\max}^2)$, so that $a_0^{F_1}$ is not an independent parameter.
- Slight isospin-dependence in above relations (negligible impact w/ current statistics)
- Unbinned non-extended ML fit using full **4-d rate** and complete *BABAR* data in $q^2 \in [0.2, 10.2]$ GeV². **No normalization.**

BGL fits setup – incorporating $|V_{cb}|$

- **Normalization** needed to get $|V_{cb}|$ obtained from **HFLAV**:

Mode	$\mathcal{B}(\bar{B} \rightarrow D^* \ell^- \bar{\nu}_\ell)(\%)$	τ_B (ps)	$\Gamma_{\text{tot}} \times 10^{15}$ (GeV)
B^0	4.88 ± 0.10	1.518 ± 0.004	21.16 ± 0.38
B^+	5.59 ± 0.21	1.638 ± 0.004	22.46 ± 0.79

- Add the $\Gamma_{\text{tot}} \equiv \int (d\Gamma/dw)dw = \mathcal{B}/\tau$ as Gaussian constraints.
- Nominal results quoted using the MILC result as a Gaussian constraint:

$$h_{A_1}(1)|_{\text{MILC}} = 0.906 \pm 0.013$$

Unitarity, stability checks and global minima

- Six fit variables: $\{a_0^f, a_1^f, a_1^{F1}, a_0^g, a_1^g, |V_{cb}|\}$. *BABAR*-only version (without $|V_{cb}|$) also provided.
- BGL setup is even more complicated than CLN. Significant effort to ensure **stability** and robustness of the **global minimum**.
- **1000** fit iterations with randomized start values within $[-1, 1]$ for the BGL coefficients.
- Fit parameters scaled appropriately such that MINOS cov. matrix diag. elements are all ~ 1 . HESSE and MINOS cov. matrices matches.

Systematics and final results

- Most **systematics cancel** since *BABAR* part uses **no normalization**.
- Many cross-checks performed with isospin-separated (different slow pions), e/μ separated fits. Primary remnant source is the background.
- Final $N = 1$ BGL expansion results, including systematic uncertainties:


$a_0^f \times 10^2$	$a_1^f \times 10^2$	$a_1^{F1} \times 10^2$	$a_0^g \times 10^2$	$a_1^g \times 10^2$	$ V_{cb} \times 10^3$
1.29 ± 0.03	1.63 ± 1.00	0.03 ± 0.11	2.74 ± 0.11	8.33 ± 6.67	38.36 ± 0.90

- Final CLN results:

$\rho_{D^*}^2$	$R_1(1)$	$R_2(1)$	$ V_{cb} \times 10^3$
0.96 ± 0.08	1.29 ± 0.04	0.99 ± 0.04	38.40 ± 0.84


- Our BGL $|V_{cb}|$ is **consistent** with WA $|V_{cb}|_{\text{excl.}}$ and remains in **tension** with WA $|V_{cb}|_{\text{incl.}}$.

BABAR results – FF shapes



- Dashed curves denote 1σ error bands
- *BABAR* FF's significantly different from CLN-WA. CLN p -value: 0.0017
- Within *BABAR*, CLN/BGL seem consistent.
- LCSR'08 at $q^2 = 0$ (large recoil) has huge errors, but central value more consistent with *BABAR*.

BABAR results – $R_{1,2}$ ratios



- Included the $R_{1,2}$ curves (w/o errors) from the original Caprini paper.

- BABAR BGL: $R_1(1) \approx 1.2$ and the slope is positive; $R_2 \approx 1$ and flat.
- Consistent with HQET, although no HQET constraints used in the fit. It's what the data prefers.
- Note: CLN $R_{1,2}$ shape is fixed. Too constrained?

BABAR-BGL semi-tauonic predictions

- We mostly follow the Gambino'17 prescriptions with $\sim 15\%$ uncertainty on the **HQET** estimation of $P_1(1)$ (scalar FF).
- At maximum recoil, instead of LCSR, we use the *BABAR*-BGL prediction from $F_1(w_{\max})$.

$R(D^*)$ Description	Value
HFLAV'19 WA exp.	0.295 ± 0.014
Fajfer'12 CLN	0.252 ± 0.003
Ligeti'17 BGL	0.257 ± 0.003
Gambino'17 BGL	0.260 ± 0.008
HFLAV'19 SM avg.	0.258 ± 0.005
BABAR BGL	0.253 ± 0.005

- $\sim 2.8\sigma$ dev. in $R(D^*)$ w/ *BABAR*-BGL.
- Predictions stable, but disagreement on the **errors**.
- Effect of the **FF's** on exp. **efficiencies**?

- P_τ : -0.483 ± 0.027 (*BABAR*-BGL), $-0.38 \pm 0.51_{-0.16}^{+0.21}$ (*Belle*'17)
- $F_L^{D^*}$: $\underbrace{+0.454 \pm 0.011}_{\text{predictions}}$ (*BABAR*-BGL), $\underbrace{+0.60 \pm 0.08 \pm 0.035}_{\text{measurements}}$ (*Belle*'19)

Summary

- First **tagged 4d** $\bar{B} \rightarrow D^* \ell \bar{\nu}_\ell$ angular analysis using *BABAR* data
- *BABAR* **FF**'s w/ BGL/CLN consistent and deviate with **CLN-WA**.
- *BABAR* + HFLAV-BF: $|V_{cb}|$ consistent between BGL and CLN-WA.
 $\sim 3\sigma$ exclusive-inclusive **tension persists**.
- Need **lattice** $w > 1$ data asap for joint fit with unbinned *BABAR*! Short PRL submitted. Long PRD under preparation.
- Looking forward to Belle *tagged 4d fit* results.

Summary

- First **tagged 4d** $\bar{B} \rightarrow D^* \ell \bar{\nu}_\ell$ angular analysis using *BABAR* data
- *BABAR* FF's w/ BGL/CLN consistent and deviate with **CLN-WA**.
- *BABAR* + HFLAV-BF: $|V_{cb}|$ consistent between BGL and CLN-WA.
 $\sim 3\sigma$ exclusive-inclusive **tension persists**.
- Need **lattice** $w > 1$ data asap for joint fit with unbinned *BABAR*! Short PRL submitted. Long PRD under preparation.
- Looking forward to Belle *tagged 4d fit* results.

Thank you!

Backup

References

- Gambino'17: arXiv:1703.06124, arXiv:1707.09509
- BGL: hep-ph/9508211, hep-ph/9705252, arXiv:1703.08170
- CLN: arXiv:9712417
- Belle τ/D^* pol.: arXiv:1709.00129, arXiv:1901.06380
- TreeFit: arXiv:physics/0503191

The $b \rightarrow c W_L^{*-} (\rightarrow \ell^- \bar{\nu}_\ell)$ effective Hamiltonian

- In the SM, W_L^{*-} acts a left-handed off-shell **spin-1** vector boson.
- The minimal effective Hamiltonian for light leptons is:

$$\mathcal{H}_{\text{eff}} = \frac{2G_F V_{cb}^L}{\sqrt{2}} [(g^V \bar{c} \gamma_\mu b - g^A \bar{c} \gamma_\mu \gamma_5 b) \bar{\ell} \gamma^\mu \nu_L + (\dots)]$$

- Additional scalar (charged H^\pm) and tensor (Leptoquarks) operators can occur.
- **Heavy W_R^-** can induce $g_{V,A} \equiv (1 \pm \epsilon_R)$ even for **light leptons**.
- The helicity amplitudes $\{H_\pm, H_0\}$ can acquire phases from **complex ϵ_R** .

NP searches: the $\sin \chi$ terms [arXiv:1505.02873]

- In SM, the amplitudes are relatively real. $\sin \chi$ terms are zero.
- Non-zero $\Gamma_{\{6,9,10,14\}}$ would be clear sign of NP!
- Formalism developed at *BABAR* for both the ρ and D^* vector mesons.
- $\pi\pi$ S -wave under the ρ also very interesting for RH current searches in $b \rightarrow u$.

i	$f_i(\Omega)$	$\Gamma_i^{\text{tr}}(q^2)/(\mathbf{k}q^2)$
1	$P_0^0 Y_0^0$	$h_0^2 + h_{\parallel}^2 + h_{\perp}^2$
2	$P_2^0 Y_0^0$	$-\frac{1}{\sqrt{5}}(h_{\parallel}^2 + h_{\perp}^2) + \frac{2}{\sqrt{5}}h_0^2$
3	$P_0^0 Y_2^0$	$\frac{1}{2\sqrt{5}} h_{\parallel}^2 + h_{\perp}^2 - 2h_0^2 $
4	$P_2^0 Y_2^0$	$-\frac{1}{10}(h_{\parallel}^2 + h_{\perp}^2) - \frac{2}{5}h_0^2$
5	$P_2^1 \sqrt{2} \text{Re}(Y_2^1)$	$-\frac{3}{5}h_{\parallel} h_0 \cos \delta_{\parallel}$
6	$P_2^1 \sqrt{2} \text{Im}(Y_2^1)$	$\frac{3}{5}h_{\perp} h_0 \sin \delta_{\perp}$
7	$P_0^0 \sqrt{2} \text{Re}(Y_2^2)$	$-\frac{3}{2\sqrt{15}}(h_{\parallel}^2 - h_{\perp}^2)$
8	$P_2^0 \sqrt{2} \text{Re}(Y_2^2)$	$\frac{\sqrt{3}}{10}(h_{\parallel}^2 - h_{\perp}^2)$
9	$P_0^0 \sqrt{2} \text{Im}(Y_2^2)$	$\sqrt{\frac{3}{5}}h_{\perp} h_{\parallel} \sin(\delta_{\perp} - \delta_{\parallel})$
10	$P_2^0 \sqrt{2} \text{Im}(Y_2^2)$	$-\frac{\sqrt{3}}{5}h_{\perp} h_{\parallel} \sin(\delta_{\perp} - \delta_{\parallel})$
11	$P_0^0 Y_1^0$	$-\sqrt{3}h_{\perp} h_{\parallel} \cos(\delta_{\perp} - \delta_{\parallel})$
12	$P_2^0 Y_1^0$	$\frac{3}{\sqrt{15}}h_{\perp} h_{\parallel} \cos(\delta_{\perp} - \delta_{\parallel})$
13	$P_2^1 \sqrt{2} \text{Re}(Y_1^1)$	$\frac{3}{\sqrt{5}}h_{\perp} h_0 \cos \delta_{\perp}$
14	$P_2^1 \sqrt{2} \text{Im}(Y_1^1)$	$-\frac{3}{\sqrt{5}}h_{\parallel} h_0 \sin \delta_{\parallel}$

4-d rate expression

- Generic amplitude with complex \mathcal{H}_i 's for $\bar{B} \rightarrow D^*(\rightarrow D\pi)\ell^-\bar{\nu}_\ell$:

$$|\overline{\mathcal{M}}|^2 = \left| \sum_{\lambda \in \{0, \pm 1\}} \sqrt{3} \mathcal{H}_\lambda d_{\lambda,0}(\theta_V) d_{\lambda,-1}^1(\theta_l) e^{i\lambda\chi} \right|^2$$

- Expanding this out, leads to:

$$\frac{d\Gamma}{dq^2 d\Omega} = \frac{\sqrt{8\pi} |V_{cb}|^2 \eta_{EW}^2 G_F^2 \mathcal{B}^{D^* \rightarrow D\pi}}{3m_B^2 (4\pi)^4} \left\{ \sum_{i=1}^{14} f_i(\Omega) \Gamma_i(q^2) \right\}$$

- If the \mathcal{H}_i 's are real, it boils down to the usual expression:

$$\begin{aligned} \frac{d\Gamma}{dq^2 d\Omega} = & \left[\left(H_+^2 (1 - \cos \theta_l)^2 + H_-^2 (1 + \cos \theta_l)^2 \right) \sin^2 \theta_V + \right. \\ & 2H_0 \sin \theta_l \sin 2\theta_V \cos \chi \left(H_+ (1 - \cos \theta_l) - H_- (1 + \cos \theta_l) \right) \\ & \left. + 4H_0^2 \sin^2 \theta_l \cos^2 \theta_V + 2H_+ H_- \sin^2 \theta_l \sin^2 \theta_V \cos 2\chi \right] \\ & \times \frac{3}{8(4\pi)^4} G_F^2 \eta_{EW}^2 |V_{cb}|^2 \frac{kq^2}{m_B^2} \mathcal{B}(D^* \rightarrow D\pi) \end{aligned}$$

BSemiExclAdd skim


- Charmful seed $S \in \{D^{(*)0}, D^{(*)+}, D_s^{(*)+}, J/\psi\}$ and charmless $Y = n_1\pi^\pm + n_2K^\pm + n_3K_S^0 + n_4\pi^0$,
- Hadronic B_{tag} reconstruction: 2968 modes compared to 651 in [BABAR-09](#) $D\ell\bar{\nu}_\ell$ tagged analysis.
- Charmful seed $S \in \{D^{(*)0}, D^{(*)+}, D_s^{(*)+}, J/\psi\}$ and charmless $Y = n_1\pi^\pm + n_2K^\pm + n_3K_S^0 + n_4\pi^0$,

- Purity cut removed.
- In the e^+e^- rest frame:

$$\Delta E = E_{\text{tag}} - \sqrt{s}/2$$

$$m_{\text{ES}} = \sqrt{s/4 - |\vec{p}_{\text{tag}}|^2}$$

- Skim-level tag-side requirement:
 $m_{\text{ES}} > 5.27$ GeV, $|\Delta E| < 72$ MeV.



Signal side reconstruction and MC samples

- Full exclusive event topology reconstructed. No additional charged tracks. Final *BABAR* ultimate PID selectors for $\{e, \mu, \pi, K, \pi^0\}$
- **Leptons**: $|\vec{p}_{\text{lab}}| > 200$ and 300 MeV for e and μ . $\theta_{\text{lab}}^\ell \in (0.4, 2.6)$ fiducial cut. Brem-recovery for e .
- For a given event all $\{D^0, D^\pm, D^{*0}, D^{*\pm}, +\text{charmless}\}\ell$ combinations sought in **disjoint** channels. Helps cleaning up the “specific” **SL** backgrounds.
- MC samples:
 - **Generic $B\bar{B}$** MC: FF's and BF's reweighted to PDG. Used for **background** studies/estimation.
 - **FLATQ2** MC: tag-side same as in generic MC. Signal-side generated flat in $dq^2 d\Omega$ for **acceptance** calculation in the angular analysis.

The 12 individual $D^*(\rightarrow D\pi)\ell$ modes

- E_{extra} : additional good photons not used in the reconstruction.
- Only 3 cleanest D^0 modes used. Cuts on E_{extra} and CL from TreeFit.

Particle	D^0 Decay mode	Mode #	Cuts
$D^{*0} \rightarrow D^0\pi^0$	$K^-\pi^+$	1	$E_{\text{extra}} < 0.5 \text{ GeV}$, CL > 0.001
	$K^-\pi^+\pi^0$	2	$E_{\text{extra}} < 0.4 \text{ GeV}$, CL > 0.01
	$K^-\pi^+\pi^-\pi^+$	3	$E_{\text{extra}} < 0.4 \text{ GeV}$, CL > 0.01
$D^{*+} \rightarrow D^0\pi^+$	$K^-\pi^+$	4	$E_{\text{extra}} < 0.6 \text{ GeV}$, CL $> 10^{-6}$
	$K^-\pi^+\pi^0$	5	$E_{\text{extra}} < 0.6 \text{ GeV}$, CL $> 10^{-6}$
	$K^-\pi^+\pi^-\pi^+$	6	$E_{\text{extra}} < 0.6 \text{ GeV}$, CL $> 10^{-6}$

- 6 D^* modes \Rightarrow 12 $D^*\ell$ modes.
- **Truth-matching** criterion: reconstructed and generated mode must match.
- Each mode has different **acceptance/backgrounds**. Analysed **independently** and combined at the end.
- Quite different from *BABAR* $R(D^{(*)})$ analysis (*many* modes lumped together)

Final mode-wise statistics

- Due to low tagging efficiency, limited MC statistics is critical.

$[D]\pi e^\pm$ mode	N_{tot}	N_{bkgd}	N_{sim}
$[K^-\pi^+]\pi^0 e^\pm$	486	15	14396
$[K^-\pi^+\pi^0]\pi^0 e^\pm$	547	19	17806
$[K^-\pi^+\pi^-\pi^+]\pi^0 e^\pm$	350	13	3410
$[K^-\pi^+]\pi^+ e^\pm$	418	5	3990
$[K^-\pi^+\pi^0]\pi^+ e^\pm$	801	14	7795
$[K^-\pi^+\pi^-\pi^+]\pi^+ e^\pm$	453	8	4472

$[D]\pi\mu^\pm$ mode	N_{tot}	N_{bkgd}	N_{acc}
$[K^-\pi^+]\pi^0\mu^\pm$	442	18	14893
$[K^-\pi^+\pi^0]\pi^0\mu^\pm$	574	29	18441
$[K^-\pi^+\pi^-\pi^+]\pi^0\mu^\pm$	395	16	3577
$[K^-\pi^+]\pi^+\mu^\pm$	417	10	4171
$[K^-\pi^+\pi^0]\pi^+\mu^\pm$	768	22	7838
$[K^-\pi^+\pi^-\pi^+]\pi^+\mu^\pm$	461	11	4446

- Final cut: $|U| < 90$ MeV.
- $N_{\text{tot}} = 6112$, $N_{\text{bkgd}} \sim 180$ over all 12 modes.
- Clean enough that background is a *systematic* from generic $B\bar{B}$ MC.
- No additional signal-background separation required.

“Model-independent” FF fits: pseudoscalars

- **Pseudoscalar** cases: *z*-**expansion** fits for the single $f_+(q^2)$ FF from both w/ and w/o hadronic tagging at *B*-factories.
- Rate $\propto \sin^2 \theta_\ell |f_+^2(q^2)|$ factorizes \Rightarrow effectively **1d** analysis in q^2 .
- Reliable theory inputs also exist:
 - $\bar{B} \rightarrow \pi \ell^- \bar{\nu}_\ell$: both **LCSR** (large-recoil) and **Lattice** (low-recoil)
 - $\bar{B} \rightarrow D \ell^- \bar{\nu}_\ell$: non-zero recoil **Lattice** data from both MILC ([1503.07237](#)) and HPQCD ([1505.03925](#))
- Latest Belle hadronic tagged results: $\bar{B} \rightarrow \pi \ell^- \bar{\nu}_\ell$ ([1306.2781](#)) and $\bar{B} \rightarrow D \ell^- \bar{\nu}_\ell$ ([1510.03657](#)).

“Model-independent” FF fits: vector mesons

- **Vector meson** cases much more difficult than pseudoscalars:
 - **Three FF's** that require a full **4-d** amplitude analysis to disentangle.
 - Charmless $\{\rho, \omega\}$: broad resonances and difficult for both LCSR and Lattice. ρ can have significant $\pi\pi$ S -wave. Plus ρ - ω mixing.
 - Charmful D^* : published **Lattice** data only at $w = 1$ yet. LCSR at large recoil unreliable for heavy D^* . **HQET** framework helps, though.
- Long time goal at *BABAR*. But **hard**, analysis-wise as well as little theory guidance on the QCD input in the z -expansion (till pre-2017). BGL papers for $B \rightarrow D^*$ were pre-2000.
- **Grinstein** $B \rightarrow D^*$ paper in 2017 reinvigorated the discussion. Much follow-up theory activity by Gambino/Ligeti/Lattice \Rightarrow very helpful!



BGL: optimal t_0 and unitarity conditions

- Nominal choice (CLN):

$$t_0 = t_-$$

- Optimal choice:

$$t_0^{\text{opt}} = t_+ - \sqrt{t_+(t_+ - t_-)}$$




- t_0^{opt} approximately halves the $|z|$ range, for better convergence.
- The BGL coefficients must satisfy the unitarity constraints:

$$\sum_{n=0}^N |a_n^f|^2 + |a_n^{F_1}|^2 \equiv S_{1+} \leq 1, \quad \sum_{n=0}^N |a_n^g|^2 \equiv S_{1-} \leq 1$$

- In reality, far from saturation: $S \sim \mathcal{O}(10^{-2})$.

$|V_{cb}|$ tension and BGL fits: ICHEP'18


- ICHEP'18 plenary claim:



- But life is not so simple...

HQET implications from these BGL fits

- Ab initio, *huge* HQET breaking from BGL fits:






- HQS: FF ratios $R_{1,2} \rightarrow 1$.
- R_1 from Gambino looks quite weird
- LCSR+unitarity usage!

- HQET works pretty well in heavy quark spectroscopy, lifetimes etc. So, quite surprising, if true...

Preliminary non-zero recoil Lattice data



- Preliminary lattice data seems consistent with $h_{A_1}, h_V, R_1 \sim \mathcal{O}(1)$ expectation from HQET. Lattice uncertainties on these $\sim 3\%$.





- $R_1(w)$ flat-ish and $R_1(1) \sim 1.3$
- MILC/JLQCD have different systematics and ensembles. Need both for confirmation.

- R_2 uncertainties appear to be large in Lattice.

Preliminary non-zero recoil Lattice data (cntd.)




- HQET: $h_{A_2} \rightarrow 0$
- Error: $\sim 40\%$



- HQET: $h_{A_3} \rightarrow 1$
- Error: $\sim 30\%$

BABAR BGL results – FF shapes (cntd.)



- New LCSR data at large **negative** q^2 values from [1811.00983](#)
- **BABAR BGL** “not inconsistent” with these LCSR data. NB: *BABAR* fits did not include any LCSR/HQET inputs.
- **Curvature** of A_1 from CLN-WA seems to change at $q^2 \rightarrow -15 \text{ GeV}^2$.

Ambiguities in 'product of four 1-d' fits?

- **Angle** distributions: barred terms, q^2 -info **integrated** away. Very little sensitivity to q^2 -dependence.

$$\frac{d\Gamma}{d\chi} \sim \left[\left(|H_+|^2 + |H_-|^2 + |H_0|^2 \right) - |H_+ H_-| \cos 2\chi \right]$$

$$\frac{d\Gamma}{d \cos \theta_\ell} \sim \left[(1 - \cos \theta_\ell)^2 |H_+|^2 + (1 + \cos \theta_\ell)^2 |H_-|^2 + 2 \sin^2 \theta_\ell |H_0|^2 \right]$$

$$\frac{d\Gamma}{d \cos \theta_V} \sim \left[\sin^2 \theta_V \left(|H_+|^2 + |H_-|^2 \right) + 2 \cos^2 \theta_V |H_0|^2 \right]$$





- $d\Gamma/dw$ has info only on sum of **FF squares**:

$$\frac{d\Gamma}{dw} \sim \sqrt{w^2 - 1} (1 - 2wr + r^2) \left[|H_+|^2 + |H_-|^2 + |H_0|^2 \right]$$

- Can the **individual FF's** be extracted **unambiguously** in a fit to product of four 1-d projections?
- **BABAR-BGL'19** instead employs a full **4-d** fit.

Comparisons w/ Belle-17 data

Overlay, *not* a fit:




- Attempted to add unpublished Belle'17 data as a Gaussian constraint to the overall *BABAR* NLL.
- Fit significantly deviating from *BABAR*-only if entire 40×40 cov. matrix used. Single set of 10×10 cov. matrix ok \Rightarrow not used in the end.
- Note: only mild diff. in 1-d projections between CLN-WA and *BABAR*-BGL.


Data/MC comparisons: *BABAR*

- FLATQ2 accepted MC (incl. eff. effects) weighted by the BGL results should **match** the data in all **multi-dim.** distributions.

1-d distributions:



3-d in angles, q^2 -integrated:



Lattice'19 MILC slide

