



Contribution ID: 580

Type: Parallel talk

## Towards two-loop computations in four dimensions with the Loop-Tree Duality

Thursday, 11 July 2019 17:30 (20 minutes)

We propose a new algorithm based on the Loop-Tree Duality theorem to renormalise and calculate two-loop diagrams. The ultraviolet singularities are locally cancelled in a systematic way and at the integrand level, allowing for a full four-dimensional numerical implementation of the method. In particular, we apply the method to calculate the  $H \rightarrow \gamma\gamma$  amplitude at two-loop level, and find an excellent agreement with already available literature results. We also present other advantages of the Loop-Tree Duality formalism, such as the possibility to write unintegrated amplitudes in a universal way, regardless of the particle running inside the loop.

**Primary author:** DRIENCOURT-MANGIN, Félix (IFIC, UV-CSIC)

**Co-authors:** RODRIGO, German (IFIC CSIC-UV); SBORLINI, German (Università di Milano, INFN Milano and IFIC-Valencia); TORRES BOBADILLA, William Javier (IFIC CSIC-UV)

**Presenter:** DRIENCOURT-MANGIN, Félix (IFIC, UV-CSIC)

**Session Classification:** Quantum Field and String Theory

**Track Classification:** Quantum Field and String Theory