

Vetoing Tops without B -Tagging¹

EPS 2019 - Ghent University

Richard Ruiz

Center for Cosmology, Particle Physics, and Phenomenology (CP3)
Universite Catholique de Louvain



12 July 2019



¹w/ folks from Durham+Pittsburgh and NIKHEF+Paris

Motivation

Inclusive production rate of top quarks at hadron colliders is, well, **big**...





- The LHC generates $L \sim 1 \text{ fb}^{-1}/\text{day} \implies 1\text{M } t\bar{t}$ pairs/day
 - ▶ Single- / pair-production background to **all** measurements and searches
- **Subtle bit**: standard tools, i.e., b -tagging, are not perfect
 - ▶ $\varepsilon^b \sim 0.7 \implies 10\%$ top pairs survive cull
 - ▶ Actually $> 10\%$ since b -tagging requires: $|\eta^j| \lesssim 2.5$, $p_T^j > 25 - 30$ GeV
- **Question**: Can such “irreducible backgrounds” be further reduced?

Revisiting an Old Idea: (Central) Jet Vetoes

In color-singlet processes, like weak boson scattering:

Absence of central color flow

⇒ absence of central, high- p_T jets (“rapidity gap”²)



Basis for Central Jet Veto³:

- Reject events with any jet satisfying $p_T^j > 25 - 30$ GeV, $|\eta^j| < 2 - 3$
- Crucial to Higgs physics but not perfect...

²Dokshitzer, Khoze, Troyan ('86)

³Barger, et al, PRD ('91) + PLB ('95); Bjorken, PRD ('94)

Limitations of Jet Vetoes

- 1 Almost uniquely, top are a background *all*⁴ measurements + searches
 - ▶ heavy quark flavor- / *b*-vetoing is very good but not perfect
- 2 Tight/stringent, central jet vetoes can alleviate this, but at cost of sensitivity to missing higher order corrections⁵
 - ▶ Substantial pQCD theory uncertainty despite high precision
- 3 Low signal efficiency for colorless, high-mass BSM⁶
 - ▶ At *pp* colliders, even the most exotic, colorless BSM is accompanied by high- p_T , initial-state radiation

⁴Literally, just listen to any ATLAS or CMS person about their analysis

⁵E.g., the pQCD/hep-ph literature on jet vetoes on [[InSpireHep](#)]

⁶For example: Sleptons by Tackmann, et al, [[1603.03052](#)]; SSM W'/Z' by Fuks, RR [[1701.05263](#)]; and heavy neutrinos by Pascoli, RR, et al [[1805.09335](#), [1812.08750](#)]

Reenvisioning⁷ the Jet Veto:


Event-Based⁸ Jet Vetoes

⁷Pascoli, **RR**, et al [PLB ('18), [1805.09335](#)]

⁸aka: "safe jet veto," "dynamic jet veto," or "phase space-dependent jet veto"


Leptons in Vector Boson Fusion

For production of leptons in VBF/VBS: $p_T^{\ell_k}, S_T \sim M_{VV} \gg p_T^{jVBS} \sim M_V/2$



Leptons in Vector Boson Fusion


For production of leptons in VBF/VBS: $p_T^{\ell_k}, S_T \sim M_{VV} \gg p_T^{jVBS} \sim M_V/2$



An idea: on event-by-event basis, set $p_T^{\text{Veto}} = p_T^{\ell_1}$

- VBF events pass by construction.
- What about other color-singlet processes, e.g., high-mass Drell-Yan?

Top Quark Background vs Event-Based Jet Vetoes



$pp \rightarrow t\bar{t}Z \rightarrow 1\mu + 3e + 2j_b + \cancel{E}_T$
candidate event [1509.05276]

Textbook kinematics:


- $m_{ee} = 93$ GeV
- $\cancel{E}_T = 57$ GeV

Typically,

- $p_T^{e3} \sim \frac{M_Z}{2} \sim 45$ GeV
 - $p_T^{e1} \sim \frac{m_t}{4} \left(1 + \frac{M_W^2}{m_t^2}\right) \sim 50$ GeV
 - $p_T^{b1} \sim \frac{m_t}{2} \left(1 - \frac{M_W^2}{m_t^2}\right) \sim 60$ GeV
- $p_T^{b1} > p_T^{\ell k} \implies$ event vetoed!

Setting $p_T^{\text{Veto}} = p_T^{\ell_1}$ can **reduce** top background **without** b -jet tagging!
- Di-/triboson backgrounds nontrivially impacted, too! [Backup slides]

Case Study I: sleptons and generalizing event-based jet veto definitions⁹




⁹Fuks, Nordstrom, RR, Williamson [1901.09937]

Leptons vs Hadrons

Consider the signature $pp \rightarrow \mu^+ \mu^- + X$ at NLO+PS(LL)


- Generator cuts + $p_T^\ell > 15$ GeV + $|\eta^\ell| < 2.4$ + anti- k_T ($R = 1$)

Plotted: $r_{j_1}^{\ell_1} = p_T^{\ell_1} / p_T^{j_1}$



Good (poor) S/B separation for $|m_{\tilde{\mu}} - m_{\tilde{\chi}}| / m_{\tilde{\mu}} \gg 0$ (~ 0)

Plotted: $r_{H_T}^{\ell_2} = p_T^{\ell_2}/H_T$, $H_T = \sum_{k \in \{\text{clusters}\}} E_T^k$



Choices of leptonic and hadronic observables is param. space-dependent
 Nevertheless, separation between $t\bar{t}$ from $V^{(*)}V^{(*)}$ is clear. :)

As a benchmark, we considered CMS's Run II search [1806.05264]

- Changes: relaxed leading $p_T^{\ell_1}$ cut and swapped out $p_T^{\text{Veto}} = 25$ GeV
- + FxFx1j merging, MPI/UE tuning


Plot: Sensitivity change using $H_T^{\text{Veto}} = p_T^{\ell_2}$ in $(m_{\tilde{\chi}}, m_{\tilde{\mu}})$ -space

darker = improvement

- Improved sensitivity
for larger mass
splittings

- Worse for small
mass splitting

- Other ratios show
opposite behavior
[1901.09937]



Summary, Conclusions, and Outlook

A class of jet vetoes has been investigated for high-mass, multi- ℓ searches, one based on comparing lepton vs hadronic activities, e.g., $p_T^{\text{Veto}} = S_T$


- 1 High signal efficiency ✓
 - ▶ New scheme reveals $> 90 - 95\%$ signal acceptance with little-to-no dependence on mass scales [1805.09335]
- 2 High background rejection ✓
 - ▶ Redesigned analysis gives better reduction of background
- 3 Low/less sensitivity to missing higher order corrections ✓
 - ▶ Substantial reduction in QCD theory uncertainty
 \implies less need for high-precision resummation [1812.08750]
- 4 Universality ✗
 - ▶ Collider signature and param space dependence [1901.09937]
- 5 Analytic Control ✗
 - ▶ Resummation beyond parton shower-precision less clear
- 6 Broad applicability to BSM looks promising (see backup slides)



Thank you.

Backup I:

Case Study II: Heavy Neutrinos at the LHC¹⁰




¹⁰Pascoli, RR, et al [PLB ('18), [1805.09335](#), JHEP ('19), [1812.08750](#)]

Jet vetoes tries to discriminate jet activity in signal and background.

Consider $pp \rightarrow N\ell \rightarrow 3e/\mu + X$ at NLO+PS(LL)+anti- k_T ($R = 1$)


Plotted: $p_T^{j_1}$ for CC DY signal + SM Bkg



Separation according to **Born color** clear but more overlap for larger m_N

Event-based jet vetoes discriminate against hadronic and leptonic activities


Plotted: $r_{j_1}^{\ell_1} = p_T^{\ell_1} / p_T^{j_1}$ for CC DY signal + SM Bkg



Similar behavior for other ratios, e.g., S_T/H_T , $p_T^{\ell_k}/H_T$

- Not universal, e.g. degenerate mass limit in $\tilde{\mu} \rightarrow \mu \tilde{\chi}^0$ [[1901.09937](#)]

Plot: LHC 14 sensitivity to **active-sterile neutrino mixing** (coupling²) vs **heavy N mass** (m_N), in search for $pp \rightarrow \mu^\pm e^\mp \ell_X + X$ ($\ell_X = e, \mu, \tau_h$)¹¹



Improved sensitivity up to 10 – 11 \times with $\mathcal{L} = 3 \text{ ab}^{-1}$.

¹¹See [1812.08750] for various lepton flavor permutations, uncertainty plots, etc


Backup II:

Di- and Tribosons vs Event-Based Jet Vetoes

Diboson and Triboson Processes


Di- / triboson processes contain large amounts of hadronic activity

- Significant cancelations in Born-like configurations¹²
- Nontrivial color flow, despite naïve, color-singlet nature



¹²i.e., Radiation zeros (super interesting!) Mikaelian ('78); +Brown, et al ('78-'79); Zhu ('80); Brodsky, et al ('82,'83); Baur, Han, Ohnemus [PRL ('94), [hep-ph/9403248](https://arxiv.org/abs/hep-ph/9403248)]

Irreducible Bkgs vs Event-Based Jet Vetoes



Inclusive diboson production driven by $pp \rightarrow VV'j + X$

- Manifests as large K -factor in $pp \rightarrow W^+W^-$ at NNLO [1408.5243]
- $p_T^\ell \sim \frac{M_V}{2} + \frac{p_T^j}{2n_b}$ ($n_b = \#$ bosons)
- \Rightarrow large $p_T^j > p_T^\ell$ tail

Jets mistagged/IDed as τ_h/e , i.e., “fake leptons,” major irreducible bkg:

- E.g., $pp \rightarrow W^+W^-j$
- Subleading jets more likely to be mis-IDed than leading jet

Setting $p_T^{\text{Veto}} = p_T^{\ell_1}$ can **reduce** irreducible backgrounds