


Sterile Neutrino searches with the ICARUS detector

*Alessandro Menegolli
University and INFN Pavia*

*on behalf of the
ICARUS collaboration*

*European Physical Society
Conference on High
Energy Physics*

*Ghent (Belgium)
July 10-17, 2019*



The full list of the Collaboration: <https://icarus.fnal.gov/collaboration>



Catania (INFN and Univ.)
GSSI
LNGS
INFN Milano Bicocca
INFN Napoli
Padova (INFN and Univ.)
Pavia (INFN and Univ.)



CINVESTAV



Brookhaven (BNL)
Colorado State
FNAL
Houston
Pittsburgh
Rochester
SLAC
Southern Methodist Univ.
Texas (Arlington)



Spokesperson: C. Rubbia, INFN GSSI
more than 90 collaboration members

International Partner





Neutrino Platform NP01

Many thanks for the major contributions to the Far Detector cryogenics and cosmic ray tagger from our partners at CERN, INFN-Bologna, INFN-Lecce, INFN-Milano, INFN-Napoli, INFN-Genoa, INFN-LNS.

Liquid Argon TPC: an “electronic bubble chamber”


- LAr-TPCs: ideal detectors for neutrino physics and nucleon decay:
 - 3D reconstruction with high (mm^3) spatial granularity.
 - Homogeneous, full-sampling calorimetry for contained particles.
 - Scintillation light can provide fast signals for timing/triggering.
 - Electrons can drift for several meters (if Argon is sufficiently pure).
 - LAr is dense and cheap: very large masses (ktons) are realistic.
- First proposed by C. Rubbia in 1977: long R&D at INFN and CERN culminated in first large-scale experiment: **ICARUS-T600** at LNGS (2010-2013):



- ICARUS was exposed to CNGS beam and cosmics for 3 years, confirming expected performance and obtained important physics results.
- It proved the maturity of the LAr-TPC technique for large-scale experiments (DUNE).


- High electron lifetime: > 7 ms (impurity concentration < 40 ppt) over whole run. Crucial step towards future larger detectors

2014 JINST 9 P12006




- Excellent spatial/calorimetric reconstruction. Accurate dE/dx measurement with fine sampling ($0.02X_0$). Particle ID from dE/dx vs. range

AHEP (2013) 260820





- Momentum of escaping muons measured by multiple Coulomb scattering. Average $\sim 15\%$ resolution on stopping muons ($0.5 \div 5$ GeV/c)


JINST 12P04010




- ν_e CC event (electron-initiated EM showers) separation from NC background with π^0 (γ -initiated showers): crucial for oscillation physics.
- LAr-TPC provides 3 handles:
 - Visual identification of γ conversion gap.
 - Reconstruction of π^0 invariant mass.
 - dE/dx: calorimetric accuracy and fine sampling (2% X_0) allow measuring dE/dx on each wire: single MIP corresponds to an electron.



High-energy CNGS ν_e CC interaction:



- ICARUS searched for sterile ν oscillations through ν_e appearance in the CNGS beam.
- $L/E \sim 36 \text{ km/GeV}$, far from LSND value $\sim 1 \text{ km/GeV}$ \rightarrow "sterile-like" oscillation was averaged out, canceling energy dependence.
- $7.9 \cdot 10^{19}$ pots analyzed (~ 2650 ν interactions).
- Expected $\sim 8.5 \pm 1.1$ ν_e background events in absence of anomaly, mostly from intrinsic ν_e beam contamination.
- Estimated ν_e identification efficiency $\sim 74\%$ with negligible background from misidentification.
- 7 events observed \rightarrow no evidence of oscillation.
- Most of LSND allowed region is excluded - except for small area around $\sin^2 2\theta \sim 0.005$, $\Delta m^2 < 1 \text{ eV}^2$
- Similar result by OPERA with same CNGS beam and different detection technique.





Eur. Phys. J. C
(2013) 73:2599

The SBN project

8

$$L/E_\nu \sim 600 \text{ m} / 700 \text{ MeV} \sim \mathcal{O}(1 \text{ m/MeV})$$

**T600 also off-axis on NUMI beam:
Asset for DUNE**



ICARUS T600


FAR DETECTOR:
T600 – 476 ton

ICARUS


MicroBooNE
89 ton

NEAR DETECTOR:
SBND – 112 ton

- The experiment is expected to clarify the sterile anomaly by precisely/independently measuring **both** ν_e appearance and ν_μ disappearance:




ν_e appearance: LSND 99% CL region covered at 5 σ level



3-5 σ ν_μ disapp. SBN sensitiv.

- Using the same detector technology for all the 3 detectors will greatly reduce the systematic errors: **SBND** (near detector) will provide the "initial" beam composition and spectrum.
- The great ν_e identification capability of LAr-TPC will help reduce the NC background.
- During SBN operations, ICARUS will also collect ~ 2 GeV neutrinos from **NuMI** (Neutrino Main Injector) Off-Axis beam. This will be an asset for the future long-baseline project as DUNE.



ICARUS at FNAL is facing a more challenging experimental condition than at LNGS, requiring the recognition of $O(10^6)$ ν interactions among 11 kHz of cosmic rays.




- Therefore, T600 underwent an intensive overhauling at CERN in the **Neutrino Platform** framework from 2015 to 2017, before shipping to US.
- Several technology developments were introduced *while maintaining the already achieved performance at LNGS run*:
 - new cold vessels, with a purely passive insulation;
 - renovated LAr cryogenics/purification equipment;
 - improvement of the cathode planarity;
 - upgrade of the PMT system: higher granularity and ns time resolution;
 - new faster, higher-performance read-out electronics.
- **3 m concrete overburden** to remove contribution from charged hadrons/ γ 's.
- External cosmic ray tagger to correlate residual muons with TPC signals.

ICARUS@SBN exploits 90 PMTs per TPC (5% coverage, 15 phe/MeV) that provides:

- Sensitivity to low energy events (~ 100 MeV)
- Good spatial resolution (≤ 50 cm)
- \approx ns timing resolution
- Possible cosmics identification by PMT space/time pattern





- All PMTs tested at room temperature in a dedicated dark room at CERN
- A subset of 60 PMTs tested immersed in LAr to compare the PMT performance in cryogenic environment to room temperature



New TPC readout electronics


- Outside the cryostat
- Serial 12-bit ADC, fully synchronous in the whole detector
- CAEN A2795 64-chan modules.
- More compact layout: both analog+digital electronics hosted on a single flange.



- Lower noise ~ 1200 e- equivalent ($\sim 20\%$ S/N improvement w.r.t. LNGS electronics).
- Shorter shaping time ($\sim 1.5 \mu\text{s}$ for all planes) and drastic reduction of undershoot after large signals.
- Induction 2 signal keeps bipolar shape \rightarrow allows calorimetric measurement in this plane, to improve ν_e identification efficiency by $\sim 20\%$.


*New electronics extensively tested on a
50-liter TPC@CERN*

JINST 13 (2018) P12007




*Placement of
ICARUS
(August 2018)*

*Chimneys installation
(October 2018)*




*Feedthrough installation
(December 2018)*




Readout electronics

*Power
supply*




Installing the readout electronics (May 2019)


- Top cold shields and top CRT support installed.
- Installation of proximity cryogenics completed.
- ICARUS Vacuum phase started June 5th!
- Side CRT installation also ongoing.
- Director's Review in December 2018 recognized the great progress of SBN.





Connectivity test



Side CRT installation






- All the feedthrough flanges and the mini-crates with the TPC wire read-out electronics (576 channels + optical links) has been installed.
- A test of the full readout chain, from wires to DAQ, has been performed in April/May for all the mini-crates :
 - Allowed to check readout and set baseline for future noise monitoring
 - Noise measured on random triggers and test pulses
 - Noise RMS ~ 1700 e $^-$, not too far from ~ 1200 e $^-$ measured in CERN 50-liter setup: grounding conditions were still far from optimal



*The successful readout test confirms
the good performance of the full TPC electronics!*

- A detailed understanding of detector-related systematics and their correlation across near/far detectors **will be crucial** to SBN physics.
- **Common reconstruction tools** and oscillation analysis are therefore fundamental.
- ICARUS joined the **LArSoft** framework: mutual sharing of algorithms and tools and cross-check between different reconstruction approaches.
- Full simulation performed with realistic geometry and signals from all sub-detectors (TPC,PMT,CRT).


Scheme of detector geometry



electron lifetime
(reco vs. simulation)

angle between sim/reco direction for
EM showers

- TPC/trigger electronics installation to be completed and tested by summer 2019.
- PMT electronics installation also to be completed during the summer.
- ICARUS expected to be ready to fill by end of September.
- After cryogenics commissioning, cool down and filling, ICARUS T600 should be operational in the last quarter of 2019.
- Commissioning of CRT, DAQ, trigger and slow controls will follow.
- Data-taking for physics is expected by the end of this year.



- The ICARUS-T600 successful 3-year run at LNGS proved that LAr-TPC technology is mature and ready for large-scale neutrino physics experiments.
- ICARUS searched for LSND-like anomaly via ν_e appearance in the CNGS beam. The negative result constrained significantly the allowed parameter region.
- The SBN project at FNAL is expected to clarify the sterile neutrino puzzle, by looking at both appearance and disappearance channels with three LAr-TPCs.
- After an extensive refurbishing, ICARUS is being installed as the SBN far detector at FNAL. Data taking expected in 2019, near detector in 2021.
- ICARUS will see first neutrinos by the end of this year !



Thank you!

Backup

M. Diwan, X. Qian, J. Stewart, B. Viren, E. Worcester, M. Worcester, C. Zhang
Brookhaven National Laboratory, USA

B. Behera, T. Boone, I. Caro Terrazas, C. Hilgenberg, J. Mueller, M. Mooney, H. Rogers,
D. Warner, R.J. Wilson
Colorado State University, USA

W. F. Badgett, L.F. Bagby, M. Betancourt, K. Biery, S. Brice, J. Brown, G. Cerati, R. Doubnik, A. Fava,
F.G. Garcia, M. Geynisman, S. Hahn, C. James, W. Ketchum, G. Lukhanin, S. Marcocci, A. Mazzacane, T.
Nichols, A. Prosser, R. Rameika, R. Rechenmacher, A. Schukraft, A. Soha, D. Torretta, P. Wilson, M.
Wospakrik, J. Zennamo, M. Zuckerbrot
Fermi National Accelerator Laboratory, USA

D. Cherdack, A. Wood

University of Houston, USA

V. Bellini, C. Petta, C. Suter, F. Tortorici
INFN Sez. di Catania and University, Catania, Italy

C. Rubbia

INFN GSSI, L'Aquila, Italy

C. Vignoli

INFN LNGS, Assergi (AQ), Italy

R. Benocci, M. Bonesini, A. Falcone, M. Torti
INFN Sez. di Milano Bicocca, Milano, Italy

A. Cocco

INFN Sez. di Napoli, Napoli, Italy

A. Braggiotti, S. Centro, C. Farnese, D. Gibin, A. Guglielmi, G. Meng,
F. Pietropaolo, F. Varanini, S. Ventura

INFN Sez. di Padova and University, Padova, Italy

A. Menegolli, C. Montanari, A. Rappoldi, G.L. Raselli, M. Rossella, A. Scaramelli

INFN Sez. di Pavia and University, Pavia, Italy

V. Paolone, H. Su, A. Wickremasinghe

University of Pittsburgh, USA

M. Convery, L. Domine, F. Drielsma, B. Nelson, G. Petrillo, H. Tanaka, K. Terao, Y.T. Tsai, T. Usher

SLAC National Accelerator Laboratory, Stanford, USA

H. Budd, R. Howell, K.S. McFarland

University of Rochester, USA

J. Asaadi, H. Carranza, A. Chatterjee, Z. Williams, J. Yu

University of Texas (Arlington), USA

T. Coan

Southern Methodist University (USA)

O.G. Miranda, G. Moreno Granados

CINVESTAV-IPN (Mexico)




CINVESTAV




Neutrino Platform NP01

- *ICARUS Spokesman: C. Rubbia, GSSI*
- *7 INFN groups, 9 USA institutions, 1 Mexico institution, more than 90 collaborators.*
- *Major contributions to the ICARUS cryogenics and cosmic ray tagger from our partners at CERN and INFN (Bologna, Lecce, Milano, Napoli, Genoa and LNS).*

- The sterile neutrino scenario is far from understood and needs a definitive clarification
- Some “anomalies” from accelerators (LSND), reactor, neutrino sources, point out to flavour transitions in the $\Delta m^2 \sim 1 \text{ eV}^2$ range
- However, no evidence of oscillations in ν_μ disappearance data (MINOS, IceCube)
- Tension between ν_e appearance and ν_μ disappearance results. *Measuring both channels with the same experiment will help disentangle*





Combined analyses



- A comparison between far/near detector is crucial for any accelerator experiment, with a better control of backgrounds and systematics

SBN satisfies these requirements: it could have a crucial role in solving the sterile neutrino puzzle!

- ICARUS at LNGS was also exposed to atmospheric neutrinos (exposure ~ 0.74 kt year)
first observation of atmospheric neutrinos with a LAr-TPC
- 14 events found ($8 \nu_e \text{ CC} + 6 \nu_\mu \text{ CC}$) vs. 18 expected (taking into account: triggering, filtering and scanning efficiencies)
- Very good benchmark for the forthcoming SBN experiment: similar energy/features.
Useful to develop filtering and reconstruction tools



- Surrounds the cryostat with two layers of plastic scintillators: 1100 m²
- Tags incident cosmic or beam-induced muons with high efficiency (95%) giving spatial and timing coordinates of the track entry point.
- Reconstructed CRT hits are matched to activity in the LAr volume.
- Few ns time resolution allows measuring direction of incoming/outgoing particle propagation via time of flight

TOP:

*~ 400 m²: roof+angled parts
Will catch ~80% cosmic ms
2 strip layers (X+Y)
SiPM readout*

SIDES:

*~ 500 m² on four sides
Old MINOS veto modules
parallel strips
SiPM readout*

BOTTOM:

*~ 200 m², already installed
D-Chooz veto modules
2 parallel layers
PMT readout*

