

The dominant contributions to the κ_{λ} sensitivity derive from the di-boson decay channels $\gamma\gamma$, ZZ, WW and from the ggF and ttH production modes [5].

- A simultaneous fit is performed to $(\kappa_{\lambda, \kappa_F})$ and $(\kappa_{\lambda, \kappa_V})$, the modifiers of the Higgs boson coupling to fermions and to massive vector bosons.
- Other fit configurations (e.g. $\kappa_{\lambda}, \kappa_{F}, \kappa_{V}$) result in nearly no sensitivity on



References

[1] ATLAS Collaboration, *Combination of searches for Higgs boson pairs in pp collisions at* √s =13 TeV with the ATLAS detector, arXiv:1906.02025, 2019;
[2] G. Degrassi, P. P. Giardino, F. Maltoni and D. Pagani, *Probing the Higgs self coupling via single Higgs production at the LHC*, JHEP **12** (2016) 080, arXiv: 1607.04251;
[3] F. Maltoni, D. Pagani, A. Shivaji and X. Zhao, *Trilinear Higgs coupling determination via single-Higgs differential measurements at the LHC*, Eur. Phys. J. **C77** (2017) 887, arXiv:1709.08649;
[4] D. de Florian et al., Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector, (2016), arXiv: 1610.07922 [hep-ph];
[5] ATLAS Collaboration, *Constraint of the Higgs boson self-coupling from Higgs boson differential production and decay measurements*, ATL-PHYS-PUB-2019-009, 2019.