



Contribution ID: 658

Type: Poster

Calibration of the highly segmented SoLid antineutrino detector

Monday 15 July 2019 18:30 (1h 30m)

SoLid is a short baseline neutrino experiment, which is currently operating a 1.6 tons detector at the SCK•CEN BR2 research reactor in Belgium. SoLid will address the study of the so called Reactor Antineutrino Anomaly (RAA), whose origin could be the existence of a light sterile neutrino state with a mass around the eV scale. In addition, it will perform a new measurement of the antineutrino energy spectrum produced by the ^{235}U isotope, which will help in the understanding of the 5-MeV distortion observed in previous reactor antineutrino experiments.

SoLid leverages a novel technology, combining PVT cubes of $5\times 5\times 5\text{ cm}^3$ dimensions and $^6\text{LiF:ZnS(Ag)}$ screens of $\sim 250\text{ }\mu\text{m}$ thickness. To detect antineutrino interactions, signals are readout by a network of wavelength shifting fibers and SiPMs. The fine granularity (12800 cells) provides powerful tools to distinguish signal from background, but presents a challenge in ensuring homogeneous detector response and calibrating the energy scale and neutron detection efficiency. In this poster the methods that have been developed for the calibration of such a segmented detector will be described. In addition, the calibration results will be presented.

Primary author: MANZANILLAS, Luis (Centre National de la Recherche Scientifique (FR))

Presenter: MANZANILLAS, Luis (Centre National de la Recherche Scientifique (FR))

Session Classification: Wine & Cheese Poster Session

Track Classification: Detector R&D and Data Handling