## 4th International Conference on the Initial Stages in High-Energy Nuclear Collisions



Contribution ID: 85

Type: not specified

## Lambda polarization as a probe of initial state

Wednesday 20 September 2017 18:00 (20 minutes)

In hydrodynamic approach to heavy ion collisions, hadrons with nonzero spin produced out of the fluid can acquire polarization via spin-vorticity thermodynamic coupling mechanism [1]. The hydrodynamical quantity steering the polarization is the thermal vorticity, that is minus the antisymmetric part of the gradient of four-temperature field  $\beta^{\mu} = u^{\mu}/T$ .

Based on this idea, it has been shown in the framework of cascade+viscous hydro model, UrQMD+vHLLE [2] that in Au-Au collisions at RHIC Beam Energy Scan (BES) the mean polarization of Lambda hyperons grows with decreasing collision energy up to 1.5% at  $\sqrt{s} = 7.7$  GeV RHIC Au-Au collisions. This goes in line with recent measurements of Lambda polarization by STAR experiment [3].

We show how the excitation function of the mean polarization is related to vorticity in the initial state and explore effects of initial state fluctuations on it. Finally, we propose a new polarization observable which can be probed in heavy ion collisions at LHC energies.

[1] F. Becattini, V. Chandra, L. Del Zanna, E. Grossi, Ann. Phys. 338 (2013) 32.

[2] I. Karpenko, F. Becattini, arXiv:1610.04717, to be published in EPJC.

[3] STAR collaboration, arXiv:1701.06657

Authors: Dr KARPENKO, Iurii (INFN Florence); Prof. BECATTINI, Francesco (Unversity of Florence)

Presenter: Dr KARPENKO, Iurii (INFN Florence)

Session Classification: CGC / Vorticity