

Charm meson physics at BESIII

Thursday 28 September 2017 14:55 (20 minutes)

BESIII Collaboration has accumulated the world's largest e^+e^- collision samples at $E_{cm} = 3.773, 4.009, 4.18$ GeV. From analyses of pure leptonic decay $D(s)^+ \rightarrow l^+ \nu$ ($l = \mu$ or τ), semileptonic decays of $D \rightarrow K(\pi)l^+ \nu$ ($l = e$ or μ), $D^+ \rightarrow K^+ \pi^0 e^+ \nu$, $D^+ \rightarrow \omega e^+ \nu$, $D^0 \rightarrow f_0(980) e^+ \nu$, $D_s^+ \rightarrow \eta e^+ \nu$, and $D_s^+ \rightarrow \eta' e^+ \nu$, we report the determinations of their absolute branching fractions, CKM matrix elements $|V_{cs}(d)|$, the $D(s)^+$ decay constants, the form factors of D semi-leptonic decays. These are important to calibrate the LQCD calculations of decay constants and form factors and to test the CKM unitarity. We have performed an amplitude analysis of $D^0 \rightarrow K^+ \pi^- \pi^+ \pi^-$, and have measured the asymmetries of $D^+ \rightarrow K_S^0 L^+ K^+ \pi^0$ and $D^0 \rightarrow K_S^0 L^+ \pi^0 \pi^0$, as well as the branching fractions for $D^0 \rightarrow PP$ and some decays containing two K Ss. In addition, we also measure the branching fractions for $D_s^+ \rightarrow \omega \pi^+$, ωK^+ and baryon decay of $p\pi$.

Author: DOU, Zhenglei

Presenter: DOU, Zhenglei

Session Classification: Hadron decays

Track Classification: Hadron decays