

# Free Energy of a Large- $N$ Pion Gas and Chiral Symmetry Restoration

Santiago Cortés<sup>1,†</sup>, Ángel Gómez Nicola<sup>2</sup>, and John Morales<sup>3</sup>

<sup>1</sup>Departamento de Física, Univ. de Los Andes, 111711 Bogotá, Colombia.

<sup>2</sup>Departamento de Física Teórica II. Univ. Complutense. 28040 Madrid. Spain.

<sup>3</sup>Departamento de Física, Univ. Nacional de Colombia, 111321 Bogotá, Colombia.

†js.cortes125@uniandes.edu.co



## Abstract

We study thermal properties of a large- $N$  massless pion gas using a low-energy QCD approach given by an  $O(N+1)/O(N)$  Nonlinear Sigma Model. We build diagrammatically the associated finite free energy to  $\mathcal{O}(TM^3)$  in the pion mass expansion through an effective vertex that considers all the contributions coming from closed diagrams. Subsequently, we calculate finite order parameters such as the quark condensate and its respective derivative, i.e., the scalar susceptibility, in the chiral limit, along with their associated critical exponents. These results are compared with our previous unitarized scattering analyses for the chiral transition universality class, thus showing a reasonable agreement both with lattice simulations and these resonant studies.

## 1. Introduction and Formalism

Low-energy QCD phenomena such as chiral symmetry restoration are essential to understand the behavior of hadron matter created in heavy-ion collision experiments such as LHC-ALICE and RHIC, a fact showed and confirmed by lattice simulations [1]. Our purpose is to study the critical behavior of a Large- $N$  pion gas through its free energy by considering an  $O(N+1)/O(N)$  Nonlinear Sigma Model with an explicit chiral symmetry breaking term given by the following Lagrangian [2]:

$$\mathcal{L}_{NL\text{SM}} = \frac{1}{2} \left[ \delta_{ab} + \frac{1}{NF^2} \frac{\pi_a \pi_b}{1 - \pi^2/NF^2} \right] \partial_\mu \pi^a \partial^\mu \pi^b + NF^2 M^2 \sqrt{1 - \frac{\pi^2}{NF^2}}, \quad (1)$$

$$\pi^2 = \sum_{a=1}^N \pi_a \pi_a,$$

where  $M^2$  and  $\sqrt{NF}$  are respectively the pion mass and the pion decay constant in the chiral limit. Large- $N$  closed dominant diagrams ( $\mathcal{O}(N)$ ) are obtained from the mass vertex in (1). Momentum contributions are subdominant either by their large- $N$  contribution (i.e.,  $\mathcal{O}(N^0)$ ) or their associated Feynman rules are canceled due to its odd character. Bearing this in mind, we proceed to build a massive effective thermal vertex as it is shown in the Figure below.

$$\begin{aligned} & \text{Diagram} = -\frac{M^2}{8NF^2} \left[ \text{Diagram 1} + \frac{1}{2} \text{Diagram 2} + \frac{5}{16} \text{Diagram 3} \right. \\ & \left. + \frac{7}{32} \text{Diagram 4} + \dots + 8(-1)^{k+1} \binom{1/2}{k+2} \text{Diagram 5} \right] = -\frac{M^2}{8NF^2} g\left(\frac{G_1}{F^2}\right) \end{aligned}$$

This vertex takes into account all possible insertions of thermal tadpoles coming from diagrams with six or more external legs. Its Feynman rule is written in terms of a function  $g(x)$  that reads


$$g(x) = -\frac{8}{x^2} \left[ \sqrt{1-x} - 1 + \frac{x}{2} \right] = -8 \sum_{k=0}^{\infty} (-1)^k \binom{1/2}{k+2} x^k = 1 + \frac{x}{2} + \frac{5}{16}x^2 + \dots \quad (2)$$

$G_1(M, T)$  is taken as in [3]. Since this is not a scattering approach, we do not consider any combinatory factor linked with the way the pion lines are attached either with external legs or loops. Now, we are able to diagrammatically construct the partition function for  $N$  massless pions.

## 2. Free Energy and Order Parameters


### 2.1 Free Energy

By taking the limit  $M/F \rightarrow 0$  and considering large- $N$  dominant contributions up to  $\mathcal{O}(M^3T)$ , we obtain a free energy that does not need to be renormalized after examining the contributions coming from the closed diagrams showed in the next Figure [4]. The dot (a) represents the constant term that comes from the expansion of (1), whereas the Daisy diagram (e) depicts multiple insertions of the mass vertex along a single loop. Any other diagram not listed in the figure is either large- $N$  subdominant or contributes as  $\mathcal{O}(M^4)$  (e.g., foam-like diagrams), which are divergent terms avoided in this limit.



### 2.2 Scalar Quark Condensate and Susceptibility

First and second derivatives of this partition function allows us to respectively find the quark condensate and the scalar susceptibility, whose thermal evolution in the chiral limit  $M \rightarrow 0^+$  is plotted in the following figures [4].



We also plot the  $f_0(500)$ -saturated scalar susceptibility [5], along with the well-known ChPT results [3]. We find in an exact way that the quark condensate scales as  $(1 - T^2/T_c^2)^\beta$  with  $\beta = 0.5$  in the critical region, whilst the critical exponent of the susceptibility is  $\gamma = 0.75$ , where  $\chi_S(T) \sim (1 - T^2/T_c^2)^{-\gamma}$ . Furthermore, we check that the condensate cancels out exactly at a critical temperature  $T_c = \sqrt{12}F \approx 174$  MeV, thus showing that the system undergoes a second-order phase transition.

## 3. Conclusions

Our results show that this critical behavior rationally agrees with those given for the universality classes expected from lattice simulations. However, our numerical values obtained for  $\gamma$  and  $T_c$  differ from our previous scattering analysis since this approach only takes into account contributions coming from Nambu-Goldstone bosons.

## References

- [1] Y. Aoki, S. Borsanyi, S. Durr, Z. Fodor, S. D. Katz, S. Krieg and K. K. Szabo, JHEP **0906**, 088 (2009).
- [2] A. Dobado and J. Morales, Phys. Rev. D **52**, 2878 (1995).
- [3] P. Gerber and H. Leutwyler, Nucl. Phys. B **321**, 387 (1989).
- [4] S. Cortés, A. Gómez Nicola and J. Morales, Phys. Rev. D **94**, no. 11, 116008 (2016).
- [5] S. Cortés, A. Gómez Nicola and J. Morales, Phys. Rev. D **93**, no. 3, 036001 (2016).