

AdS/QCD Modified Soft Wall Model and Light Meson Spectra

Tuesday 26 September 2017 20:45 (15 minutes)

We analyze here the mass spectrum of light vector and scalar mesons applying the novel approach developed in [1], where a modified soft wall model that includes a UV-cutoff at a finite z -position in the AdS space is used, thus introducing an extra energy scale. For this model, we found that the masses for the scalar and vector spectra are well fitted within $\delta_{\text{RMS}} = 6.88\%$ for these states, with non-linear trajectories given by two common parameters, the UV locus z_0 and the quadratic dilaton profile slope κ . We also concluded that in this model the $f_0(500)$ scalar resonance cannot be fitted holographically as a $q\bar{q}$ state since we could not find a trajectory that included this pole. This result is in agreement with the most recent phenomenological and theoretical methods.

[1] N. R. F. Braga, M. A. Martin Contreras and S. Diles,
Phys. Lett. B 763, 203 (2016).

Authors: CORTES, Santiago (Universidad de los Andes); Dr MARTÍN CONTRERAS, Miguel Ángel (Universidad de los Andes, Bogotá (Colombia)); Prof. ROLDÁN, José Rolando (Universidad de los Andes, Bogotá (Colombia))

Presenter: CORTES, Santiago (Universidad de los Andes)

Session Classification: Poster session

Track Classification: Spectroscopy of mesons