



Contribution ID: 42

Type: not specified

Confronting jet quenching with jet grooming: splitting function and jet mass distribution in heavy ion collisions

Monday, July 17, 2017 6:00 PM (20 minutes)

We present the first calculations of the momentum sharing and angular separation distributions between the leading subjects inside a reconstructed jet, as well as the jet mass distribution modification in heavy ion collisions. These observables are sensitive to the early and late stages of the in-medium parton shower evolution and allow us to probe the quark-gluon plasma across a wide range of energy scales. We use the medium-induced splitting functions obtained in the framework of soft-collinear effective theory with Glauber gluon interactions to calculate the subjet distributions. Qualitative and in most cases quantitative agreement between theory and preliminary CMS measurements suggests that the parton shower in heavy ion collisions can be dramatically modified early in the branching history. Predictions for the subjet angular distribution is also presented which will illuminate the nature of the medium-induced radiations. On the other hand, using renormalization group techniques we resum the jet mass at next-to-leading logarithmic accuracy for groomed jets with small radii, and we include the medium contributions consistently. We find that the jet mass modification is sensitive to the medium scale and allows for a precise extraction of the medium properties.

Primary author: Dr CHIEN, Yang-Ting (Massachusetts Institute of Technology)

Presenter: Dr CHIEN, Yang-Ting (Massachusetts Institute of Technology)

Session Classification: Measurements and Modeling