

$|V_{ub}|$ from QCD Sum Rules on the Light-Cone

Patricia Ball

IPPP, Durham

FPCP07, 13 May 2007

Based on Ball/Zwicky, hep-ph/0406232; Ball, hep-ph/0611108.



Theory Input for Semileptonic Decays

Form factors:

$$\begin{aligned} & \langle \pi(p) | \bar{u} \gamma_\mu (1 - \gamma_5) b | B(p + q) \rangle \\ &= (q + 2p)_\mu f_+(q^2) + \frac{m_B^2 - m_\pi^2}{q^2} q_\mu (f_0(q^2) - f_+(q^2)) \end{aligned}$$

$$0 \leq q^2 \leq (m_B - m_\pi)^2 \quad \longleftrightarrow \quad m_\pi \leq E_\pi \leq \frac{1}{2m_B} (m_B^2 - m_\pi^2)$$

$$0 \leq q^2 \leq 26.4 \text{ GeV}^2 \quad \longleftrightarrow \quad 0.14 \text{ GeV} \leq E_\pi \leq 2.6 \text{ GeV}$$

Theoretical methods:

- lattice → D. Becirevic's talk today
- SCET/dispersive constraints → I. Stewart's talk Tue
- QCD sum rules on the light-cone → this talk!

QCD Sum Rules on the Light-Cone

Basic quantity: correlation function:

$$i \int d^4 y e^{iqy} \langle \pi(p) | T[\bar{u}\gamma_\mu b](y) [m_b \bar{b} i\gamma_5 d](0) | 0 \rangle \stackrel{\text{LCE}}{=} \sum_n T_H^{(n)} \otimes \phi_\pi^{(n)}$$

- $\phi_\pi^{(n)}$: π distribution amplitudes (DAs)
- $T_H^{(n)}$: perturbative amplitudes
- n : twist
- LCE: light-cone expansion

$$= 2p_\mu \left(f_+(q^2) \frac{m_B^2 f_B}{m_B^2 - p_B^2} + \text{higher poles and cuts} \right) + \dots$$

- B meson described by **Euclidean current** + plus analytical continuation

QCD Sum Rules on the Light-Cone

Features of LCSRs:

- terms in LCE ordered in powers of $1/m_b \rightarrow$ need to include **higher-twist terms** ($n > 2$)
- $\sum T_H^{(n)} \otimes \phi_\pi^{(n)}$ implies **factorization** – valid at higher twist?
 - calculate $O(\alpha_s)$, known for
 - T2 (π (Khodjamirian et al. 97, Ball et al. 97), ρ (Ball/Braun 98))
 - T3 (π (Ball/Zwicky 2001))
 - \rightarrow **factorization OK**
- use standard SR techniques: Borel-transformation, continuum model
 - introduce irreducible **systematic uncertainty** $\sim 10\%$

QCD Sum Rules on the Light-Cone

- Ball/Zwicky 04:


$$f_+(0) = 0.258 \pm 0.031$$

with theory input for leading-twist π distribution amplitude $\phi_{\pi;2}$

- Ball/Zwicky 05: constrain $\phi_{\pi;2}$ from experimental q^2 spectrum of $B \rightarrow \pi e \nu$:

$f_+(0) \approx 0.27$ and


$$|V_{ub}| = (3.2 \pm 0.4) \cdot 10^{-3}$$



BZ 04

Results for $B \rightarrow \rho e \nu$ also available — but less experimental information.

Theory Assisted by Experiment



2006 BaBar results for q^2 spectrum in $B \rightarrow \pi e \nu$ in **12 bins** (up from 5 bins in 2005)

Strategy: Parametrise form factor, fit to data, extract $|V_{ub}|f_+(0)$.

Form Factor Parametrisations

- **Becirevic/Kaidalov (BK)** :

$$f_+(q^2) = \frac{f_+(0)}{(1 - q^2/m_{B^*}^2)(1 - \alpha_{\text{BK}} q^2/m_B^2)},$$

where α_{BK} determines the **shape** of f_+ and $f_+(0)$ the **normalisation**;

Form Factor Parametrisations

- **Becirevic/Kaidalov (BK)** :

$$f_+(q^2) = \frac{f_+(0)}{(1 - q^2/m_{B^*}^2) (1 - \alpha_{\text{BK}} q^2/m_B^2)},$$

where α_{BK} determines the **shape** of f_+ and $f_+(0)$ the **normalisation**;

- **Ball/Zwicky (BZ)**:

$$f_+(q^2) = f_+(0) \left(\frac{1}{1 - q^2/m_{B^*}^2} + \frac{r q^2/m_{B^*}^2}{(1 - q^2/m_{B^*}^2) (1 - \alpha_{\text{BZ}} q^2/m_B^2)} \right),$$

with the two **shape parameters** α_{BZ}, r and the **normalisation** $f_+(0)$; BK is a variant of BZ with $\alpha_{\text{BK}} := \alpha_{\text{BZ}} = r$.

Form Factor Parametrisations

- the **AFHNV parametrisation** (Flynn et al.), based on an $(n + 1)$ -subtracted Omnes representation of f_+ :

$$f_+(q^2) \stackrel{n \gg 1}{\approx} \frac{1}{m_{B^*}^2 - q^2} \prod_{i=0}^n [f_+(q_i)^2 (m_{B^*}^2 - q_i^2)]^{\alpha_i(q^2)},$$

$$\text{with } \alpha_i(s) = \prod_{j=0, j \neq i}^n \frac{s - s_j}{s_i - s_j};$$

the **shape parameters** are $f_+(q_i^2)/f_+(q_0^2)$ with q_0^2, \dots, q_n^2 the subtraction points; the **normalisation** is given by $f_+(0)$.

Form Factor Parametrisations

- the **BGL** parametrisation based on analyticity of f_+ :

$$f_+(q^2) = \frac{1}{P(q^2)\phi(q^2, q_0^2)} \sum_{k=0}^{\infty} a_k(q_0^2) [z(q^2, q_0^2)]^k, \quad \sum_k a_k^2 \leq 1,$$

$$z(q^2, q_0^2) = \frac{\{(m_B + m_\pi)^2 - q^2\}^{1/2} - \{(m_B + m_\pi)^2 - q_0^2\}^{1/2}}{\{(m_B + m_\pi)^2 - q^2\}^{1/2} + \{(m_B + m_\pi)^2 - q_0^2\}^{1/2}}$$

q_0^2 : free parameter, determines maximum $|z|$; define

- BGLa** : $q_0^2 = 20.1 \text{ GeV}^2, |z| < 0.28$

- BGLb** : $q_0^2 = 0, |z| < 0.52$

- systematic expansion in the **small parameter** z ; **truncate** at k_{\max} ;


choose $k_{\max} = 2$ for **BGLa** and $k_{\max} = 3$ for **BGLb**.

$|V_{ub}|f_+(0)$ from data

Param.	$ V_{ub} f_+(0)$	Remarks
BK	$(9.3 \pm 0.3 \pm 0.3) \times 10^{-4}$	$\chi_{\min}^2 = 8.74/11$ dof $\alpha_{\text{BK}} = 0.53 \pm 0.06$
BZ	$(9.1 \pm 0.5 \pm 0.3) \times 10^{-4}$	$\chi_{\min}^2 = 8.66/10$ dof $\alpha_{\text{BZ}} = 0.40_{-0.22}^{+0.15}$, $r = 0.64_{-0.13}^{+0.14}$
BGLa	$(9.1 \pm 0.6 \pm 0.3) \times 10^{-4}$	$\chi_{\min}^2 = 8.64/10$ dof
BGLb	$(9.1 \pm 0.6 \pm 0.3) \times 10^{-4}$	$\chi_{\min}^2 = 8.64/9$ dof
AFHNV	$(9.1 \pm 0.3 \pm 0.3) \times 10^{-4}$	$\chi_{\min}^2 = 8.64/8$ dof
SCET	$(8.0 \pm 0.4) \times 10^{-4}$	from $B^- \rightarrow \pi^- \pi^0$ (Arnesen et al.) (tree-level, no $1/m_b$ corrections)

All parametrisations agree – model-independent result!

Fitted Form Factor



Left panel: **best-fit form factors f_+ as a function of q^2** . The line is an overlay of all five parametrisations.

Right panel: **best-fit form factors normalised to BGLa**.

Solid line: BK, long dashes: BZ, short dashes: BGLb, short dashes with long spaces: AFHNV.

Results for $|V_{ub}|$

Procedure 1: take FF from theory calculation, fit to **BK** and extract $|V_{ub}|$ from experimental partial branching ratio
($q^2 \leq 16 \text{ GeV}^2$ for LCSR, $q^2 \geq 16 \text{ GeV}^2$ for lattice)

LCSR	$f_+(0) = 0.26 \pm 0.03, \quad \alpha_{\text{BK}} = 0.63_{-0.21}^{+0.18}$ $ V_{ub} = (3.5 \pm 0.6(\text{th}) \pm 0.1(\text{exp})) \times 10^{-3}$ $ V_{ub} f_+(0) = (9.0_{-0.6}^{+0.7} \pm 0.4) \times 10^{-4}$
HPQCD	$f_+(0) = 0.21 \pm 0.03, \quad \alpha_{\text{BK}} = 0.56_{-0.11}^{+0.08}$ $ V_{ub} = (4.3 \pm 0.7 \pm 0.3) \times 10^{-3}$ $ V_{ub} f_+(0) = (8.9_{-0.9}^{+1.2} \pm 0.4) \times 10^{-4}$
FNAL	$f_+(0) = 0.23 \pm 0.03, \quad \alpha_{\text{BK}} = 0.63_{-0.10}^{+0.07}$ $ V_{ub} = (3.6 \pm 0.6 \pm 0.2) \times 10^{-3}$ $ V_{ub} f_+(0) = (8.2_{-0.8}^{+1.0} \pm 0.3) \times 10^{-4}$

Results for $|V_{ub}|$

Procedure 2: take FF from theory, fit to experimentally determined shape, **BGLa**, obtain $f_+(0)$, extract $|V_{ub}|$ from full branching ratio.

LCSR	$f_+(0) = 0.26 \pm 0.03$ $ V_{ub} = (3.5 \pm 0.4(\text{shape}) \pm 0.1(\mathcal{B})) \times 10^{-3}$
HPQCD	$f_+(0) = 0.21 \pm 0.03$ $ V_{ub} = (4.3 \pm 0.5 \pm 0.1) \times 10^{-3}$
FNAL	$f_+(0) = 0.25 \pm 0.03$ $ V_{ub} = (3.7 \pm 0.4 \pm 0.1) \times 10^{-3}$

- reduced theoretical uncertainty as **shape** of FF is fixed by experimental data
- reduced experimental uncertainty as total $\mathcal{B}(B \rightarrow \pi e \nu)$ can be used

Summary

- form factor calculations from QCD sum rules on the light-cone in mature shape, no scope for major improvement
- LCSR predictions for small and moderate $q^2 < 16 \text{ GeV}^2$
 \longleftrightarrow LQCD predictions for large $q^2 > 16 \text{ GeV}^2$
- reduce error of $|V_{ub}|$ determination by **fixing shape of form factor from experiment** instead of theory data
- model-independent result: $|V_{ub}|f_+(0) = (9.1 \pm 0.7) \times 10^{-4}$
- both LCSR and FNAL prefer **small** $|V_{ub}| \sim 3.6 \times 10^{-3}$
- HPQCD points at **larger** $|V_{ub}| \sim 4.3 \times 10^{-3}$
- new analysis using Omnes relations, including also Belle/Cleo data and fitting to all LQCD data (Flynn/Nieves 07):
 - $|V_{ub}| = (3.90 \pm 0.32(\text{stat}) \pm 0.18(\text{syst})) \times 10^{-3}$ and
 $|V_{ub}|f_+(0) = (8.8 \pm 0.8) \times 10^{-4}$