



A DIGRESSION ON DIVERGENCES IN ELECTROMAGNETISM

Ezio Todesco

European Organization for Nuclear Research (CERN)



FOREWORD

- The equations ruling the construction of an electromagnet have some aspects that can be used as a paradigm in teaching some aspects of physics and mathematics



CONTENTS

- Maxwell equations
- The field of a current line
 - Analytic functions and Taylor series
 - Convergence domain and singularities
- The two divergences of electromagnetism
- An outlook on scales

- Maxwell equations

$$\vec{\nabla} \cdot \vec{B} = 0$$

$$\vec{\nabla} \times \vec{E} + \frac{1}{c} \frac{\partial B}{\partial t} = 0$$

$$\vec{\nabla} \cdot \vec{E} = 4\pi\rho$$


$$\vec{\nabla} \times \vec{B} - \frac{1}{c} \frac{\partial E}{\partial t} = \frac{4\pi}{c} \vec{J}$$

$$\vec{B} = \vec{\nabla} \times \vec{A}$$

$$\vec{E} = -\vec{\nabla}\phi - \frac{1}{c} \frac{\partial A}{\partial t}$$

- Hide few monsters:

- The presence of c means that they cannot be invariant with Galileo
- So Maxwell has inside the special relativity



James Clerk Maxwell,
Scottish
(13 June 1831 - 5 November 1879)



Ridley Scott, Alien (1979)

- Maxwell equations

$$\vec{\nabla} \cdot \vec{B} = 0$$

$$\vec{\nabla} \times \vec{E} + \frac{1}{c} \frac{\partial B}{\partial t} = 0$$

$$\vec{\nabla} \cdot \vec{E} = 4\pi\rho$$


$$\vec{\nabla} \times \vec{B} - \frac{1}{c} \frac{\partial E}{\partial t} = \frac{4\pi}{c} \vec{J}$$

$$\vec{B} = \vec{\nabla} \times \vec{A}$$

$$\vec{E} = -\vec{\nabla}\phi - \frac{1}{c} \frac{\partial A}{\partial t}$$

- Hide few monsters:

- Equations have divergences for infinitely small charge density and infinitely large current density
- So the charge cannot be a point, it has some dimension



James Clerk Maxwell,
Scottish
(13 June 1831 - 5 November 1879)



CONTENTS

- Maxwell equations
- The field of a current line
 - Analytic functions and Taylor series
 - Convergence domain and singularities
- The two divergences of electromagnetism
- An outlook on scales


MAXWELL EQUATIONS

- **Maxwell equations** for magnetic field

$$\nabla \cdot \mathbf{B} = \frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} = 0 \quad \nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t}$$

- In absence of charge and magnetized material

$$\nabla \times \mathbf{B} = \left(\frac{\partial B_y}{\partial z} - \frac{\partial B_z}{\partial y}, \frac{\partial B_z}{\partial x} - \frac{\partial B_x}{\partial z}, \frac{\partial B_x}{\partial y} - \frac{\partial B_y}{\partial x} \right) = 0$$



James Clerk Maxwell,
Scottish
(13 June 1831 - 5 November 1879)

- If $\frac{\partial B_z}{\partial z} = 0$ (constant longitudinal field), then

$$\frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} = 0 \quad \frac{\partial B_x}{\partial y} - \frac{\partial B_y}{\partial x} = 0$$

- A complex function of complex variables is **analytic** if it coincides with its power series


$$f(z) = \sum_{n=1}^{\infty} C_n z^{n-1} \quad f_x(x, y) + if_y(x, y) = \sum_{n=1}^{\infty} C_n (x + iy)^{n-1} \quad (x, y) \in D$$

on a domain D !

- Note: domains are usually a painful part, we talk about it later
- A necessary and sufficient condition to be analytic is that

$$\begin{cases} \frac{\partial f_x}{\partial x} - \frac{\partial f_y}{\partial y} = 0 \\ \frac{\partial f_x}{\partial y} + \frac{\partial f_y}{\partial x} = 0 \end{cases}$$

called the Cauchy-Riemann conditions



Augustin Louis Cauchy
French
(August 21, 1789 - May 23, 1857)

DEFINITION OF FIELD HARMONICS

- If $\frac{\partial B_z}{\partial z} = 0$

Maxwell gives

$$\frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} = 0$$

$$\frac{\partial B_y}{\partial y} + \frac{\partial B_x}{\partial x} = 0$$

$$\begin{cases} \frac{\partial f_x}{\partial x} - \frac{\partial f_y}{\partial y} = 0 \\ \frac{\partial f_x}{\partial y} + \frac{\partial f_y}{\partial x} = 0 \end{cases}$$

and therefore the function $B_y + iB_x$ is analytic

$$B_y(x, y) + iB_x(x, y) = \sum_{n=1}^{\infty} C_n (x + iy)^{n-1} \quad (x, y) \in D$$

where C_n are **complex coefficients**

- Advantage: we reduce the description of a function from \mathbb{R}^2 to \mathbb{R}^2 to a (simple) series of complex coefficients
 - Attention !! **We lose something** (the function outside D)





Georg Friedrich Bernhard Riemann,
German
(November 17, 1826 - July 20, 1866)

FIELD OF A CURRENT LINE

- Field given by a current line (**Biot-Savart law**)
 - Differential form (international system)


$$d\vec{B}(\vec{r}) = \frac{\mu_0}{4\pi} \frac{\vec{I} \times d\vec{r}}{|\vec{r}|^3}$$



Félix Savart,
French
(June 30, 1791-March 16, 1841)

- Infinite current line**
 - A factor two is given by the atan integration


$$\vec{B}(\vec{r}) = \frac{\mu_0}{2\pi} \frac{\vec{I} \times \vec{r}}{|\vec{r}|^2}$$



Jean-Baptiste Biot,
French
(April 21, 1774 - February 3, 1862)

- Field in a **centre of a circular loop**, radius r

$$B = \frac{\mu_0 I}{2r}$$



FIELD OF A CURRENT LINE: COMPLEX NOTATION

- Field given by a current line (**Biot-Savart law**)

- Infinite current line

$$B_x(x, y) = -\frac{\mu_0 I}{2\pi} \frac{y - y_0}{(x - x_0)^2 + (y - y_0)^2},$$

$$B_y(x, y) = \frac{\mu_0 I}{2\pi} \frac{x - x_0}{(x - x_0)^2 + (y - y_0)^2},$$

- Complex notation

$$B_y(x, y) + iB_x(x, y) = \frac{\mu_0 I}{2\pi} \frac{(x - x_0) - i(y - y_0)}{(x - x_0)^2 + (y - y_0)^2}$$



- Using the relation

$$\frac{a - ib}{a^2 + b^2} = \frac{a - ib}{(a + ib)(a - ib)} = \frac{1}{a + ib}$$

- We obtain the compact very useful notation

$$B(z) = \frac{\mu_0 I}{2\pi(z - z_0)}$$

$$\vec{B}(\vec{r}) = \frac{\mu_0}{2\pi} \frac{\vec{I} \times \vec{r}}{|\vec{r}|^2}$$



Félix Savart,
French
(June 30, 1791-March 16, 1841)



Jean-Baptiste Biot,
French
(April 21, 1774 - February 3, 1862)

- Field given by a current line (**Biot-Savart law**)

$$B(z) = B_y(z) + iB_x(z)$$



$$B(z) = \frac{I\mu_0}{2\pi(z - z_0)} = -\frac{I\mu_0}{2\pi z_0} \frac{1}{1 - \frac{z}{z_0}}$$

using

$$\frac{1}{1-t} = 1 + t + t^2 + t^3 + \dots = \sum_{n=1}^{\infty} t^{n-1} \quad |t| < 1 !!!$$

and defining a R_{ref} as a scale length we get

$$B(z) = -\frac{I\mu_0}{2\pi z_0} \sum_{n=1}^{\infty} \left(\frac{z}{z_0}\right)^{n-1} = -\frac{I\mu_0}{2\pi z_0} \sum_{n=1}^{\infty} \left(\frac{R_{ref}}{z_0}\right)^{n-1} \left(\frac{x + iy}{R_{ref}}\right)^{n-1}$$



Félix Savart,
French
(June 30, 1791-March 16, 1841)



Jean-Baptiste Biot,
French
(April 21, 1774 - February 3, 1862)

FIELD HARMONICS OF A CURRENT LINE


- Now we can compute the **multipoles of a current line at z_0**

$$B(z) = -\frac{I\mu_0}{2\pi z_0} \sum_{n=1}^{\infty} \left(\frac{z}{z_0}\right)^{n-1} = -\frac{I\mu_0}{2\pi z_0} \sum_{n=1}^{\infty} \left(\frac{R_{ref}}{z_0}\right)^{n-1} \left(\frac{x+iy}{R_{ref}}\right)^{n-1} \quad |x+iy| < |z_0|$$

$$B_y + iB_x = 10^{-4} B_1 \sum_{n=1}^{\infty} (b_n + ia_n) \left(\frac{x+iy}{R_{ref}}\right)^{n-1}$$

$$B_1 = -\frac{I\mu_0}{2\pi} \operatorname{Re}\left(\frac{1}{z_0}\right)$$

$$b_n + ia_n = -\frac{I\mu_0 10^4}{2\pi z_0 B_1} \left(\frac{R_{ref}}{z_0}\right)^{n-1}$$




2. FIELD HARMONICS OF A CURRENT LINE

- Multipoles given by a current line **decay with the order**

$$b_n + ia_n = -\frac{I\mu_0 10^4}{2\pi z_0 B_1} \left(\frac{R_{ref}}{z_0}\right)^{n-1}$$

$$\ln(|b_n + ia_n|) = \ln\left(\frac{|I|\mu_0 10^4}{2\pi R_{ref} B_1}\right) + n \ln\left(\frac{R_{ref}}{|z_0|}\right) = p + nq$$




- The slope of the **decay is the logarithm of $(R_{ref}/|z_0|)$**
 - At each order, the multipole decreases by a factor $R_{ref}/|z_0|$
 - The decay of the multipoles tells you the ratio $R_{ref}/|z_0|$, i.e. where is the coil w.r.t. the reference radius –
 - like a radar** ... one can detect assembly errors in the magnet through magnetic field shape

2. FIELD HARMONICS OF A CURRENT LINE

- Multipoles given by a current line **decay with the order**

$$b_n + ia_n = -\frac{I\mu_0 10^4}{2\pi z_0 B_1} \left(\frac{R_{ref}}{z_0}\right)^{n-1}$$

$$\ln(|b_n + ia_n|) = \ln\left(\frac{|I|\mu_0 10^4}{2\pi R_{ref} B_1}\right) + n \ln\left(\frac{R_{ref}}{|z_0|}\right) = p + nq$$



- The **semilog scale** is the natural way to plot multipoles
 - This is the point of view of Biot-Savart
- But usually **specifications are on a linear scale**
 - In general, multipoles must stay below one or a fraction of units – see later
 - This explains why **only low order multipoles**, in general, are relevant

- Maxwell equations
- The field of a current line
 - Analytic functions and Taylor series
 - Convergence domain and singularities
- The two divergences of electromagnetism
- An outlook on scales

3. VALIDITY LIMITS OF FIELD HARMONICS

- When we expand a function in a **power series we lose something**

$$\frac{1}{1-t} = 1 + t + t^2 + t^3 + \dots = \sum_{n=1}^{\infty} t^{n-1} \quad |t| < 1$$


- Example 1. In $t=1/2$, the function is

$$\frac{1}{1-1/2} = \frac{1}{1/2} = 2$$

and using the series one has

$$1 + \frac{1}{2} + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^3 + \dots = 1 + 0.5 + 0.25 + 0.125 = 1.875 + \dots$$

not bad ... with 4 terms we compute the function within 7%



3. VALIDITY LIMITS OF FIELD HARMONICS

- When we expand a function in a **power series we lose something**

$$\frac{1}{1-t} = 1 + t + t^2 + t^3 + \dots = \sum_{n=1}^{\infty} t^{n-1} \quad |t| < 1$$


- Ex. 2. In $t=1$, the function is infinite

$$\frac{1}{1-1} = \frac{1}{0} = \infty$$

and using the series one has

$$1 + 1 + (1)^2 + (1)^3 + \dots = 1 + 1 + 1 + 1 + \dots$$

which diverges ... this makes sense



3. VALIDITY LIMITS OF FIELD HARMONICS

- When we expand a function in a **power series we lose something**

$$\frac{1}{1-t} = 1 + t + t^2 + t^3 + \dots = \sum_{n=1}^{\infty} t^{n-1} \quad |t| < 1$$


- Ex. 3. In $t=-1$, the function is well defined

$$\frac{1}{1+1} = \frac{1}{2}$$

BUT using the series one has

$$1 + (-1) + (-1)^2 + (-1)^3 + \dots = 1 - 1 + 1 - 1 + \dots$$

even if the function is well defined, the series does not work: we are **outside the convergence radius**



3. VALIDITY LIMITS OF FIELD HARMONICS

- When we expand a function in a **power series we lose something**

$$\frac{1}{1-t} = 1 + t + t^2 + t^3 + \dots = \sum_{n=1}^{\infty} t^{n-1} \quad |t| < 1$$


- Ex. 3. In $t=-1$, the function is well defined

$$\frac{1}{1+1} = \frac{1}{2}$$

BUT using the series one has

$$1 + (-1) + (-1)^2 + (-1)^3 + \dots = 1 - 1 + 1 - 1 + \dots$$

even if the function is well defined, the series does not work: we are **outside the convergence radius**



3. VALIDITY LIMITS OF FIELD HARMONICS

- Now let us consider $t=-2$

$$\frac{1}{1-t} = 1 + t + t^2 + t^3 + \dots = \sum_{n=1}^{\infty} t^{n-1} \quad |t| < 1$$

$$\frac{1}{1+2} = \frac{1}{3}$$

using the series one has


$$1 + (-2) + (-2)^2 + (-2)^3 + \dots = 1 - 2 + 4 - 8 + \dots$$

therefore when **we can resum the series** (if we manage to work out the original function) to

$$1 - 2 + 4 - 8 + 16 - 32 + \dots = \frac{1}{3}$$

see more on G. H. Hardy, Divergent series

<https://archive.org/details/DivergentSeries>



- Maxwell equations
- The field of a current line
 - Analytic functions and Taylor series
 - Convergence domain and singularities
- The two divergences of electromagnetism
- An outlook on scales

- Maxwell equations present a **divergence for a point charge**
 - Stored energy is proportional to the square of electric field

$$U = \int \frac{\epsilon_0 E^2}{2} dV$$

- Integrating over a sphere larger than R

$$U = 4\pi \frac{\epsilon_0}{2} \int_R^\infty \frac{e^2}{16\pi^2 \epsilon_0^2 r^4} r^2 dr = \frac{e^2}{8\pi\epsilon_0 R}$$

- We obtain a **divergence proportional to the size of the sphere**
- The classical electron radius is defined by equating this energy to half of the rest mass energy

$$U(r_e) = \frac{mc^2}{2}$$

$$r_e = \frac{e^2}{4\pi\epsilon_0 mc^2}$$

CLASSICAL RADIUS OF ELECTRON

- One can prove that the energy associated to a charge e on the surface of a sphere whose radius is r_e is $mc^2/2$

$$U = \frac{1}{4\pi\epsilon_0} \iint_{r=r_e} \frac{dq_1 dq_2}{r_1 - r_2} = \frac{mc^2}{2}$$

- So the total energy associated to electric field in space and to the charges on the sphere surface is equal to the electron mass
- **The electron classical radius is 2.8×10^{-15} m**

$$r_e = \frac{e^2}{4\pi\epsilon_0 mc^2}$$

- The scale length of **Thompson scattering** is the classical electron radius:

$$\sigma_t = \frac{8\pi}{3} \left(\frac{e^2}{4\pi\epsilon_0 mc^2} \right)^2$$

DIVERGENCE OF A CURRENT LOOP

- A less known divergence is related to the energy stored in the **magnetic field**
- We consider a **loop of current** of radius r_L and current I
- The loop creates a dipole-like magnetic field whose energy is

$$U = \int \frac{B^2}{2\mu_0} dV$$

- Close to the current line the field is proportional to the inverse of the distance to the current line

$$B = \frac{\mu_0 I}{2\pi r}$$

DIVERGENCE OF A CURRENT LOOP

- And one can approximate the integral with the **contribution close to the current line**

$$U \approx \frac{2\pi r_L}{2\mu_0} 2\pi \int_{r_w}^R \left(\frac{\mu_0 I}{2\pi r} \right)^2 r dr = \frac{r_L \mu_0 I^2}{2} \int_{r_w}^R \frac{dr}{r} = \frac{r_L \mu_0 I^2}{2} \ln \left(\frac{R}{r_w} \right)$$

where r_w is the wire radius

- The energy **diverges with the logarithm of the radius of the wire**, so a point charge also has a divergence in the energy associated to the magnetic field, not only to the electric field
- The **inductance of a loop of current depends on the wire size**

$$U = \frac{1}{2} LI^2 \qquad L \approx r_L \mu_0 \ln \left(\frac{R}{r_w} \right)$$

- Maxwell equations
- The field of a current line
 - Analytic functions and Taylor series
 - Convergence domain and singularities
- The two divergences of electromagnetism
- An outlook on scales



A REMINDER ON BOHR RADIUS

- The classical electron radius 10^{-15} m is about **four order of magnitudes smaller than the size of atoms**
- Bohr radius can be computed from the electrostatic potential of an electron in the field created by a proton, whose energy is

$$E = \frac{p^2}{2m} + \frac{e^2}{4\pi\epsilon_0 r}$$

- Plus the quantization of the angular momentum

$$\oint p dr = nh$$

- For a circular orbit

$$pr = n \frac{h}{2\pi} = n\hbar$$

A REMINDER ON BOHR RADIUS

- Replacing the momentum using quantization, one has

$$E(r) = \frac{h^2}{8\pi^2 r^2 m} + \frac{e^2}{4\pi\epsilon_0 r} = \frac{e^2}{4\pi\epsilon_0} \frac{1}{r^2} \left(\frac{\epsilon_0 h^2}{2\pi m e^2} + r \right)$$

- And minimizing the energy one finds the Bohr radius

$$\frac{dE}{dr} = \frac{e^2}{4\pi\epsilon_0} \frac{1}{r^3} \left(-\frac{\epsilon_0 h^2}{\pi m e^2} + r \right) \qquad r_B = \frac{\epsilon_0 h^2}{\pi m e^2}$$

- The Bohr radius is 5.3×10^{-11} m

ALPHA: THE FINE STRUCTURE CONSTANT


- What is **rarely found in textbooks** is that the ratio between the Bohr radius and the classical electron radius is $\approx (137)^2$, the square of an important physical constant

$$r_B = \frac{\epsilon_0 h^2}{\pi m e^2} = \frac{4\epsilon_0^2 h^2 c^2}{e^4} \frac{e^2}{4\pi\epsilon_0 m c^2} = \left(\frac{2\epsilon_0 h c}{e^2} \right)^2 r_e$$

- The quantity between brackets is the inverse of the **fine structure constant**

$$\alpha = \frac{e^2}{2\epsilon_0 h c} \approx \frac{1}{137}$$

- It has a very misleading name, since in reality is the **coupling constant of electromagnetism** e^2 normalized through h and c – *and is the perturbative expansion parameter of QED*



ALPHA AND COMPTON LENGTH

- In between the Bohr radius and the classical electron radius one finds another very important quantity

$$\frac{r_e}{\alpha} = \frac{2\varepsilon_0 hc}{e^2} \frac{e^2}{4\pi\varepsilon_0 mc^2} = \frac{\hbar}{mc}$$

- This is called **reduced Compton wavelength**

$$r_C = \frac{\hbar}{mc}$$

- Whose size is 3.8×10^{-13} m

