

Solar Experiments

European Strategy for Future Neutrino Physics

Gioacchino Ranucci - I.N.F.N. Milano

CERN □ October, 1, 2009

1-Solar neutrinos summary

2-Perspectives for further oscillation parameter
constraints

3-R&D challenges for the future



Neutrino production in the Sun

The pp chain reaction
The CNO cycle

In our star > 99% of the energy is created in this reaction



In the Sun < 1% More important in heavier stars

There are different steps in which energy (and neutrinos) are produced



Neutrino production in the Sun

Neutrino energy spectrum as predicted by the Solar Standard Model (SSM)



John Norris Bahcall

([Dec. 30, 1934](#) – [Aug. 17, 2005](#))

${}^7\text{Be}$: 384 keV (10%)
862 keV (90%)
Pep: 1.44 MeV

Surface metallicity
composition controversy
still open: High Z vs Low Z

Solar neutrino experiments: a more than four decades long saga

Radiochemical experiments:

Homestake (Cl)

Gallex/GNO (Ga)

Sage (Ga)

Real time Cherenkov experiments

Kamiokande/Super-Kamiokande


SNO

Scintillator experiments

Borexino



Long standing discrepancy between measured and predicted fluxes: SNP
Culminated with a crystal clear proof that neutrino oscillates



SOLAR PLUS
KAMLAND (Reactor ν 's)

Neutrino oscillations !

$$\Delta m^2 = 7.59^{+0.19}_{-0.21} \times 10^{-5}$$

$$\theta_{12} = 34.4^{+1.3}_{-1.2}$$

Phys. Rev. Lett. 101:111301, 2008

MSW matter enhanced flavor conversion
LMA solution

Given these fundamental achievements what's next?

- recent and future implications
 - from Borexino ,SNO ,Super-Kamiokande
 - note: SNO no more operating
- future prospects
 - KamLAND
 - SNO+
 - LENS
 - CLEAN, MOON, XMASS (multidisciplinary projects)



Borexino@LNGS: real time detection

Scintillator:

270 t PC+PPO in a 150 μm thick nylon vessel


Nylon vessels:

Inner: 4.25 m
Outer: 5.50 m

Neutrino electron scattering

$$\nu e \rightarrow \nu e$$

Carbon steel plates



Stainless Steel Sphere:

2212 photomultipliers
1350 m^3

Design based on the principle of graded shilding


Water Tank:

γ and n shield
 μ water Č detector
208 PMTs in water
2100 m^3



20 legs



Unprecedented intrinsic ultra low background levels allow the ${}^7\text{Be}$ spectroscopy measure



$49 \pm 3_{\text{stat}} \pm 4_{\text{syst}} \text{ cpd}/100\text{tons}$ for 862 keV ${}^7\text{Be}$ solar ν
 $\Phi({}^7\text{Be}) = (5.12 \pm 0.51) \times 10^9 \text{ cm}^{-2}\text{s}^{-1}$ SSM; H.M. $(5.08 \pm 0.56) \times 10^9 \text{ cm}^{-2}\text{s}^{-1}$
L.M. $(4.55 \pm 0.5) \times 10^9 \text{ cm}^{-2}\text{s}^{-1}$




Striking first confirmation of the predicted MSW transition between vacuum and matter regimes to be improved with a future 5% measure

Other limits obtained by Borexino after 192 days of data taking - the best in the literature

- 1- Limits on pp e CNO solar fluxes; with the Luminosity constraint:

$$\Phi_{pp}(\text{Borexino data}) / \Phi_{pp}(\text{SSM}) = 1.004^{+0.008}_{-0.020}$$

$$\Phi_{CNO}(\text{Borexino data}) / \Phi_{CNO}(\text{SSM}) < 3.8 \text{ (90\% C.L.)}$$



- 2- upper limit on the neutrino magnetic moment: $5.4 \cdot 10^{-11} \mu_B$ (90% C.L.)
Gemma : $3.2 \cdot 10^{-11} \mu_B$ (90% C.L.)

- 3- day-night asymmetry:

$$ADN^\nu = 0.02 \pm 0.04_{stat}$$



The low threshold measurement of the ${}^8\text{B}$ solar neutrinos


Major background sources:

- 1) Muons;
- 2) Gammas from neutron capture;
- 3) Radon emanation from the nylon vessel;
- 4) Short lived ($t < 2$ s) cosmogenic isotopes;
- 5) Long lived ($t > 2$ s) cosmogenic isotopes (${}^{10}\text{C}$);
- 6) Bulk ${}^{232}\text{Th}$ contamination (${}^{208}\text{Tl}$);

$\text{Rate}_{>2.8\text{ MeV}} = (0.26 \pm 0.04 \text{ stat} \pm 0.02 \text{ sys}) \text{ counts/day /100 tons}$

- ${}^7\text{Be}$ and ${}^8\text{B}$ flux measured with the same detector
- Borexino ${}^8\text{B}$ flux above 5 MeV agrees with existing data
- Neutrino oscillation is confirmed by the ${}^8\text{B}$ of Borexino at 4.2 sigma

The Borexino ${}^8\text{B}$ spectrum






Sudbury Neutrino Observatory

cc






NC





- measures total ^{8}B ν flux from the Sun
- equal cross section for all active ν flavors

ES



SNO Fluxes: 3 Phases


— stat — stat + syst



Art MacDonald@Neutel 2009

$$\frac{\phi_{\text{CC}}^{\text{SNO}}}{\phi_{\text{NC}}^{\text{SNO}}} = 0.301 \pm 0.033 \text{ (total).}$$

p-value for consistency of NC/CC/ES in the salt & NCD phases + D₂O
NC(unconstr) is 32.8%



Solar + KamLAND fit results

$$\Delta m^2 = 7.59^{+0.19}_{-0.21} \times 10^{-5} \text{ eV}^2$$

$$\phi_{8B} = 4.91 \times 10^6 \text{ cm}^2 \text{ s}^{-1} (\pm \sim 7\%)$$

$$\theta_{12} = 34.4^{+1.3}_{-1.2} \text{ degrees}$$

$$\theta_{12} = 33.9^{+2.4}_{-2.2} \text{ deg (previous)}$$

Art MacDonald@Neutel 2009

The accuracy on θ_{12} and ϕ_{8B} will improve with new data analysis: SNO LETA

Recent results:


- SNO NCD results agree well with previous SNO phases. Minimal correlation with CC. Different systematics.
- New precision on θ

Future solar analysis:

- LETA (Low Energy Threshold Analysis)
- 3-neutrino analysis
- *hep* flux
- Day-night, other variations

STATUS OF THETA13


Schwetz et al NJPhys.10:113011,2008



c.f. Fogli et al, 2008 & MINOS

$$\sin^2\theta_{13} \leq \begin{cases} 0.060 \text{ (0.089)} & \text{(solar+KamLAND)} \\ 0.027 \text{ (0.058)} & \text{(CHOOZ+atm+K2K+MINOS)} \\ 0.035 \text{ (0.056)} & \text{(global data)} \end{cases}$$

New SNO analysis, soon with lower threshold, smaller uncertainties on CC/NC so $\sin^2\theta_{12}$. Hence also $\sin^2\theta_{13}$





Solar Neutrino future prospects in SK



ν_e survival probability

(at best fit parameter)



Transition from vacuum to matter osc.
Upturn is expected in ${}^8\text{B}$ spectrum.



Cern - 1 October, 2009

Giacomo Ranucci - I.N.F.N. Sez. di Milano

Aim to reduce background in SK




BG is 70% reduced as SK-I below 5.5MeV
Energy correlated systematic error is half as SK-I

Years

Sensitivity of the upturn measurement

In the case of $(\sin^2\theta, \Delta m^2) = (0.30, 7.9 \times 10^{-5})$






- First target : 2 sigma level up-tern discovery for 3 years observation. (or exclude the up-tern)
- Need to enlarge fiducial volume with low BG as large as possible
- Also the reduction of the energy correlated systematic error is important.

(1) Enlarge fiducial volume to 22.5kton with low B.G.

(2) Half energy correlated systematic error as SK-1.

The black line shows the 13.3kton ($<5.5\text{MeV}$), 22.5kton ($>5.5\text{MeV}$) fiducial volume with the same energy correlated error as SK-1.

Solar ν measurements in SK-III



- Most of the reconstruction tools & reduction criteria are retuned recently.
- The improved reconstruction of event vertex & direction leads to 10% better angular resolution compared to SK-I.
- SK-III has lower background level in central area in low-energy region.

Confusing non-zero θ_{13} with non-standard interactions in the solar neutrino sector




FIG. 1: Region of oscillation parameters allowed ($\Delta\chi^2 = 1, 4$) by KamLAND data (independent of FCI) superimposed to the solar LMA solutions obtained in the standard case $\epsilon_{e\tau}^{dV} = 0$ and for two representative non-standard cases with FCI having equal amplitude and opposite sign ($\epsilon_{e\tau}^{dV} = \pm 0.2$).

A. Palazzo, J. W. F. Valle Arxiv:0909.1535

the possibility to disentangle the different behavior at intermediate energies could perhaps become realistic in high-energy experiments with a lowered threshold.

In this respect the new ${}^8\text{B}$ data expected from **Borexino**, **Super-K-III**, and from the low energy threshold analysis (LETA) underway in the **SNO** collaboration, may play an important role.





FIG. 3: Solar ν_e survival probability (averaged over the ${}^8\text{B}$ ν production region) for three representative cases.

θ_{12}


Recent BOREXINO




Future running: BOREXINO, KAMLAND, SuperK, SAGE (Source calibration?)
Being Developed: SNO+ (Recently Funded), LENS, MOON, CLEAN, XMASS

Other Interest: (BOREXINO, SNO+)
 Measure pep (transition region),
 Solar Models (Metallicity): CNO




The transition region (1 – 5 MeV) is a sensitive place to look for sub-dominant effects:
 Non-standard Int, Mass Varying ν



Solar Neutrino in KamLAND



nuclear fusion reaction in the sun



Standard Solar Model (SSM)


J.N. Bahcall and A.M. Serenelli, Astro. Phys. J. 621, 85 (2005)

Model	pp	pep	hep	⁷ Be	⁸ B	¹³ N	¹⁵ O	¹⁷ F
BP04(Yale)	5.94	1.40	7.88	4.86	5.79	5.71	5.03	5.91
BP04(Garching)	5.94	1.41	7.88	4.84	5.74	5.70	4.98	5.87
BS04	5.94	1.40	7.86	4.88	5.87	5.62	4.90	6.01
BS05(¹⁴ N)	5.99	1.42	7.91	4.89	5.83	3.11	2.38	5.97
BS05(OP) GS98	5.99	1.42	7.93	4.84	5.69	3.07	2.33	5.84
BS05(AGS OP) AGS05	6.06	1.45	8.25	4.34	4.51	2.01	1.45	3.25
BS05(AGS,OPAL)	6.05	1.45	8.23	4.38	4.59	2.03	1.47	3.31

Test low abundance of heavy element (AGS05)

-10% -38%

S₃₄ : 2.5% S_{1,14} : 8.4%



Radioactive BG reduction (⁸⁵Kr) : 10⁻⁴~10⁻⁵ after the 2nd purification




SNO+


- \$300M of heavy water removed and returned to Atomic Energy of Canada Limited (every last drop)
- SNO detector to be filled with liquid scintillator
 - 50-100 times more light than Čerenkov
- linear alkylbenzene (LAB)
 - compatible with acrylic, undiluted
 - high light yield, long attenuation length
 - safe: high flash point, low toxicity
 - cheaper than other scintillators
- physics goals: *pep* and *CNO* solar neutrinos, geo neutrinos, reactor neutrino oscillations, supernova neutrinos, double beta decay with Nd
- Approved and funded on June 2009



LENS Detection Scheme



The Indium Low Energy Neutrino Tag



Unique:

- Specifies ν Energy
 $E_\nu = E_e + Q$
- Complete LE nu spectrum
- Lowest Q known → 114 keV
→ access to 95.5% pp nu's
- Target isotopic abundance ~96%
- Powerful delayed coinc. Tag
Can suppress bgd = $10^{11} \times$ signal

Downside:


- Bgd from ${}^{115}\text{In}$ radioactivity to (pp nu's only) → rate = $10^{11} \times$ signal

Tools:



- Time & Space coinc. → Granularity (10^7 suppression already)
- Energy Resolution—important for In betas < 500 keV; $\sum \text{Tag} = 613 \text{ keV}$
- Other analysis cuts

Technology and Bgd Control

< Towards Hi Precision fluxes >



- Hi Quality InLS
 - New Detector Design
 - Background Analysis Insights
- In decay bgd suppressed → S/N ~3 for first time



	Status
Design of Detector	Cubic Lattice Chamber
InLS: In content Light attenuation $L(1/e)$ Signal Eff	>8% >8m 900 Pe/MeV
Indium Mass(1900 pp/5y)	10 ton
Total Mass	125 ton
PMT's	13,300
Neutrino detection eff.	64%

Raghavan@Physun workshop

Expected Result: Low Energy Solar ν -Spectrum


LENS Signal

[SSM(low CNO) + LMA
xDetection Efficiency ε]

pp: $\varepsilon=64\%$; ^7Be ^others: $\varepsilon > 85\%$
→ Rate: pp 40 /y /t ln
→ 2000 pp ev. / 5y → $\pm 2.5\%$
→ Design Goal: S/N ≥ 3

Access to pp spectral
Shape for the first time

- Valid Event: Coincidences only
Not Singles—
→ Radiopurity $< 10^{-13}\text{g/g}$, e.g. C14 OK
- ALL Bgd: Only Random Coinc.
- MEASURED Live with Signal at long
delays—No uncertainty of bgd
- Signal extracted by exponential fit
with known lifetime and stripped
- Bgd only for pp's—In decay $< 500\text{ keV}$



The CLEAN Approach

Scalable technology based on detection of scintillation in liquified noble gases. No E field.
Ultraviolet scintillation light is converted to visible light with a wavelength-shifting film.

Liquid neon and liquid argon are bright scintillators (30,000 - 40,000 photons/MeV).

Do not absorb their own scintillation.


Are inexpensive (Ar: \$2k/ton, Ne: \$60k/ton).

Are easily purified underground.


Exhibit effective pulse shape discrimination.

Exchange of targets allows better characterization of radioactive backgrounds

Self-shielding



Pulse-shape discrimination



Fast component: < 10 ns

Slow component: 1.6 μ s (LAr), 15 μ s (LNe)

Discriminate based on fraction of light in
first 100 ns (Fprompt)

D. N. McKinsey and J. M. Doyle, J. Low Temp. Phys. 118, 153 (2000).


D. N. McKinsey and K. J. Coakley, Astropart. Phys. 22, 355 (2005).

M. Boulay, J. Lidgard, and A. Hime, nucl-ex/0410025

M. Boulay and A. Hime, Astropart. Phys. 25, 179 (2006).

Mini-CLEAN central detector


Single-phase LAr/LNe detector; high light detection and simplicity
Cryostat under construction; to be installed in SNOLAB in 2009





What's XMASS

Multi purpose low-background and low-energy threshold experiment with liq. Xe


- Xenon detector for Weakly Interacting **MASSive** Particles (**DM search**)
- Xenon **MASSive** detector for solar neutrino (**pp/⁷Be**)
- Xenon neutrino **MASS** detector (**$\beta\beta$ decay**)




Solar neutrino




Dark matter






Multilayer PL plates and PL fiber planes with thin ^{100}Mo source film for pp-Be7 solar ν and $\beta\beta$ with 30 meV.



- A. Low $Q=0.17$ MeV, large CC of 680 & 220 SNU for pp & ${}^7\text{Be}-\nu$.
- B. Real time studies of inverse β rays in delayed coincidence with the β decays from ${}^{100}\text{Tc}$.
- C. 9-units 0.27 ton 100m^3 , 32 units, 1 ton 300 m^3


Detector $\neq \beta\beta$ source
Select $\beta\beta$ sources
Solar ν as well



H. Ejiri, et al., PRL, 85, 2000.

H. Ejiri, J. Phys. Soc. Japan 74 2005.


H. Ejiri European Phys. J. 162 2008



Hanohano a mobile deep ocean detector

10 kiloton liquid scintillation

Deploy and retrieve from barge



100 kton-scale
scintillator
detector
concepts

Rome - 3 July, 2009

Gioacchino Ranucci - I.N.F.N. Sez. di Milano

LENA,

DETECTOR LAYOUT

Cavern

height: 115 m, diameter: 50 m
shielding from cosmic rays: ~4,000 m.w

Muon Veto

plastic scintillator panels (on top)
Water Cherenkov Detector
1,500 phototubes
100 kt of water
reduction of fast
neutron background

Steel Cylinder

height: 100 m, diameter: 30 m
70 kt of organic liquid
13,500 phototubes

Buffer

thickness: 2 m
non-scintillating organic liquid
shielding external radioactivity

Nylon Vessel

parting buffer liquid
from liquid scintillator

Target Volume

height: 100 m, diameter: 26 m
50 kt of liquid scintillator

vertical design is favourable in terms of rock pressure and buoyancy forces



Conclusions

Solar neutrino physics is a field that has accumulated epochal successes, especially over the past decade, which constitute a solid heritage and foundation for the next round of precision experiments aimed to the accurate determination of the oscillation parameters.

Nevertheless , the perspectives of additional insights in the sub-MeV solar spectrum with the current and future solar experiments, as well as the low threshold investigation of the ${}^8\text{B}$ spectrum, pave the way for new interesting outputs especially for the non zero hint of θ_{13} and for further improvement on θ_{12} .

So this mature field still is in condition to express interesting promises for the further deepening of the understanding of the neutrino mass-mixing properties

**" I tell you: one must have chaos within oneself,
to give birth to a dancing star."** ~ Friedrich
Nietzsche from [Thus Spoke Zarathustra](#)