


# WG1 VBF report

*C. Bertella, P. Govoni, Y. Haddad, A. Karlberg, B. Jäger*  
LHC Higgs XS workshop: 14-July-2017



**Imperial College  
London**



**University of  
Zurich<sup>UZH</sup>**




EBERHARD KARLS  
**UNIVERSITÄT  
TÜBINGEN**







# Introduction

- New organization of the WG1 XS subgroups: focus group on VBF
- New CMS convener: Yacine Haddad
- New egroup: lhc-higgs-vbf: please subscribe!
- A twiki page is under construction:  
<https://twiki.cern.ch/twiki/bin/view/LHCPhysics/LHCHXSWGVBFB>
- Had our first meeting on 15th of June (<https://indico.cern.ch/event/643726/>)
- Define the roadmap for the VBF group
  - Theory inputs needed by the LHC analysis
  - Focus on major studies to be released in short/middle/long term scale (up to 1 year)
  - We tried to gather a list of topics of interest, to collect interest of people
  - Additional suggestions are welcome!

# LHC analyses status




- Many new results @ 13 TeV
- More analyses ongoing with the full luminosity




# LHC analyses status

ATLAS-CONF-2015-044



Parameter	SM prediction	Best-fit value	Uncertainty	
			Stat	Syst
$\sigma_{\text{VBF}}/\sigma_{\text{ggF}}$	$0.082 \pm 0.009$	$0.11^{+0.03}_{-0.03}$	$+0.03$ $-0.02$	$+0.02$ $-0.01$

ATLAS-CONF-2017-047



Quantity	Result	Uncertainty			SM prediction
		Total	Stat.	Exp. Th.	
$\sigma_{\text{VBF}}/\sigma_{\text{ggF}}$ [10 <sup>-2</sup> ]	21.5	$+8.5$ $-6.3$	$(+7.3$ $-5.6)$	$+2.8$ $-1.7)$	$7.9^{+0.4}_{-0.6}$

# Signal model status

- Signal model by Powheg+Pythia8 and MadGraph5\_aMC@NLO
- VBF modeling uncertainty rather important for  $H \rightarrow WW$  systematic model
  - Mainly coming from matching uncertainty - estimated from generator
    - Run-2  $H \rightarrow WW$ : impact of signal strength +14/-5% ([ATLAS-CONF-2016-112](#))
      - Comparison between Powheg and MadGraph5\_aMC@NLO
- Parton Shower and Underlying Event systematics play an important role in VBF selections
  - Run-2 ATLAS  $H \rightarrow ZZ$  order of 5% ([ATLAS-CONF-2017-043](#))
    - Comparison of Pythia8 and Herwig7
  - Run-2 CMS  $H \rightarrow \gamma\gamma$  order of 7% ([HIG-16-040](#))
    - Comparison of Pythia8 and Herwig++ tunes
    - CMS is providing now eigenvalue envelopes from the experiment's Pythia tune fit

# Main theory limitation: ggH contamination

- Predictions for higher jet multiplicities extremely important for measurement of VBF production where ggH is an irreducible background
- NNLOPS will be used for Run-2 results
- ggH QCD scale estimated using Stewart-Tackmann method
  - Run-2 ATLAS  $H \rightarrow ZZ$  order of 11% ([ATLAS-CONF-2017-043](#))
  - Run-2 CMS  $H \rightarrow \gamma\gamma$  order of 29(10)% yields(migration) ([HIG-16-040](#))
  - Run-2 ATLAS  $H \rightarrow WW$ : impact of signal strength 8%(6%) for 3(2)-jets inclusive ([ATLAS-CONF-2016-112](#))

[D. Gillberg' slides](#)


Cross sections and fractional uncertainties

STXS	sig	stat	mu	res	mig01	mig12	pTH	qm_b	qm_top	Tot
VBF1	0.27	+/- 0.00	+7.9%	+7.9%	+3.9%	+16.2%	-2.5%	-2.4%	+0.1%	+20.3%
VBF2	0.36	+/- 0.00	+7.9%	+7.9%	+3.9%	+16.2%	-0.9%	-1.1%	+0.2%	+20.1%

- Run-1 approach documented in YR3: uncertainties from MCFM parametrized as a function of  $\Delta\phi(H,jj)$  — which is correlated to the third jet  $p_T$ 
  - NB: these uncertainties are from 8 TeV
- Alternative option: uncertainties from G. Luisoni ([slides](#))

# Request from the collaborations

- LHC analyses will provide results using categories defined by the Simplified Template Cross Sections method
  - Stage 0: Closest correspondence to Run 1
  - Stage 1: “Minimal” hoped-for splitting for Run 2
- Provide theory uncertainties for the "stage 1" binning of the STXS
- QCD uncertainties: How are the uncertainties correlated in  $m_{jj}$  regions ?
  - Option 1: single overall yield  $\Delta\mu$  plus  $m_{jj}$  shape/migration  $\Delta m_{jj}$
  - Option 2: uncorrelated sources  $\Delta_{\text{low}}$  and  $\Delta_{\text{high}}$  for low and high  $m_{jj}$
- Discussion is ongoing with WG2
  - [F. Tackmann' slides](#)



# Request from the collaborations

- ggF uncertainties in VBF-like regions are becoming more relevant, could there be recommendations with the latest updates?
  - Topic in between ggF and VBF WGs
- Several analyses use MVA discriminants to separate VBF from ggF, starting from all events with (at least) 2 jets. → need to assess uncertainty on both normalization and shape of such MVA discriminants
  - YR3 addresses the problem but needs to be revisited.

# Central jet veto


- Jet activity in between the two tagging jets is strongly suppressed in VBF, which is not the case in QCD induced Hjj background.
- Hence, requiring no central jet activity can help significantly reduce the Hjj background without losing too much of the VBF signal.
- When matching NLO VBF to a parton shower, various showers give significantly different central jet activity and disagree widely on the Njets distribution ([Nason, Oleari: 0911.5299](#)).
- The third jet distributions stabilise when NLO VBF Hjjj is matched with a parton shower ([Jäger, Schissler, Zeppenfeld: 1405.6950](#)).

# Central Jet Veto (preliminary study)

- Third jet in VBF described at LO in MC sample currently used by experimental collaborations
- NNLO calculation might shed some light on parton shower models
- Preliminary studies (with F. Dreyer) suggest pt-ordered showers agree well with NNLO calculation

$$\min(y_{j1}, y_{j2}) < y_{j3} < \max(y_{j1}, y_{j2})$$

$$p_{t,\text{tag}} > 25 \text{ GeV}, \quad M_{jj} > 600 \text{ GeV}, \quad \Delta y_{jj} > 4.5, \quad y_{j1}y_{j2} < 0, \quad y_j < 4.5$$




# Central Jet Veto (preliminary study)

- The NNLO corrections tend to follow the same pattern as without the CJV - lower fiducial cross sections due to softer tagging jets.
- Corrections are in general of the same size as without the CJV (5-6% for fiducial xsec and up to 10-15% in certain kinematic regions)


$$\min(y_{j1}, y_{j2}) < y_{j3} < \max(y_{j1}, y_{j2})$$

$$p_{t,\text{tag}} > 25 \text{ GeV}, \quad M_{jj} > 600 \text{ GeV}, \quad \Delta y_{jj} > 4.5, \quad y_{j1}y_{j2} < 0, \quad y_j < 4.5$$




# Background modelling

- ggH background estimated using NNLOPS (2nd jet LO)
- Recent work suggests the xsec is overestimated under VBF cuts  
[J. Andersen, T. Hapola, A. Maier, J. Smillie]
- Main uncertainty comes from background modelling
- A dedicated study of backgrounds and optimisation of VBF cuts could be valuable (cross over with ggH sub-group)




# Background modelling

- ggH background estimated using NNLOPS (2nd jet LO)
- Recent work suggests the xsec is overestimated under VBF cuts  
[\[J. Andersen, T. Hapola, A. Maier, J. Smillie\]](#)
- Main uncertainty comes from background modelling
- A dedicated study of backgrounds and optimisation of VBF cuts could be valuable (cross over with ggH sub-group)



# Higher order corrections

- Currently no way of including NNLO-QCD and NLO-EW corrections at event generation
- A MiNLO type reweighting would be ideal but is not feasible at the moment
- Instead: use the fact that NNLO-QCD corrections are rather flat and well approximated by shower except for in the jet rapidity separation
- Feasibility studies under way



# VBF WG1 roadmap: short term

- Simplified Template Cross Sections uncertainty
- Parton shower accuracy:
  - Comparisons between different merged samples and between different showering options
  - Possible new recommendations on the central value and on the uncertainties to be adopted in the analyses
- CJV systematics:
  - CJV definition used by experimental collaboration may not be trivial (third jet information as input for MVA)
  - Third jet in VBF described at LO in MC sample currently used by experimental collaborations
  - Compare third jet kinematics of VBF Hjj NLO with VBF Hjjj NLO

# VBF WG1 roadmap: medium/long term

- Higher-order corrections (NNLOPS)
- Merging of VBF +2jet NLO and VBF + 3jets NLO
- ggH background uncertainties:
  - Large contamination of theory uncertainties from ggHjj in signal
  - ggHjj signal overestimated by current generators
- High Higgs  $p_T$ :
  - First measurements of the high  $p_T$  Higgs spectrum (above 0.5 TeV) are being published (CMS HIG-17-010)
  - Dedicated study at higher orders in QCD and EW and with additional jets are needed to be compared to data



# Backup

# Parametrization of VBF uncertainties

Bin	QCD uncertainties (Option 2)					EW uncertainties		
	$\Delta_{\text{low}}$	$\Delta_{\text{high}}$	$\Delta_{\mu}^{\text{VH}}$	$\Delta_{200}$	$\Delta_{25}$	$\Delta_{\text{Sud}}$	$\Delta_{\text{hard}}$	$\Delta_{W,Z,\gamma}^{\text{VH}}$
$p_T^j [0,200]$	$\approx 1$	$\approx 1$	$\approx 1$	$-1$		$y_1$	*	
VBF cuts	$\approx 0$	$\approx 1$	$\approx 0$	$-x_1$	$0$	$y_2$	*	
$p_T^{Hjj} [0, 25]$		$z$		$-x_1 z$	$+1$	$\dots$	*	
$p_T^{Hjj} [25, \infty]$		$1 - z$		$-x_1(1 - z)$	$-1$	$\dots$	*	
VH cuts	?	?	$\approx 1$	$-x_2$		$y_3$	*	$1$
Rest	$\approx 1$	$\approx 0$	?	$-x_3$		$y_4$	*	
$p_T^j [200, \infty]$	?	?	?	$+1$		$1 - y_i$	*	

# VBF NNLO Corrections

- Experiments would like to profit from computations of fully differential NNLO QCD + NLO EWK cross sections, likely via 1D reweighting of NLO samples
- Discussions ongoing to determine the appropriate variable and phase space



[YR4](#)