Winter school on Physics with Trapped Charged Particles

Report of Abstracts

Abstract ID : 41

Investigation of two-frequency Paul traps for antihydrogen production

Content

Radio-frequency (rf) Paul traps operated with multifrequency rf trapping potentials provide the ability to independently confine charged particle species with widely different charge-to-mass ratios. In particular, these traps may find use in the field of antihydrogen recombination, allowing antiproton and positron clouds to be trapped and confined in the same volume without the use of large superconducting magnets. We explore the stability regions of two-frequency Paul traps and perform numerical simulations of small samples of multispecies charged-particle mixtures of up to twelve particles that indicate the promise of these traps for antihydrogen recombination.

Summary

Primary author(s) : Dr. LEEFER, Nathan (Helmoltz-Institut Mainz; Department of Physics, University of California Berkeley); Mr. KRIMMEL, Kai (Helmholtz-Institut Mainz; QUANTUM, Institut für Physik, JGU Mainz); BERTSCHE, William Alan (University of Manchester (GB)); Prof. BUDKER, Dmitry (Helmoltz-Institut Mainz; QUANTUM, Institut für Physik, JGU Mainz; Department of Physics, University of California Berkeley; Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley); Prof. FAJANS, Joel (University of California Berkeley (US)); Prof. FOLMAN, Ron (Ben-Gurion University of the Negev); Prof. HÄFFNER, Hartmut (Department of Physics, University of California Berkeley); Prof. SCHMIDT-KALER, Ferdinand (Helmholtz-Institut Mainz; QUANTUM, Institut fur Physik, JGU Mainz)

Presenter(s) : Mr. KRIMMEL, Kai (Helmholtz-Institut Mainz; QUANTUM, Institut für Physik, JGU Mainz)

Status: SUBMITTED

Submitted by KRIMMEL, Kai on Thursday 30 November 2017