
New standard physics at the LHC

Yuval Grossman

Cornell

M. Galanti, A. Giammanco, YG, Y. Kats, E. Stamou, J. Zupan, 1505.02771

Measuring quark polarization

- It will be really cool if we can measure polarization of quarks
 - Examples include $H \rightarrow b\bar{b}$ and $\tilde{b} \rightarrow b\tilde{g}$
 - Polarization carries information about the Dirac structure of the couplings, not only their strengths
- It seems impossible since hadronization washes away the polarization of the quarks
- Think of b quark that hadronized into B meson
- Can we measure the polarization of b quarks?

Heavy baryons!

- Despite hadronization, bottom baryons partly retain the polarization of the b quark
 - Evidences observed at LEP in $Z \rightarrow b\bar{b}$
 - About 8% of the b quarks hadronized into baryons
-

- How to measure the b and c polarizations at the LHC?
- Can we calibrate the measurement on SM samples?
- Can we use it for discovering/characterizing new physics?

Theory of polarization loss

Time scales of polarization loss

Start with the known statement:

“The top quark decays before it hadronized and thus it keeps its polarization”

- Is that statement correct?
- How large are the corrections?
- What is the situation for $\Gamma_t = 30 \text{ MeV}$?

Time scales of polarization loss

Start with the known statement:

“The top quark decays before it hadronized and thus it keeps its polarization”

- Is that statement correct?
- How large are the corrections?
- What is the situation for $\Gamma_t = 30 \text{ MeV}$?

The relevant depolarization scale is

$$\frac{\Lambda_{\text{QCD}}^2}{m_t} \sim 1 \text{ MeV}$$

Recall the Hydrogen atom

- The 21 cm line measures the hyperfine splitting
- Consider a proton with spin up that “meets” an electron with spin down
 - Very fast the electron decay to the $n = 1$ state
 - Then the atom is in a superposition of a $J = 0$ and $J = 1$ state
 - The atom then oscillates between the two states, and the proton depolarizes at time scale of $1/\Delta E$
- What is the situation when a proton with spin up “meets” an electron with spin up?
- Hyperfine interactions in bound states lead to depolarization

b and c quarks

We now move to bound states of QCD, $Q\bar{q}$

- In the heavy quark limit (we use $Q = b$)
 - No polarization is retained for mesons
 - All baryons decay strongly to Λ_b
 - For baryons in the Heavy quark limit $\mathcal{P}(Q) = \mathcal{P}(\Lambda_b)$
 - There are corrections of order Λ_{QCD}/m_b
- Open questions
 - How large are these corrections?
 - Can we understand them theoretically?
 - Can we measure them?

Determining the depolarization

How much polarization is retained?

For interpreting polarization measurement, we need to know

$$r \equiv \frac{\mathcal{P}(\Lambda_b)}{\mathcal{P}(b)}$$

Bottom line:

- We can calculate it, but the result involves unknown hadronic parameters and thus we cannot predict it
- We hope that r can be measured in some decay modes
- We can show, theoretically, that r is universal (up to a known, small running)
- Thus, measuring r once, we can use the value of r to probe the b polarization in other processes

Calculation of r

Falk and Peskin, hep-ph/9308241

$$r \equiv \frac{\mathcal{P}(\Lambda_b)}{\mathcal{P}(b)}$$

- Main $1/m_b$ effect is due to $\Sigma_b - \Sigma_b^*$ oscillation
- The light DOFs in a $\Sigma_b^{(*)}$ (Λ_b) are a $J = 1$ ($J = 0$) state
- We need $x \equiv \Delta M/\Gamma_\Sigma$ with $\Delta M \equiv m(\Sigma_b^*) - m(\Sigma_b)$.
 - $\Delta M \sim \Lambda_{QCD}^2/m_b \sim 20$ MeV
 - $\Gamma(\Sigma \rightarrow \Lambda_b + \pi) \sim \# \times \Lambda_{QCD} \sim 10$ MeV
- While formally $O(1/m_b)$ effect, it is numerically $O(1)$
- $\Sigma_b - \Sigma_b^*$ oscillation washes out the b polarization

Calculations

- With some approximation we get

$$r = \frac{1 + (1 + 4w_1)A/9}{1 + A}$$

- r depends on two hadronization parameters

$$A = \frac{P(\Sigma_b)}{P(\Lambda_b)} \quad w_1 = \frac{P(|S_Z^{\text{light}}| = 1)}{P(S^{\text{light}} = 1)}$$

- No direct measurement of A yet.
 - Statistical hadronization model: $A \sim 2.6$
 - Pythia tunes (light hadron data): $A = 0.35 \pm 0.10$
- DELPHI $w_1 = -0.36 \pm 0.44$ and CLEO $w_1 = 0.71 \pm 0.13$

Determining A and w_1

If b is polarized transversely, r is different

$$r_L \approx \frac{1 + (0.23 + 0.38w_1)A}{1 + A} \quad r_T \approx \frac{1 + (0.62 - 0.19w_1)A}{1 + A}$$

- In principle we can measure both A and w_1
- Yet, it involves measuring r , so it will not help us to determine r
- The bottom line, it will be very hard to calculate r

Universality

Why we like universality

While we cannot calculate r we can measure it

- Consider a decay where, assuming the SM, we know the initial polarization of the b quark
- The point is that r is almost the same in any high energy process
- Basically, the b does not care much about the overall energy, just the fact that it is in a jet
- We can in fact hope to measure r in several processes and compare them to verify universality

Where to measure it at the LHC?

- Top decay
 - Maximal polarization; Large cross section; Clean sample
- $Z \rightarrow b\bar{b}$
 - Large polarization; Large cross section; Large QCD background ($S/B \approx 1/15$)
 - LEP data gave about 30% uncertainty in the value of r
- QCD production
 - Large cross section; Small and only transverse polarization; Good for LHCb
 - Looking for correlation between the two bs to get r_L

How to measure it?

$$b \rightarrow X \ell \nu$$

- Large rate (about 10%) and high sensitivity to θ , the angle between the polarization and the lepton direction

$$\frac{d\Gamma}{d \cos \theta} \propto (1 + \alpha \mathcal{P} \cos \theta)$$

where

$$\alpha_\ell \approx 0.26 \quad \alpha_\nu = 1$$

- B mesons only dilute the effect, no need to veto on them. Maybe we can still do better by demanding a Λ ?
- Can we use neutrinos? (probably yes)
- Can we use exclusive decay? (need to study it)

What about charm?

- Same formalism ensures polarization retention as in b
- Potentially promising decay mode: $\Lambda_c \rightarrow PK\pi$ with 6.7% BR (vs 3% for semileptonic)
- About 6% of charms end up inside baryons
- Can we do it with charms from $W \rightarrow c\bar{s}$ decays?
- Can also be done at Belle2 using $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$
- Can be done also using D^* . We do not need baryons to keep polarization information

Conclusions

Conclusions

- Measuring b and c polarizations is interesting
- We need calibrations from known SM processes
- It seems possible at the LHC
- Maybe can also be done with B^* and D^* decays
- Maybe possible also for s quarks

Y. Kats,