



Low Energy Neutrons

FLUKA Beginner's Course


Introduction

- In FLUKA we call neutrons below 20 MeV **low energy neutrons**
- Neutron interactions at higher energy are handled by FLUKA nuclear models
- Transport and interactions of neutrons with energies below 20 MeV are handled by a dedicated library

Why are low Energy Neutrons special?

- The neutron has no charge → can interact with nuclei at low energies, e.g. meV
- Neutron cross sections (σ) are complicated → cannot be calculated by models → we rely on data files

Typical neutron cross section



Evaluated Nuclear Data Files

- Evaluated nuclear data files (ENDF, JEFF, JENDL...)
 - typically provide neutron σ (cross sections) for $E < 20\text{MeV}$ for all channels
 - σ are stored as continuum + resonance parameters

Point-wise and Group-wise cross sections

- In neutron transport codes in general two approaches are used: **point-wise** ("continuous" cross sections) and **group-wise** transport
- Point-wise follows cross section precisely but it can be time and memory consuming
- Group approach is widely used in neutron transport codes because it is fast and gives good results for most application

Complex programs (NJOY, PREPRO...) convert ENDF files to **point-wise or group-wise cross sections**, including Doppler broadening etc.

Group Transport Technique

- The energy range of interest is divided in a given number of discrete intervals (“**energy groups**”). Each group is identified by a number increasing with decreasing energy. Group 1 is the group of highest energy (in the present FLUKA library, energies between 19.64 and 20 MeV)
- Elastic and inelastic reactions are not simulated as exclusive processes, but by group-to-group **transfer probabilities** (**downscattering matrix**)
- **Downscattering matrix**: if a neutron in a given group undergoes a scattering event and loses energy, it will be transferred to a group of lower energy (each of the lower energy groups having a different probability)
- If the neutron does not lose enough energy to be in another group, it will stay in the same group (**in-scattering**)
- In the thermal region neutrons can gain energy. This is taken into account by an **upscattering matrix**, containing the transfer probability to a group of higher energy

Group wise treatment

- Convert σ to energy groups like histograms, the energy width is different for each group in order to better represent resonances
- Each group i contains the <average> σ_i :

$$\langle \sigma_i \rangle = \frac{\int_{E_{i,low}}^{E_{i,high}} \sigma(E) \Phi(E) dE}{\int_{E_{i,low}}^{E_{i,high}} \Phi(E) dE}$$

where Φ is the fluence

- Advantage: fast
- Disadvantage: effects like self shielding are not accurately reproduced (see later), angular distributions replaced by a discrete number of angles (3 in FLUKA) conserving their first n moments (6 in FLUKA)


Fluence

- An assumption is needed about the neutron spectrum to be used as a weighting function for calculating the average cross section on each group

For instance it can be shown that in most cases, between 1 eV and 1 MeV:

$$\Phi(E)dE = C \frac{dE}{E} \quad \left(\frac{1}{E} \text{ spectrum}\right)$$

- In any case, the error is small if the group width is small



Angular distribution [1/2]

- The probability distribution of the scattering angle for each group-to-group transfer is represented by a Legendre polynomial expansion truncated at the (N+1)th term:

$$\sigma_s(g \rightarrow g', \mu) = \sum_{i=0}^N \frac{2i+1}{4\pi} P_i(\mu) \sigma_s^i(g \rightarrow g')$$

$\mu = \bar{\Omega} \cdot \bar{\Omega}' =$ scattering angle

where N = chosen Legendre order of anisotropy

Angular distribution [2/2]

- In FLUKA, $N=5$
- The scattering angular probabilities are obtained by a discretization of a P5 Legendre polynomial expansion of the actual scattering angular distribution which **preserves its first 6 moments**.
- Result of this P5 expansion is a set of **3 discrete polar angle cosines and 3 corresponding probabilities**, i.e. for a given transfer $g \rightarrow g'$ only three values are possible for the **polar** angle
- The **azimuthal** angle is sampled from a uniform distribution and can have any value between 0 and 2π

Self shielding

- The group structure is necessarily coarse with respect to the **resonance structure** in many materials
- A resonance in a material present in a dilute mixture or as a small piece cannot affect much a smooth neutron flux (so-called "**infinite dilution**")
- But if an isotope exhibiting large resonances is **very pure** or is present with a **large fractional abundance**, it can act as a "**neutron sink**", causing sharp dips in the neutron spectrum corresponding to each resonance → an apparent decrease in σ
- This effect, which results in a lower reaction rate $\sigma\Phi$, is called **self-shielding** and is necessarily **lost in the process of cross section averaging** over the width of each energy group, unless a special correction is made

The FLUKA Low Energy Neutron Library [1/3]

- FLUKA uses the **multigroup** transport technique
- The **energy boundary** below which multigroup transport takes over depends in principle on the cross section library used. In the present library it is 20 MeV.
- Both fully biased and semi-analog approaches are available
- Number of groups: 260 of approximately equal logarithmic width, the actual energy limits of each group can be found in the manual (or can be printed to *.out file)
- N.B. the **group with the highest energy has the number 1**, the group with the lowest energy has number 260
- 31 thermal groups, with 30 upscattering groups
- Energy range of library: 0.01 meV - 20 MeV

The FLUKA Low Energy Neutron library [2/3]

- Based on recent versions of evaluated nuclear data files: ENDF/B-VI.8, ENDF/B-VII.0, Jendl-3.3, Jendl-3.4, Jeff-3.1,...
- About 270 isotopes/materials available
- Almost all materials available at 2 temperatures: 87K, 296K
- Some also at 4K, 120K, 430K
- **Doppler broadening** at the relevant temperature is taken into account
- The library handles also gamma generation, energy deposition by kerma factors, residual nuclei production, secondary neutrons, fission neutrons
- For some isotopes/materials: **self-shielding**, **molecular binding**, **correlated gamma generation**, **point-wise transport**
- NB: Because of the group technique the energy of a transported neutron below 20 MeV is only defined within the resolution of the groups

The FLUKA Low Energy Neutron library [3/3]

Energy weighting:

- Averaging inside each energy group according to the following weighting function. In order of increasing energy:
 - a Maxwellian at the relevant temperature in the thermal range
 - a $1/E$ spectrum in the intermediate energy range

But it is important to realize that the groups in the present library are very narrow! Therefore the energy weighting is not very important

Gamma generation

- Gamma generation from $(n,x\gamma)$ reactions is possible only for those elements for which data are available in the evaluated nuclear data files (see manual for a complete list)
- Gamma generation is performed by a multigroup scheme as well
- Number of gamma groups: 42
 - NB number of γ groups different from number of neutron groups!
- Energy range: 1keV - 50 MeV
 - NB γ energy range different from neutron energy range!
- The actual energy of the generated photon is sampled randomly in the energy interval corresponding to its gamma group
- Exception: 2.2 MeV from $H(n,\gamma)^2H$ reaction, 478 keV photon from $^{10}B(n, \alpha\gamma)$ and gamma cascades from $Cd(n,\gamma)$ and $Xe(n,\gamma)$
- Capture gammas as well as gammas from inelastic reactions like (n,n') are included
- The neutron library only creates gammas, the transport is done by the EMF module (like all other gammas in FLUKA)

Energy deposition

- Energy deposition by neutrons below 20 MeV is estimated by means of **kerma factors**
- For some materials with gamma production the kerma values of some groups (mainly at high energies) are problematic (see manual). The reason is inconsistent data in the evaluated data files. Effort was addressed to apply corrections to improve the situation but there are still some materials with problematic kermas.
- The user should check carefully the results of simulations with these materials. However, the effect should vanish in a typical simulation.

Secondary and Fission Neutrons


- Neutrons from (n,xn) reactions are taken into account implicitly by a group-dependent **non-absorption probability**, i.e. a factor by which the weight of a neutron is multiplied after a collision
- If the only possible processes are scattering and capture the non-absorption probability is smaller than 1. If also $(n,2n)$ is possible, the factor is bigger than 1
- **Fission neutrons** are treated separately by a group-dependent fission probability
- Fission neutrons are emitted isotropically with an energy sampled from a fission spectrum appropriate for the isotope and neutron energy
- The fission neutron multiplicity was obtained separately from the evaluated data files
- The fission fragments are not transported, their energy is deposited at the spot

Charged particle generation

- **Recoil protons** from **hydrogen** and **protons** from $^{14}\text{N}(n,p)$ are produced and transported explicitly (i.e. like other protons)
- That means that detailed kinematics of elastic scattering, continuous energy loss with energy straggling, delta ray production, multiple and single scattering, are all taken into account
- If **point-wise transport** has been requested, **α and ^3H fragments from neutron capture in ^6Li and ^{10}B** can also be transported explicitly
Note that for this purpose you cannot use the cross sections of natural Li and B, but you must define a compound of ^6Li and ^7Li , or one of ^{10}B and ^{11}B
- All other charged secondaries produced in low energy neutron reactions are not transported but their energy is deposited at the point of interaction using **kerma factors**

Residual nuclei production

- **Residual nuclei:** nuclei that are the result of a reaction and are at rest, e.g. ^{28}Al after a neutron capture reaction of ^{27}Al
- For all materials data are available for estimating residual nuclei production by low energy neutrons. Command **RESNUCLEi** allows to request separately residual nuclei from low energy neutrons and from high energy particles
- For Ti, Ga the residual nuclei information is based on different evaluations than the transport



*Using the
Low Energy Neutron Library*

Available Materials

- Section 10.4.1.2 of manual gives a list of available materials
- Example:

Material		Temp.	Origin	RN	Name	Identifiers			Gam
9Be	Beryllium 9	296K	ENDF/B-VIIR0	Y	BERYLLIU	4	9	296	Y
9Be	Beryllium 9	87K	ENDF/B-VIIR0	Y	BERYLLIU	4	9	87	Y
B	Natural Boron	296K	ENDF/B-VIIR0	Y	BORON	5	-2	296	Y
B	Natural Boron	87K	ENDF/B-VIIR0	Y	BORON	5	-2	87	Y
10B	Boron 10	296K	ENDF/B-VIIR0	Y	BORON-10	5	10	296	Y
10B	Boron 10	87K	ENDF/B-VIIR0	Y	BORON-10	5	10	87	Y
11B	Boron 11	296K	ENDF/B-VIIR0	Y	BORON-11	5	11	296	Y
11B	Boron 11	87K	ENDF/B-VIIR0	Y	BORON-11	5	11	87	Y
C	Free gas natural Carbon	296K	ENDF/B-VIIR0	Y	CARBON	6	-2	296	Y
C	Graphite bound nat. Carbon	296K	ENDF/B-VIIR0	Y	CARBON	6	-3	296	Y
C	Free gas natural Carbon	87K	ENDF/B-VIIR0	Y	CARBON	6	-2	87	Y
N	Natural Nitrogen	296K	ENDF/B-VIIR0	Y	NITROGEN	7	-2	296	Y
N	Natural Nitrogen	87K	ENDF/B-VIIR0	Y	NITROGEN	7	-2	87	Y
14N	Nitrogen 14	296K	ENDF/B-VIIR0	Y	NITRO-14	7	14	296	Y
14N	Nitrogen 14	87K	ENDF/B-VIIR0	Y	NITRO-14	7	14	87	Y
16O	Oxygen 16	296K	ENDF/B-VIR8	Y	OXYGEN	8	16	296	Y
16O	Oxygen 16	87K	ENDF/B-VIR8	Y	OXYGEN	8	16	87	Y
19F	Fluorine 19	296K	ENDF/B-VIR8	Y	FLUORINE	9	19	296	Y
19F	Fluorine 19	87K	ENDF/B-VIR8	Y	FLUORINE	9	19	87	Y
23Na	Sodium 23	296K	JENDL-3.3	Y	SODIUM	11	23	296	Y
23Na	Sodium 23	87K	JENDL-3.3	Y	SODIUM	11	23	87	Y
Mg	Natural Magnesium	296K	JENDL-3.3	Y	MAGNESIU	12	-2	296	Y
Mg	Natural Magnesium	87K	JENDL-3.3	Y	MAGNESIU	12	-2	87	Y
27Al	Aluminium 27	296K	ENDF/B-VIIR0	Y	ALUMINUM	13	27	296	Y
27Al	Aluminium 27 SelfShielded	296K	ENDF/B-VIIR0	Y	ALUMINUM	13	1027	296	Y

RN: residual nuclei

Gam: Gamma production

Name: name to be used
in LOW-MAT card

Identifiers: to be used in
LOW-MAT card

Using the Low Energy Neutron Library

- How to activate low energy neutron transport?
 - Implicit: giving a **DEFAULTS** cards (except with default **EM-CASCA**), or not giving a **DEFAULTS** card at all
That means: **you are using the library in almost any simulation** (unless you are using the default **EM-CASCA** or you have switched it off explicitly with a **LOW-BIAS** or **PART-THR** card)
 - Explicit: giving a **LOW-NEUT** card
 - ◆ NB you don't need it in most practical cases! (See below)
- What must the user do?
 - To set correspondence between the actual material and the material in the low neutron library (**LOW-MAT** card), if not done by default.
 - ◆ NB you don't need it in most practical cases!
 - To set transport thresholds with **PART-THR**, if defaults are not ideal for the actual problem
 - To request special features like point wise cross sections or cross section printing (**LOW-NEUT**)

Input Cards: LOW-NEUT [1/4]

This card activates low-energy neutron transport.

- WHAT(1): number of neutron groups of the library
 - Default: 260
- WHAT(2): number of gamma groups of the library
 - Default: 42
- WHAT(3): maximum energy (in GeV) of the library
 - Default: 0.02
- WHAT(5): number of thermal groups
 - Default: 31


LOW-NEUT	n-groups: 260 ▼ Print: ▼	γ -groups: 42 ▼ XS: ▼	E _{max} : 0.02 ▼
----------	-----------------------------	---------------------------------	---------------------------

- The defaults for WHAT(1) - WHAT(3) and WHAT(5) are fine. **The only reason for changing them would be if using a different library (there was one until recently, but it has been suppressed)**

→ **Don't use them!**

Input Cards: LOW-NEUT [2/4]

- WHAT(4): printing flag, neutron cross section information is written to *.out file
 - ◆ 0.0 Minimum
 - ◆ 1.0 integral cross sections, kerma factors and probabilities
 - ◆ 2.0 additionally downscattering matrices and gamma matrices
 - ◆ 3.0 additionally scattering probabilities and angles
 - ◆ 4.0 residual nuclei information
- Default: 0.0 (minimum)



Input Cards: LOW-NEUT [3/4]

- The output for **WHAT(4) = 1**
 - Group energy limits, average energies, velocities and momenta, thermal velocities, gamma group limits
 - For each material: availability of residual nuclei information (the line: "RESIDUAL NUCLEI INFORMATIONS AVAILABLE" indicates the possibility to use option **RESNUCLEi** with **WHAT(1)= 2.0**
 - for each neutron energy group in each material:
 - ◆ **SIGT** = total cross section
 - ◆ **SIGST** = "scattering" cross section: $\sigma(n,n) + 2\sigma(n,2n) + 3\sigma(n,3n)$
 - ◆ **PNUP** = upscatter probability, is 0.0 in non thermal groups
 - ◆ **PNABS** = Probability of Non-ABSORption (= scattering)
 $PNABS = SIGST/SIGT$, and can sometimes be > 1 because of (n,xn) reactions
 - ◆ **GAMGEN** = GAMMA GENeration probability = gamma production cross section divided by **SIGT** and multiplied by the average number of γ per (n, γ) reaction
 - ◆ **NU*FIS** = fission neutron production = fission cross section divided by **SIGT** and multiplied by ν , the average number of neutrons per fission
 - ◆ **EDEP** = Kerma contribution in GeV per collision
 - ◆ **PNEL, PXN, PFISS, PNGAM** = partial cross sections, expressed as probabilities (i.e., ratios to **SIGT**). In the order: non-elastic, (n,xn) , fission, (n,gamma)

Input Cards: LOW-NEUT [4/4]

- WHAT(6): $i_0 + 10 * i_1$:
 - $i_0 = 1$: available point wise cross sections used and explicit and correlated ${}^6\text{Li}(n,\gamma){}^7\text{Li}$, ${}^6\text{Li}(n,t){}^4\text{He}$, ${}^{10}\text{B}(n,\alpha\gamma){}^7\text{Li}$, ${}^{40}\text{Ar}(n,\gamma){}^{41}\text{Ar}$, ${}^x\text{Xe}(n,\gamma){}^{x+1}\text{Xe}$ and ${}^{113}\text{Cd}(n,\gamma){}^{114}\text{Cd}$ photon cascade requested
 - ◆ = 0: ignored
 - ◆ =< -1: resets to the default (point wise cross sections are not used)
 - $i_1 = 1$, fission neutron multiplicity forced to 1, with proper weight to compensate for the "wrong" multiplicity
 - ◆ = 0, ignored
 - ◆ =< -1: resets to the default (normal fission multiplicity)
 - Default = -11., unless option **DEFAULTS** is present with **SDUM = CALORIME, ICARUS, NEUTRONS** or **PRECISIO**, in which case the default is 1.0 (point wise cross sections are used when available and fission multiplicity is not forced)

The screenshot shows the LOW-NEUT input card interface. At the top, there are three dropdown menus: 'n-groups: 260', 'γ-groups: 42', and 'Emax: 0.02'. Below these are 'Print:' and 'XS:' dropdown menus. A message reads '----- PART-THR ... STDP : 2 cards hidden'. A dropdown menu is open for the 'XS:' parameter, showing the following options: 'Default', 'Pt-wise XS', 'Fission mult', 'Pt-XS+Fission', and 'reset'.

Input Cards: LOW-MAT [1/3]

- The **LOW-MAT** card sets the correspondence between **FLUKA** materials and the low energy neutron cross sections
- If a material has the same name as a name given in the list of low neutron materials, the correspondence between material and low energy neutron transport is set automatically, and a **LOW-MAT** card is not necessary. The **first material** with the right name is taken. This is always a material at room temperature.
- That means that for the predefined material **HYDROGEN** hydrogen bound in water is used, not the free gas one
- If you want to use low energy neutron transport in H_2 gas you have to do this explicitly by a **LOW-MAT** card

Input Cards: LOW-MAT [2/3]

- WHAT(1): Name of the material (single element/isotope only!)
 - ◆ In **flair** this can be chosen from a pull down menu
- WHAT(2), WHAT(3) and WHAT(4): the 3 identifiers from table 10.4.1.2 of the manual
- SDUM: name of the material from table 10.4.1.2 of the manual
- In **flair** there is only one pull down menu for all identifiers and the name
- If you want to use the predefined materials at a temperature different from 296K, it is mandatory to give a **LOW-MAT** card with the proper identifiers

Input Cards: LOW-MAT [3/3]

- Setting the correspondence between a material and low energy neutron transport cross sections:
 - First create the material with a MATERIAL card and give it a name in SDUM
 - Give a LOW-MAT card with WHAT(1) = the name you gave in the SDUM of the MATERIAL
 - Give in WHAT(2), WHAT(3) and WHAT(4) of the LOW-MAT card the numerical identifiers (table 10.4.1.2 in manual) of the material you want to use, be careful to use the one with the **right temperature**
 - Give in SDUM of the LOW-MAT card the name provided in the same table

Example: compound at 87K

- Example: sodium iodide (NaI) at 87K
 - Create a material sodium and give it some name (SODIU_87), do the same with iodine (IOD_87)
 - Give a **LOW-MAT** card for SODIU_87, choose the right cross sections identifiers (those for 87K) and name (see list in manual), do the same for iodine
 - Create a material NAI_87 by giving first a **MATERIAL** card and then a corresponding **COMPOUND** card with the right composition

MATERIAL	Name: SODIU_87	#	p: 0.97
Z: 11	Am: 22.99	A:	dE/dx: ▼
LOW-MAT	Mat: SODIU_87 ▼	LowMat: 23Na. Sodium 23, 87K ▼	
MATERIAL	Name: IOD_87	#	p: 4.93
Z: 53	Am: 126.9	A:	dE/dx: ▼
LOW-MAT	Mat: IOD_87 ▼	LowMat: 129I. Iodine 129 (2), 87K ▼	
MATERIAL	Name: NAI_87	#	p: 3.667
Z:	Am:	A:	dE/dx: ▼
COMPOUND	Name: NAI_87 ▼	Mix: Atom ▼	Elements: 1..3 ▼
f1: 1.0	M1: IOD_87 ▼	f2: 1.0	M2: SODIU_87 ▼
f3:	M3: ▼		

MATERIAL	11.	22.99	0.97		SODIU_87
LOW-MAT	SODIU_87	11.	23.	87.	SODIUM
MATERIAL	53.	126.9	4.93		IOD_87
LOW-MAT	IOD_87	53.	129.	87.	129-I
MATERIAL			3.667		NAI_87
COMPOUND	1.0	IOD_87	1.0	SODIU_87	NAI_87

Input Cards: LOW-BIAS [1/2]

This card sets an energy cut-off during low-energy neutron transport on a region by region basis and/or non-analog absorption.

However, for an overall energy cut-off, it is preferable to use PART-THR

- WHAT(1): number of the group to apply a transport cut-off, i.e. neutrons in groups the number of which \geq WHAT(1) are not transported. Remember that the group with the highest energy has the number 1.
 - Default: 0.0 (no cut-off)
 - *flair* automatically matches the group number to the upper energy boundary of each group

The screenshot shows the LOW-BIAS input card interface. The card title is "LOW-BIAS". The "Ecut:" parameter is set to "19.155 MeV" and is currently selected in a dropdown menu. The "Reg:" parameter is set to "No cut-off". The "NonAnalogue:" parameter is set to "No cut-off". The "Survival:" parameter is set to "Step:". The "to Reg:" parameter is set to "7". The "Step:" parameter is set to "0.0". The dropdown menu for "Ecut:" lists the following energy values: 20.000 MeV, 19.640 MeV, 19.155 MeV (highlighted), 18.682 MeV, 18.221 MeV, 17.771 MeV, 27.333 MeV, 16.905 MeV, and 16.487 MeV. The interface also shows a "COMPO" section and a "requests non-analogue absorption and/or low energy neutron transport on a region" section with parameters 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100.

Input Cards: LOW-BIAS [2/2]

- **Analog absorption:** a neutron does not exist any more after an absorption process
- **Non-analog absorption:** the neutron is never killed at any interaction, but lives on with a lower weight, accounting for the relative probability of absorption and scattering.
Capture gammas are created with a weight equal to that of the surviving neutron
- **WHAT(2):** Group limit for non-analog absorption (neutrons in groups \geq WHAT(2) undergo non-analog absorption)
 - Default: 230 (non-analog absorption for all thermal neutrons)
- **WHAT(3):** non-analog survival probability
 - Default : 0.85 or 0.95, **DEFAULTS**-dependent)
- **WARNING: Only experts should modify the non-analog absorption survival probability!**
- If no **LOW-BIAS** card is given, non-analog absorption depends on the **DEFAULTS** card (see manual)
- The change of weight of the particle is taken into account (see lecture about biasing)

Transport cut-offs

- **Transport cut off**: a particle is not transported if its energy is lower than a cut off energy
- Transport cut offs for neutrons can be set to save CPU time
- Use cut offs with care, you could miss important effects like activation, dose, secondary particles,...
- For activation, thermal neutrons are very important. If you are interested in activation never cut off low energy neutrons!
- To set a general transport cut off for neutrons, please give the energy of the cut off in the **PART-THR** card, no matter if high or low energy neutrons. That was different in previous versions: the card **LOW-BIAS** was needed for low energy neutrons. **LOW-BIAS** is still needed if one wants set a cut-off region by region
- If there is no interest in low-energy neutron transport, but that feature is implicit in the **DEFAULTS** option chosen, it is suggested to request **LOW-NEUT**, and to use **PART-THRes** with an energy cutoff **WHAT(1) = 0.020**

Self-shielding [1/3]

- Self-shielded materials in FLUKA:

- ^{27}Al at 296K, 87K, 4K, 430K
- natA , ^{40}Ar at 296K, 87K
- natFe at 296K, 87K, 4K, 430K
- natCu at 296K, 87K, 4K, 430K
- ^{181}Ta at 296K, 87K
- natW at 296K, 87K, 4K, 430K
- ^{197}Au at 296K, 87K
- natPb at 296K, 87K
- ^{208}Pb at 296K
- ^{209}Bi at 296K, 87K
 - ◆ Special case: cast iron ($\text{natFe} + 5\% \text{C}$) at 296K, 87K, 4K, 430K (see example further on)

Self-shielding [2/3]

- When to use these materials?
 - Bulky (huge) pieces that are very pure (containing only one isotope)
- When not to use self-shielded materials?
 - "small" iron, copper, lead, aluminum pieces
 - Thin gold foils (but a self-shielded 100 μ m Au foil is available)
 - Diluted materials
- How to use self-shielded materials?
 - Define your material with a **MATERIAL** card
 - Give additionally a **LOW-MAT** card and give the proper identifiers in WHAT(2)-WHAT(4) and SDUM
 - If you have to use self-shielded and non self-shielded materials of the same element you need to define 2 different materials
 - Attention: predefined materials like iron, copper and lead are not self-shielded, you have to give a **LOW-MAT** card to use them self-shielded

Self-shielding [3/3]

- **Cast iron** is iron with a significant amount of carbon
- There is a self-shielded material cast iron in the low energy neutron library which is prepared to be used for creating a compound of iron and roughly 5% carbon. The amount of carbon doesn't need to be exactly 5%.
- **How to create self-shielded cast iron?**
 - Define a material iron called FeCarbSS (or any other name you like) with a **MATERIAL** card (parameters as for natural iron)
 - Insert a **LOW-MAT** card for FeCarbSS with the proper **identifiers for cast iron** in WHAT(2)-WHAT(4) and SDUM
 - Insert a **MATERIAL** card to declare a compound material called CastFe (or any other name you like)
 - Insert a **COMPOUND** card for defining CastFe as a compound of FeCarbSS and CARBON (predefined)

Artifacts of discrete angular distribution

- Artifacts can arise when a neutron is **likely to scatter only once** (thin foil, regions of low density like gases), due to the discrete angular distribution (**only 3 angles are possible for each $g \rightarrow g'$**)
- The user should be aware of such artifacts and interpret results of scattering at thin foils and gases carefully
- Because the 3 angles are different for each $g \rightarrow g'$ and the azimuthal angle is sampled from a continuous distribution, the artifact disappears when the neutrons have the possibility of scattering two or more times.
- Information about which angles and probabilities are used for each group can be obtained by setting `WHAT(4) = 3` in the **LOW-NEUT** card. The information is then written to the *.out file (see manual chapter 9: Output)

Materials with molecular binding

- Available materials with molecular bindings at 296K:
 - H (natural isotopic amount) in H_2O , CH_2
 - 1H in H_2O , CH_2
 - 2D in D_2O
 - C in graphite
- Use of these materials makes the thermal neutron calculation more realistic and can affect the energy and spatial distributions
- Example: CH_2 (polyethylene) including molecular binding
 - Create a material hydrogen and give a corresponding **LOW-MAT** card that refers to H bound in CH_2
 - Give a **COMPOUND** card that creates CH_2 as a compound of bound H and normal carbon
- For hydrogen, **H bound in water is the default**, because it is the first in the list of low energy neutron materials

Summary

- Most of the simulations in FLUKA use the low energy neutrons, implicitly via the **DEFAULTS** card
- Low neutron transport in a material is enabled by a **LOW-MAT** card, **only needed** if the material's name is not one of those in the neutron cross section library, or temperature, self-shielding or molecular binding are different from the default ones
- Use self-shielded materials properly for "bulky" and "pure" (in isotopic composition) materials
- Don't give a **LOW-MAT** card for compounds